

BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE- liquid bandage liquid

Premier Brands of America Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Premier Brands Liquid Bandage

Active ingredients

Benzethonium chloride 0.2%

Dyclonine hydrochloride 0.75%

Purpose

First aid antiseptic

Topical analgesic

Uses

- first aid to prevent bacteria contamination in minor cuts and scrapes

Warnings

For external use only.

Flammable:

do not use while smoking or near heat or flame.

Do not use

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor
- on infected areas or wounds that are draining
- with other first aid products such as lotions and creams
- over sutures
- on mucous membranes

Ask a doctor before use if you have

- deep puncture wounds
- deep cuts
- animal bites
- serious bleeding
- diabetes
- poor circulation
- serious burns

When using this product

- do not store at temperature above 120 °F

Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal.

Stop use and ask a doctor if

- conditions persists or gets worse
- infection occurs

Keep out of reach of children.

If swallowed, get medical help or call a Poison Control Center right away.

Directions

- Clean affected area
- apply a small amount on the area 1-3 times daily
- let dry
- a second coating may be applied for extra protection
- to remove, apply more Liquid Bandage and quickly wipe off
- finger nail polish remover may dissolve Liquid Bandage

Other information

- ***cap bottle tightly***
- store at room temperature away from heat
- may temporarily sting upon application
- do not allow to come in contact with floors, countertops, or other finished surfaces - will stain

Inactive ingredients

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

Questions

Call 1-866-964-0939

Principal Display Panel

Premier

For small Cuts & Wounds

Liquid Bandage

Topical Analgesic & Antiseptic

Prevents infection

Invisible & flexible

Easy to use

NET 0.3 FL OZ (9 mL)

USE LIQUID BANDAGE FOR:

Blisters, chapped & cracked finger tips, hangnails, paper cuts, shaving nicks & help in preventing the formation of calluses.

GREAT FOR:

Athletes, active lifestyles, outdoor sportment & musicians.

Premier®

NDC 56104-040-03

For Small Cuts & Wounds

LIQUID BANDAGE

Topical Analgesic & Antiseptic
Seals to Protect

- Prevents infection
- Invisible & flexible
- Easy to use

NET 0.3 FL OZ (9 mL)

USE LIQUID BANDAGE FOR:

Blisters, chapped & cracked finger tips, hangnails, paper cuts, shaving nicks & help in preventing the formation of calluses.

GREAT FOR:

Athletes, active lifestyles, outdoor sportsmen & musicians.

53-033PB

For Small
Cuts & Wounds

LIQUID BANDAGE

For Small Cuts & Wounds

LIQUID BANDAGE

Topical Analgesic & Antiseptic
Seals to Protect

NET 0.3 FL OZ (9 mL)

Distributed by:
©2019 Premier Brands of America Inc.
White Plains, NY 10601
www.premier-brands.com
Questions? Call 1-866-964-0939



MADE IN USA WITH U.S.
AND IMPORTED PARTS

DO NOT USE IF NECKBAND IS BROKEN

Drug Facts

Active ingredients
Benzethonium chloride 0.2%.....
Dyclonine hydrochloride 0.75%.....
Topical analgesic
First aid antiseptic

Purpose

Uses
■ first aid to prevent bacterial contamination in minor cuts, scrapes and burns
■ for the temporary relief of pain

Warnings
For external use only.
Flammable: Keep away from fire or flame.
Do not use ■ in the eyes ■ over large areas of the body
■ longer than 1 week unless directed by a doctor
Ask a doctor before use if you have ■ deep or puncture wounds ■ deep cuts
■ animal bites ■ serious burns
When using this product ■ avoid contact with eyes

Stop use and ask a doctor if ■ condition persists or gets worse ■ infection occurs
■ symptoms persist for more than 7 days or clear up and occur again within a few days
Keep out of reach of children. If swallowed, get medical help or call a Poison Control Center right away.


Directions ■ for adults and children 2 years of age and older ■ clean affected area
■ apply a small amount on the area 1 to 3 times daily and let dry ■ a second coating
may be applied for extra protection ■ children under 2 years of age: consult a doctor

Other information ■ cap bottle tightly ■ store between 20° to 25°C
(68° to 77°F) ■ may temporarily sting upon application ■ to remove, apply more
Liquid Bandage and quickly wipe off

Inactive ingredients nitrocellulose, SD alcohol 40
nitrocellulose, SD alcohol 40

Questions? Call 1-866-964-0939

Care ■ Wipe top of bottle clean after each use.



BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE

liquid bandage liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:56 104-040
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	0.2 mg in 9 mL
DYCLONINE HYDROCHLORIDE (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q30O)	DYCLONINE HYDROCHLORIDE	0.75 mg in 9 mL

Inactive Ingredients

Ingredient Name	Strength
ACETONE (UNII: 1364PS73AF)	
AMYL ACETATE (UNII: 92Q24NH7AS)	
CASTOR OIL (UNII: D5340Y2I9G)	
ETHYL ACETATE (UNII: 76845O8NMZ)	
PYROXYLIN (UNII: KYR8BR2X6O)	
ALCOHOL (UNII: 3K9958V90M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:56104-040-03	9 mL in 1 BOTTLE, WITH APPLICATOR; Type 0: Not a Combination Product	01/03/2018	
Marketing Information				
	Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
	OTC monograph not final	part333A	01/03/2018	

Labeler - Premier Brands of America Inc. (080051232)

Revised: 10/2019

Premier Brands of America Inc.