# MILK OF MAGNESIA- magnesium hydroxide liquid Geri-Care Pharmaceuticals, Corp

-----

#### 647 GC MOM MINT

#### Active ingredient (in each 15 mL tablespoonful)

Magnesium hydroxide 1200 mg

#### **Purpose**

Saline laxative

#### Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 1/2 to 6 hours

#### **Warnings**

#### Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are taking a prescription drug.

This product may interact with certain prescription drugs.

#### Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,** ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a

Poison Control Center right away.

#### **Directions**

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

#### Other information

- each 15 mL contains: magnesium 500 mg
- store at room temperature and avoid freezing
- close cap tightly after use

### **Inactive ingredients**

flavor, purified water, saccharin sodium, sodium hypochlorite

### package Label



\*Compare to the Active Ingredient

MINT 12 FL OZ (355 mL) REVGC1121 238-19212-4

of Phillips'® Milk of Magnesia

TAMPER-EVIDENT: Do not use if imprinted neckband is Drug Facts missing or broken.

#### Active ingredient (in each 15 mL) Magnesium hydroxide 1200 mg .....

Purpose Saline laxative

· relieves occasional constipation (irregularity).

usually produces bowel movement in ½ to 6 hours.

#### Warninas

Ask a doctor before use if you have • kidney disease

- · a magnesium-restricted diet · stomach pain, nausea, or vomiting
- · a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. This product may interact with certain prescription drugs.

#### Stop use and ask a doctor if

- . you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- · you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

- shake well before use
- . do not exceed the maximum recommended daily dose in a 24 hour period
- · dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- follow each dosage with a full glass (8oz) of liquid

3		
age	dose	
adults and children 12 years and older	30 mL to 60 mL	
children 6 to 11 years	15 mL to 30 mL	
children under 6 years	ask a doctor	

#### Other information

- each 15 mL contains: magnesium 500 mg
  store at room temperature and avoid freezing
- · close cap tightly after use

**Inactive ingredients** flavor, purified water, saccharin sodium sodium hypochlorite

\*This product is not manufactured or distributed by the owner of the registered trademark Phillips'® Milk of Magnesia.

Distributed by Power Health International Corp. 7919 20th Street Hollywood, Florida 33024 U.S.A.

Rev GC1121 938-19212-3

#### MILK OF MAGNESIA

magnesium hydroxide liquid

#### **Product Information**

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57896-003
--------------	----------------	--------------------	---------------

**Route of Administration** ORAL

#### **Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

Inactive Ingredients			
Ingredient Name	Strength		
WATER (UNII: 059QF0KO0R)			
SODIUM HYPOCHLORITE (UNII: DY38VHM5OD)			
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			

Product Characteristics				
Color		Score		
Shape		Size		
Flavor	MINT (mint)	Imprint Code		
Contains				

l	P	Packaging			
	#	# Item Code Package Description		Marketing Start Date	Marketing End Date
	1	NDC:57896-003- 12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/01/2021	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M007	01/01/2000		

# Labeler - Geri-Care Pharmaceuticals, Corp (611196254)

## **Registrant -** GCP Laboratories (965480861)

Establishment				
Name	Address	ID/FEI	Business Operations	
GCP Laboratories		965480861	manufacture(57896-003)	

Revised: 3/2025 Geri-Care Pharmaceuticals, Corp