

**ACETAMINOPHEN, CHLORPHENIRAMINE MALEATE, DEXTROMETHORPHAN HBR-**  
**acetaminophen, chlorpheniramine maleate, dextromethorphan hbr**  
**CVS Pharmacy, Inc**

-----  
**6044-CVS**

**Day Time Flu Relief Maximum Strength**

***Drug Facts***

***Active ingredients (in each caplet)***

Acetaminophen 500 mg

Dextromethorphan HBr 15 mg

***Purposes***

Pain reliever/Fever reducer

Cough suppressant

***Uses***

- temporarily relieves these symptoms due to a common cold or flu:
  - headache
  - minor aches and pains
  - cough due to minor throat and bronchial irritation
  - minor sore throat pain
- temporarily reduces fever

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain

drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- taking the blood thinning drug warfarin

**When using this product do not use more than directed.**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with a rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a doctor before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not use more than directed**
- **adults and children 12 years and over:** take 2 caplets every 6 hours, while symptoms persist
- do not take more than 6 caplets in 24 hours unless directed by a doctor
- **children under 12 years of age:** do not use

Age	Dose
adults and children 12 years of age and over	2 caplets every 6 hours
children under 12 years of age	do not use

**Other information**

Store in a dry place at 15° - 30°C (59° -86°F).

corn starch, crospovidone, FD&C blue #2 lake, FD&C red #40 lake, hypromellose, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, mineral oil, polydextrose, polyethylene glycol, povidone, sodium starch glycolate, stearic acid, talc, titanium dioxide.

1-800-231-4670

## Nighttime Flu Relief Maximu Strength

### ***Drug facts***

#### ***Active ingredients (in each caplet)***

Acetaminophen 500 mg

Chlorpheniramine maleate 2 mg

Dextromethorphan HBr 15 mg

#### ***Purposes***

Pain reliever/Fever reducer

Antihistamine

Cough suppressant

#### ***Uses***

- temporarily relieves these symptoms due to a common cold or flu:
- headache
- minor aches and pains
- cough due to minor throat and bronchial irritation
- minor sore throat pain
- runny nose
- temporarily reduces fever

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- liver disease

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children
- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with a rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a doctor before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

- **do not use more than directed**
- **adults and children 12 years and over:** take 2 caplets every 6 hours, while symptoms persist
- do not take more than 6 caplets in 24 hours unless directed by a doctor
- **children under 12 years of age:** do not use

Age	Dose
adults and children 12 years of age and over	2 caplets every 6 hours
children under 12 years of age	do not use

### ***Other information***

Store in a dry place at 15° – 30°C (59° – 86°F).

corn starch, crospovidone, FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, mineral oil, polyethylene glycol, povidone, sodium starch glycolate, stearic acid, titanium dioxide, triacetin.

1-800-231-4670

† These products are not manufactured or distributed by Haleon, distributor of Theraflu.

Distributed by: CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
© 2024 CVS/pharmacy  
CVS.com® 1-800-SHOP CVS

\*Maximum Strength per 6 hour dose.

TAMPER EVIDENT FEATURE: FLU RELIEF MAX STRENGTH CAPLETS ARE SEALED IN BLISTER PACKETS. USE ONLY IF THE INDIVIDUAL SEAL IS UNBROKEN.

READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE.  
KEEP CARTON FOR REFERENCE. DO NOT DISCARD.

DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN 6 HOURS AFTER THE LAST DOSE OF THE DAYTIME PRODUCT UNLESS DIRECTED BY YOUR DOCTOR.

DO NOT TAKE THE DAYTIME AND NIGHTTIME PRODUCTS AT THE SAME TIME.  
DO NOT TAKE MORE THAN 3 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.

CVS Health

Compare to the active ingredients in Theraflu Flu Relief Maximum Strength Daytime & Nighttime caplets†

Multi-Symptom

DAYTIME

MAXIMUM STRENGTH\*

Flu Relief

ACETAMINOPHEN

Pain reliever; Fever reducer

DEXTROMETHORPHAN HBr

Cough suppressant

Relieves: Fever, Body aches, Headache, Sore throat pain, Cough

20 CAPLETS

NIGHTTIME

MAXIMUM STRENGTH\*

Flu Relief

ACETAMINOPHEN

Pain reliever; Fever reducer

CHLORPHENIRAMINE MALEATE

Antihistamine

DEXTROMETHORPHAN HBr

Cough suppressant

Additional relief of: Runny nose

20 CAPLETS

40 TOTAL CAPLETS

### Flu Relief Maximum Strength Daytime Caplets (continued)

#### Drug Facts (continued)

**If pregnant or breast-feeding**, ask a doctor before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- do not use more than directed
- adults and children 12 years and over: take 2 caplets every 6 hours, while symptoms persist
- do not take more than 6 caplets in 24 hours unless directed by a doctor
- children under 12 years of age: do not use

Age	Dose
adults and children 12 years of age and over	2 caplets every 6 hours
children under 12 years of age	do not use

#### Other information

Store in a dry place at 15° – 30°C (59° – 86°F).

**Inactive ingredients** corn starch, crospovidone, FD&C blue #2 lake, FD&C red #40 lake, hypromellose, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, mineral oil, polydextrose, polyethylene glycol, povidone, sodium starch glycolate, stearic acid, talc, titanium dioxide.

### Flu Relief Maximum Strength Nighttime Caplets (continued)

#### Drug Facts (continued)

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with a rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a doctor before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- do not use more than directed
- adults and children 12 years and over: take 2 caplets every 6 hours, while symptoms persist
- do not take more than 6 caplets in 24 hours unless directed by a doctor
- children under 12 years of age: do not use

Age	Dose
adults and children 12 years of age and over	2 caplets every 6 hours
children under 12 years of age	do not use

### Flu Relief Maximum Strength Daytime Caplets (continued)

#### Drug Facts (continued)

**Questions? 1-800-231-4670**

### Flu Relief Maximum Strength Nighttime Caplets (continued)

#### Drug Facts (continued)

#### Other information

Store in a dry place at 15° – 30°C (59° – 86°F).

**Inactive ingredients** corn starch, crospovidone, FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, mineral oil, polyethylene glycol, povidone, sodium starch glycolate, stearic acid, titanium dioxide, triacetin.

**Questions? 1-800-231-4670**

†These products are not manufactured or distributed by Haleon, distributor of Theraflu.

40 TOTAL CAPLETS  
20 CAPLETS

Actual Size



Runny nose

**Additional relief of:**


ACETAMINOPHEN  
Pain reliever, fever reducer  
CHLORPHENIRAMINE MALEATE  
Antihistamine  
DEXTROMETHORPHAN HBr  
Cough suppressant  
Cough suppressant

**Flu Relief**  
**MAXIMUM STRENGTH\***

NIGHTTIME

20 CAPLETS

Actual Size



ACETAMINOPHEN  
Pain reliever, fever reducer  
DEXTROMETHORPHAN HBr  
Cough suppressant  
Cough suppressant  
**Relieves:**  
Fever, Body aches, Headache,  
Sore throat pain, Cough

**Flu Relief**  
**MAXIMUM STRENGTH\***

DAYTIME

Multi-Symptom

**CVS Health.**

Compare to the active ingredients in Theraflu Flu Relief Maximum Strength Daytime & Nighttime caplets†



100% money back  
guaranteed.  
CVS.com/returnpolicy

V-40056

Distributed by: CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
© 2024 CVS/pharmacy  
CVS.com • 1-800-SHOP CVS



#467276



R32091



DO NOT TAKE THE DAYTIME AND NIGHTTIME PRODUCTS AT THE SAME TIME.  
DO NOT TAKE MORE THAN 3 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.

DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN 6 HOURS AFTER  
THE LAST DOSE OF THE DAYTIME PRODUCT UNLESS DIRECTED BY YOUR DOCTOR.

READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE.  
KEEP CARTON FOR REFERENCE. DO NOT DISCARD.

TAMPER EVIDENT FEATURE: FLU RELIEF MAX STRENGTH CAPLETS ARE SEALED IN BLISTER  
PACKETS. USE ONLY IF THE INDIVIDUAL SEAL IS UNBROKEN.

\*Maximum Strength per 6 hour dose.

DO NOT TAKE THE DAYTIME AND NIGHTTIME PRODUCTS AT THE SAME TIME.  
DO NOT TAKE MORE THAN 3 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.

#### USE AS DIRECTED

##### Flu Relief Maximum Strength Daytime Caplets

### Drug Facts

#### Active Ingredients (In each caplet)

Acetaminophen 500 mg ..... Pain reliever/Fever reducer  
Dextromethorphan HBr 15 mg ..... Cough suppressant

#### Purposes

#### Uses

- temporarily relieves these symptoms due to a common cold or flu: • headache • minor aches and pains
- cough due to minor throat and bronchial irritation
- minor sore throat pain
- temporarily reduces fever

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
• more than 4,000 mg of acetaminophen in 24 hours  
• with other drugs containing acetaminophen  
• 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash  
If a skin reaction occurs, stop use and seek medical help right away.

##### Flu Relief Maximum Strength Daytime Caplets (continued)

### Drug Facts (continued)

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

#### Do not use

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- liver disease • a breathing problem such as emphysema or chronic bronchitis • cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

#### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

#### When using this product do not use more than directed.

#### Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with a rash or headache that lasts
- These could be signs of a serious condition.

##### Flu Relief Maximum Strength Nighttime Caplets

### Drug Facts

#### Active Ingredients (In each caplet)

Acetaminophen 500 mg ..... Pain reliever/Fever reducer  
Chlorpheniramine maleate 2 mg ..... Antihistamine  
Dextromethorphan HBr 15 mg ..... Cough suppressant

#### Purposes

#### Uses

- temporarily relieves these symptoms due to a common cold or flu: • headache • minor aches and pains
- cough due to minor throat and bronchial irritation
- minor sore throat pain • runny nose
- temporarily reduces fever

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
• more than 4,000 mg of acetaminophen in 24 hours  
• with other drugs containing acetaminophen  
• 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash  
If a skin reaction occurs, stop use and seek medical help right away.

##### Flu Relief Maximum Strength Nighttime Caplets (continued)

### Drug Facts (continued)

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

#### Do not use

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- liver disease • a breathing problem such as emphysema or chronic bronchitis • glaucoma • trouble urinating due to an enlarged prostate gland • cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

#### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

#### When using this product

- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

## ACETAMINOPHEN, CHLORPHENIRAMINE MALEATE, DEXTROMETHORPHAN HBR

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr kit

Product Information				
Product Type		HUMAN OTC DRUG	Item Code (Source)	
			NDC:51316-984	
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51316-984-03	1 in 1 CARTON; Type 1: Convenience Kit of Co-Package	07/10/2024	
Quantity of Parts				
Part #	Package Quantity		Total Product Quantity	
Part 1	2 BLISTER PACK		20 in 40	
Part 2	2 BLISTER PACK		20 in 40	
Part 1 of 2				
ACETAMINOPHEN, CHLORPHENIRAMINE MALEATE, DEXTROMETHORPHAN HBR				
acetaminophen, chlorpheniramine maleate, dextromethorphan hbr tablet, film coated				
Product Information				
Item Code (Source)		NDC:51316-451		
Route of Administration		ORAL		
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)			DEXTROMETHORPHAN HYDROBROMIDE	15 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)			ACETAMINOPHEN	500 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)			CHLORPHENIRAMINE MALEATE	2 mg
Inactive Ingredients				
Ingredient Name				Strength
CROSPVIDONE (UNII: 2S7830E561)				
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)				
HYPROMELLOSES (UNII: 3NXW29V3WO)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
STARCH, CORN (UNII: O8232NY3SJ)				



<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	C451
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51316-451-01	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/10/2024	

## Part 2 of 2

### ACETAMINOPHEN, DEXTROMETHORPHAN HBR

acetaminophen, dextromethorphan hbr tablet, film coated

### Product Information

<b>Item Code (Source)</b>	NDC:51316-983
<b>Route of Administration</b>	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

### Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FD&amp;C BLUE NO. 2 ALUMINUM LAKE</b> (UNII: 4AQJ3LG584)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>SODIUM STARCH GLYCOLATE TYPE A</b> (UNII: H8AV0SQX4D)	
<b>FD&amp;C RED NO. 40 ALUMINUM LAKE</b> (UNII: 6T47AS764T)	
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)	
<b>CROSPVIDONE</b> (UNII: 2S7830E561)	
<b>MEDIUM-CHAIN TRIGLYCERIDES</b> (UNII: C9H2L21V7U)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I3O)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	

### Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	C642
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51316-983-01	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/10/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/10/2024	

**Labeler** - CVS Pharmacy, Inc (062312574)