

BENZTROPINE MESYLATE- benztropine mesylate tablet
Marlex PHARMACEUTICAL, INC


BENZTROPINE
MESYLATE TABLETS, USP 0.5 mg, 1 mg and 2 mg

Rx only

DESCRIPTION

Benztropine mesylate is a synthetic compound containing structural features found in atropine and diphenhydramine.

It is a crystalline white powder, very soluble in water, designated as 3 α -(Diphenylmethoxy)-1 α H, 5 α H-tropane methanesulfonate, with the following structural formula:



C₂₁H₂₅NO•CH₄O₃S M.W. 403.54

Each tablet, for oral administration, contains 0.5 mg, 1 mg or 2 mg of benztropine mesylate.

Each tablet contains the following inactive ingredients: colloidal silicon dioxide, magnesium stearate, microcrystalline cellulose, and pregelatinized starch.

CLINICAL PHARMACOLOGY

Benztropine mesylate possesses both anticholinergic and antihistaminic effects, although only the former have been established as therapeutically significant in the management of parkinsonism.

In the isolated guinea pig ileum, the anticholinergic activity of this drug is about equal to that of atropine; however, when administered orally to unanesthetized cats, it is only about half as active as atropine.

In laboratory animals, its antihistaminic activity and duration of action approach those of pyrilamine maleate.

INDICATIONS AND USAGE

For use as an adjunct in the therapy of all forms of parkinsonism.

Useful also in the control of extrapyramidal disorders (except tardive dyskinesia - see PRECAUTIONS) due to neuroleptic drugs (e.g., phenothiazines).

CONTRAINDICATIONS

Hypersensitivity to benztropine mesylate tablets.

Because of its atropine-like side effects, this drug is contraindicated in pediatric patients under three years of age, and should be used with caution in older pediatric patients.

WARNINGS

Safe use in pregnancy has not been established.

Benztropine mesylate may impair mental and/or physical abilities required for performance of hazardous tasks, such as operating machinery or driving a motor vehicle.

When benztropine mesylate is given concomitantly with phenothiazines, haloperidol, or other drugs with anticholinergic or antidopaminergic activity, patients should be advised to report gastrointestinal complaints, fever or heat intolerance promptly. Paralytic ileus, hyperthermia and heat stroke, all of which have sometimes been fatal, have occurred in patients taking anticholinergic-type antiparkinsonism drugs, including benztropine mesylate, in combination with phenothiazines and/or tricyclic antidepressants.

Since benztropine mesylate contains structural features of atropine, it may produce anhidrosis. For this reason, it should be administered with caution during hot weather, especially when given concomitantly with other atropine-like drugs to the chronically ill, the alcoholic, those who have central nervous system disease, and those who do manual labor in a hot environment. Anhidrosis may occur more readily when some disturbance of sweating already exists. If there is evidence of anhidrosis, the possibility of hyperthermia should be considered. Dosage should be decreased at the discretion of the physician so that the ability to maintain body heat equilibrium by perspiration is not impaired. Severe anhidrosis and fatal hyperthermia have occurred.

PRECAUTIONS

General

Since benztropine mesylate has cumulative action, continued supervision is advisable. Patients with a tendency to tachycardia and patients with prostatic hypertrophy should be observed closely during treatment.

Dysuria may occur, but rarely becomes a problem. Urinary retention has been reported with benztropine mesylate.

The drug may cause complaints of weakness and inability to move particular muscle groups, especially in large doses. For example, if the neck has been rigid and suddenly relaxes, it may feel weak, causing some concern. In this event, dosage adjustment is required.

Mental confusion and excitement may occur with large doses, or in susceptible patients. Visual hallucinations have been reported occasionally. Furthermore, in the treatment of extrapyramidal disorders due to neuroleptic drugs (e.g., phenothiazines), in patients with mental disorders, occasionally there may be intensification of mental symptoms. In such cases, antiparkinsonian drugs can precipitate a toxic psychosis. Patients with mental disorders should be kept under careful observation, especially at the beginning of

treatment or if dosage is increased.

Tardive dyskinesia may appear in some patients on long-term therapy with phenothiazines and related agents, or may occur after therapy with these drugs has been discontinued. Antiparkinsonism agents do not alleviate the symptoms of tardive dyskinesia, and in some instances may aggravate them. Benztropine mesylate is not recommended for use in patients with tardive dyskinesia.

The physician should be aware of the possible occurrence of glaucoma. Although the drug does not appear to have any adverse effect on simple glaucoma, it probably should not be used in angle-closure glaucoma.

Drug Interactions

Antipsychotic drugs such as phenothiazines or haloperidol; tricyclic antidepressants (see WARNINGS).

Pediatric Use

Because of the atropine-like side effects, benztropine mesylate should be used with caution in pediatric patients over three years of age (see CONTRAINDICATIONS).

ADVERSE REACTIONS

The adverse reactions below, most of which are anticholinergic in nature, have been reported and within each category are listed in order of decreasing severity.

Cardiovascular Tachycardia.

Digestive Paralytic ileus, constipation, vomiting, nausea, dry mouth.

If dry mouth is so severe that there is difficulty in swallowing or speaking, or loss of appetite and weight, reduce dosage, or discontinue the drug temporarily.

Slight reduction in dosage may control nausea and still give sufficient relief of symptoms. Vomiting may be controlled by temporary discontinuation, followed by resumption at a lower dosage.

Nervous System Toxic psychosis, including confusion, disorientation, memory impairment, visual hallucinations; exacerbation of pre-existing psychotic symptoms; nervousness; depression; listlessness; numbness of fingers.

Special Senses Blurred vision, dilated pupils.

Urogenital Urinary retention, dysuria.

Metabolic/Immune or Skin Occasionally, an allergic reaction, e.g., skin rash, develops. If this can not be controlled by dosage reduction, the medication should be discontinued.

Other Heat stroke, hyperthermia, fever.

OVERDOSAGE

Manifestations

May be any of those seen in atropine poisoning or antihistamine overdosage; CNS depression, preceded or followed by stimulation; confusion; nervousness; listlessness; intensification of mental symptoms or toxic psychosis in patients with mental illness being treated with neuroleptic drugs (e.g., phenothiazines); hallucinations (especially visual); dizziness; muscle weakness; ataxia; dry mouth; mydriasis, blurred vision; palpitations; tachycardia; elevated blood pressure; nausea; vomiting; dysuria; numbness of fingers; dysphagia; allergic reactions, e.g., skin rash; headache; hot, dry, flushed skin; delirium; coma; shock; convulsions; respiratory arrest; anhidrosis; hyperthermia; glaucoma; constipation.

Treatment

Physostigmine salicylate, 1 to 2 mg, SC or IV, reportedly will reverse symptoms of anticholinergic intoxication.* A second injection may be given after 2 hours if required. Otherwise treatment is symptomatic and supportive. Induce emesis or perform gastric lavage (contraindicated in precomatose convulsive, or psychotic states). Maintain respiration. A short-acting barbiturate may be used for CNS excitement, but with caution to avoid subsequent depression; supportive care for depression (avoid convulsant stimulants such as picrotoxin, pentylenetetrazol, or bemegride); artificial respiration for severe respiratory depression; a local miotic for mydriasis and cycloplegia; ice bags or other cold applications and alcohol sponges for hyperpyrexia, a vasopressor and fluids for circulatory collapse. Darken room for photophobia.

DOSAGE AND ADMINISTRATION

Benztropine mesylate tablets should be used when patients are able to take oral medication.

Because of cumulative action, therapy should be initiated with a low dose which is increased gradually at five- or six-day intervals to the smallest amount necessary for optimal relief. Increases should be made in increments of 0.5 mg, to a maximum of 6 mg, or until optimal results are obtained without excessive adverse reactions.

Postencephalitic and Idiopathic Parkinsonism

The usual daily dose is 1 to 2 mg, with a range of 0.5 to 6 mg orally.

As with any agent used in parkinsonism, dosage must be individualized according to age and weight, and the type of parkinsonism being treated. Generally, older patients, and thin patients cannot tolerate large doses. Most patients with postencephalitic parkinsonism need fairly large doses and tolerate them well. Patients with a poor mental outlook are usually poor candidates for therapy.

In idiopathic parkinsonism, therapy may be initiated with a single daily dose of 0.5 to 1 mg at bedtime. In some patients, this will be adequate; in others 4 to 6 mg a day may be required.

In postencephalitic parkinsonism, therapy may be initiated in most patients with 2 mg a day in one or more doses. In highly sensitive patients, therapy may be initiated with 0.5 mg at bedtime, and increased as necessary.

Some patients experience greatest relief by taking the entire dose at bedtime; others react more favorably to divided doses, two to four times a day. Frequently, one dose a day is sufficient, and divided doses may be unnecessary or undesirable.

The long duration of action of this drug makes it particularly suitable for bedtime medication when its effects may last throughout the night, enabling patients to turn in bed during the night more easily, and to rise in the morning.

When benztropine mesylate is started, do not terminate therapy with other antiparkinsonian agents abruptly. If the other agents are to be reduced or discontinued, it must be done gradually. Many patients obtain greatest relief with combination therapy.

Benztrapine mesylate may be used concomitantly with carbidopa-levodopa, or with levodopa, in which case periodic dosage adjustment may be required in order to maintain optimum response.

Drug-Induced Extrapyramidal Disorders

In treating extrapyramidal disorders due to neuroleptic drugs (e.g., phenothiazines), the recommended dosage is 1 to 4 mg once or twice a day orally. Dosage must be individualized according to the need of the patient. Some patients require more than recommended; others do not need as much.

When extrapyramidal disorders develop soon after initiation of treatment with neuroleptic drugs (e.g., phenothiazines), they are likely to be transient. One to 2 mg of benztropine mesylate tablets two or three times a day usually provides relief within one or two days. After one or two weeks, the drug should be withdrawn to determine the continued need for it. If such disorders recur, benztropine mesylate can be reinstituted.

Certain drug-induced extrapyramidal disorders that develop slowly may not respond to benztropine mesylate.

HOW SUPPLIED

Benztrapine Mesylate Tablets, USP, are available as follows:

0.5 mg: Compressed tablet, white, round, flat – faced beveled edge tablets debossed “N” left of bisect “9” on one side and plain on other side, in bottles of 100 (NDC 10135-0813-01) and 1000 (NDC 10135-0813-10).

1 mg: Compressed tablet, white oval tablets debossed “N” left of bisect “10” on one side and plain on other side, in bottles of 100 (NDC 10135-0814-01) and 1000 (10135-0814-10).

2 mg: Compressed tablet, white, round, flat – faced beveled edge tablets debossed “N” left of bisect “11” on one side and plain on other side, in bottles of 100 (NDC 10135-0815-01) and 1000 (NDC 10135-0815-10).

Dispense in a well-closed container as defined in the USP.

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

*Duvoisin, R.C.; Katz, R.J.; Amer. Med. Ass. 206: 1963-1965, Nov. 25, 1968.

Manufactured for/ Distributed by:

Marlex Pharmaceuticals, Inc.

New Castle, DE 19720

PRINCIPAL DISPLAY PANEL

NDC 10135-0813-01

**Benztropine
Mesylate
Tablets, USP**

0.5 mg

100 Tablets

Rx only



Each tablet contains:
Benzotropine Mesylate, USP.....0.5 mg
USUAL ADULT DOSAGE: For parkinsonism, 1 to 2 mg daily. For drug induced extrapyramidal disorders, 1 to 4 mg once or twice a day. See accompanying insert.
KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN
Dispense in a well-closed container as defined in the USP.
Store at 20° to 25°C (68° to 77°F) excursions permitted to 15° to 30°C (59° to 86°F); [see USP Controlled Room Temperature].
Manufactured For/Distributed by:
Marlex Pharmaceuticals, Inc., New Castle, DE 19720
Rev. 03/25 SP



NDC 10135-0813-10

**Benzotropine
Mesylate
Tablets, USP**

0.5 mg

1000 Tablets

Rx only



Each tablet contains:
Benzotropine Mesylate, USP.....0.5 mg
USUAL ADULT DOSAGE: For parkinsonism, 1 to 2 mg daily. For drug induced extrapyramidal disorders, 1 to 4 mg once or twice a day. See accompanying insert.
KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN
Dispense in a well-closed container as defined in the USP.
Store at 20° to 25°C (68° to 77°F) excursions permitted to 15° to 30°C (59° to 86°F); [see USP Controlled Room Temperature].
Manufactured For/Distributed by:
Marlex Pharmaceuticals, Inc.
New Castle, DE 19720
Rev. 03/25 SP



NDC 10135-0814-01

**Benztropine
Mesylate
Tablets, USP**

1 mg

100 Tablets

Rx only



Each tablet contains:
Benzotropine Mesylate, USP.....1 mg
USUAL ADULT DOSAGE: For parkinsonism, 1 to 2 mg daily. For drug induced extrapyramidal disorders, 1 to 4 mg once or twice a day. See accompanying insert.
KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN
Dispense in a well-closed container as defined in the USP.
Store at 20° to 25°C (68° to 77°F) excursions permitted to 15° to 30°C (59° to 86°F); [see USP Controlled Room Temperature].
Manufactured For/Distributed by:
Marlex Pharmaceuticals, Inc., New Castle, DE 19720
Rev. 03/25 SP



NDC 10135-814-10

**Benzotropine
Mesylate
Tablets, USP**

1 mg

1000 Tablets

Rx only



Each tablet contains:
Benzotropine Mesylate, USP.....1 mg
USUAL ADULT DOSAGE: For parkinsonism, 1 to 2 mg daily. For drug induced extrapyramidal disorders, 1 to 4 mg once or twice a day. See accompanying insert.
KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN
Dispense in a well-closed container as defined in the USP.
Store at 20° to 25°C (68° to 77°F) excursions permitted to 15° to 30°C (59° to 86°F); [see USP Controlled Room Temperature].
Manufactured For/Distributed by:
Marlex Pharmaceuticals, Inc.
New Castle, DE 19720
Rev. 03/25 SP



NDC10135-0815-01

**Benzotropine
Mesylate
Tablets, USP**

2 mg

100 Tablets

Rx only

NDC 10135-0815-01

MARLEX
PHARMACEUTICALS, INC.

**Benztropine
Mesylate
Tablets, USP**

2 mg

Rx Only

100 TABLETS

Each tablet contains:
Benztropine Mesylate, USP.....2 mg

USUAL ADULT DOSAGE: For parkinsonism, 1 to 2 mg daily. For drug induced extrapyramidal disorders, 1 to 4 mg once or twice a day. See accompanying insert.

KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN

Dispense in a well-closed container as defined in the USP.

Store at 20° to 25°C (68° to 77°F) excursions permitted to 15° to 30°C (59° to 86°F); [see USP Controlled Room Temperature].

Manufactured For/Distributed by:
Marlex Pharmaceuticals, Inc., New Castle, DE 19720
Rev. 03/25 SP

10135-81501

2

N

3

NDC 10135-0815-10

**Benztropine
Mesylate
Tablets, USP**

2 mg

1000 Tablets

Rx only

NDC 10135-0815-10

MARLEX
PHARMACEUTICALS, INC.

**Benztropine
Mesylate
Tablets, USP**

2 mg

Rx Only

1000 TABLETS

Each tablet contains:
Benztropine Mesylate, USP.....2 mg

USUAL ADULT DOSAGE: For parkinsonism, 1 to 2 mg daily. For drug induced extrapyramidal disorders, 1 to 4 mg once or twice a day. See accompanying insert.

KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN

Dispense in a well-closed container as defined in the USP.

Store at 20° to 25°C (68° to 77°F) excursions permitted to 15° to 30°C (59° to 86°F); [see USP Controlled Room Temperature].

Manufactured For/Distributed by:
Marlex Pharmaceuticals, Inc.
New Castle, DE 19720
Rev. 03/25 SP

10135-81510

4

N

3

BENZTROPINE MESYLATE

benztropine mesylate tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:10135-813
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength

BENZTROPINE MESYLATE (UNII: WMJ8TL7510) (BENZTROPINE - UNII:1NHL2J4X8K)			BENZTROPINE MESYLATE	0.5 mg
Inactive Ingredients				
Ingredient Name				Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
STARCH, CORN (UNII: O8232NY3SJ)				
Product Characteristics				
Color	white	Score	2 pieces	
Shape	ROUND	Size	6mm	
Flavor		Imprint Code	N;9	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10135-813-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2025	
2	NDC:10135-813-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2025	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA204713		03/01/2025	

BENZTROPINE MESYLATE

benztropine mesylate tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:10135-814
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
BENZTROPINE MESYLATE (UNII: WMJ8TL7510) (BENZTROPINE - UNII:1NHL2J4X8K)		BENZTROPINE MESYLATE	1 mg

Inactive Ingredients				
Ingredient Name			Strength	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
STARCH, CORN (UNII: O8232NY3SJ)				
Product Characteristics				
Color	white	Score	2 pieces	
Shape	OVAL	Size	10mm	
Flavor		Imprint Code	N;10	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10135-814-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2025	
2	NDC:10135-814-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2025	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA204713	03/01/2025	

BENZTROPINE MESYLATE

benztropine mesylate tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:10135-815
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
BENZTROPINE MESYLATE (UNII: WMJ8TL7510) (BENZTROPINE - UNII:1NHL2J4X8K)		BENZTROPINE MESYLATE	2 mg
Inactive Ingredients			
Ingredient Name			Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			

CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
STARCH, CORN (UNII: O8232NY3SJ)				
Product Characteristics				
Color	white	Score	2 pieces	
Shape	ROUND	Size	7mm	
Flavor		Imprint Code	N;11	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10135-815-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2025	
2	NDC:10135-815-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2025	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA204713	03/01/2025	

Labeler - Marlex PHARMACEUTICAL, INC (782540215)

Registrant - SUNRISE PHARMACEUTICAL INC. (168522378)