

**LIDOCAINE- lidocaine patch**  
**PHARMACURE LLC**

-----

**Active Ingredient**

Lidocaine 4%

**Purpose**

Topical Anesthetic

**Uses**

Temporarily relieves mild to moderate aches and pains of muscles and joints associated with muscle soreness, strains, sprains, arthritis, simple backache, muscle stiffness, bruises

**Warnings**

**Do not use** on the face or rashes, on wounds or damaged skin, in the eyes, mouth, or other mucous membranes, on genitals, with a heating pad, any patch from a pouch that has been opened for 7 or more days.

**Ask a doctor** before use if you are allergic to topical products, you are taking a diuretic, you have high blood pressure, heart disease, kidney disease, or you are pregnant.

**When using this product** wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.

**Stop use and consult your physician if** rash, irritation, or itching develops, or condition worsens.

**If pregnant or breastfeeding**, ask a doctor before use while breastfeeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.

**For external use only**

**Keep out of reach of children.** If product is put in mouth, get medical help or contact a Poison Control Center right away. Package is not child resistant. Dispose of the used patches by folding sticky ends together.

**Directions**

**Adults and children 12 years and over:** apply 1 to 2 patches to the affected area of intact skin. Lidocaine patch should be removed after 12 hours of continuous use and remain off for at least 12 hours.

## **INSTRUCTIONS FOR USE**

- Determine area of patch application. If the pain area is smaller than the patch, patches may be cut into smaller sizes with scissors
- Safely discard the remaining unused pieces of cut patches where children and pets cannot reach
- Remove the transparent protective film (clear plastic barrier) before application of patches to the skin
- Apply immediately after removing the protective film Apply 1 patch to the affected area so that the patch covers all or most of the painful area
- Remove patch if irritation occurs

**Children under 12 years of age:** Consult a doctor.

## **Inactive ingredients**

dihydroxyaluminum aminoacetate, edetate disodium, glycerin, isopropyl myristate, polyacrylic acid, propyl p-hydroxybenzoate, propylene glycol, sodium polyacrylate, tartaric acid, titanium dioxide, tween 80, water

## **Other information**

Some individuals may not experience pain relief until several minutes or hours after applying the patch. Store in a cool, dry place. Protect the product from excessive moisture or sunlight. Store at 67-77°F (19-25°C).

## **Product label**

**DRUG FACTS****Active ingredients (in each patch):**

Lidocaine 4%

**Purpose:**

Topical Anesthetic

**USES:**

Temporarily relieves mild to moderate aches and pains of muscles and joints associated with • Muscle soreness • Strains • Sprains • Arthritis • Simple backache • Muscle stiffness • Bruises

**DIRECTIONS:****Adults and children 12 years and over:** Apply 1 to 2 patches to the affected area of intact skin. Lidocaine patch should be removed after 12 hours of continuous use and remain off for at least 12 hours.**INSTRUCTIONS FOR USE:**

- Determine area of patch application. If the pain area is smaller than the patch, patches may be cut into smaller sizes with scissors.
- Safely discard the remaining unused pieces of cut patches where children and pets cannot reach.
- Remove the transparent protective film (clear plastic barrier) before application of patches to the skin.
- Apply immediately after removing the protective film.
- Apply 1 patch to the affected area so that the patch covers all or most of the painful area.
- Remove patch if irritation occurs.

**Children under 12 years of age:** Consult a doctor.**WARNINGS:****Do not use:** • On the face or rashes • On wounds or damaged skin • In the eyes, mouth, or other mucous membranes • On genitals • With a heating pad • Any patch from a pouch that has been opened for 7 or more days**Ask a doctor before use if:** • You are allergic to topical products • You are taking a diuretic • You have high blood pressure, heart disease, or kidney disease • You are pregnant**When using this product:** • Wash hands after applying or removing patch • Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water**Stop use and consult your physician if:** • Rash, irritation, or itching develops • Condition worsens**If pregnant or breast feeding:** • Ask a doctor before use while breast feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.**Keep out of reach of children:** • If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches by folding sticky end together.

For external use only.

**OTHER INFORMATION:**

- Some individuals may not experience pain relief until several minutes or hours after applying the patch.
- Store in a cool, dry place.
- Protect product from excessive moisture or sunlight.
- Store at 67-77° F (19-25° C)

**INACTIVE INGREDIENTS:**

dihydroxyaluminum aminoacetate, edetate disodium, glycerin, isopropyl myristate, polyacrylic acid, propyl p-hydroxybenzoate, propylene glycol, sodium polyacrylate, tartaric acid, titanium dioxide, tween 80, water.



Manufactured for:

**PHARMACURE**For questions or comments  
email [info@pharmacure.info](mailto:info@pharmacure.info)

# LIDOCAINE 4% PATCH

EXTERNAL ANESTHETIC  
PAIN RELIEF

10 PATCHES  
NDC: 79643-005-01**LIDOCAINE**

lidocaine patch

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:79643-005
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>LIDOCAINE</b> (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	4 g in 100 g

**Inactive Ingredients**

Ingredient Name	Strength
<b>DIHYDROXYALUMINUM AMINOACETATE</b> (UNII: DO250MG0W6)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>ISOPROPYL MYRISTATE</b> (UNII: 0RE8K4LNJS)	
<b>POLYACRYLIC ACID (250000 MW)</b> (UNII: 9G2MAD7J6W)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	

<b>SODIUM POLYACRYLATE (2500000 MW)</b> (UNII: 05I15JNI2J)	
<b>TARTARIC ACID</b> (UNII: W4888I119H)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>WATER</b> (UNII: 059QF0KO0R)	

<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79643-005-01	10 in 1 BOX	11/29/2024	
1		14 g in 1 PATCH; Type 0: Not a Combination Product		

<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M017	11/29/2024		

**Labeler** - PHARMACURE LLC (055983858)

Revised: 10/2025

PHARMACURE LLC