

NICOTINE MINI- nicotine polacrilex lozenge
ATLANTIC BIOLOGICALS CORP.

Drug Facts

Active ingredient (in each lozenge)

Nicotine polacrilex 2 mg

Purpose

Stop smoking aid

Use

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding,

only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions (2 mg lozenge)

- **if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine lozenge
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

- **nicotine lozenge is a medicine and must be used a certain way to get the best results**
- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. **Do not chew or swallow lozenge.**
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- **do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.**
- It is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

Other information

- Phenylketonurics: Contains Phenylalanine 2.8 mg per lozenge
- store at 20-25°C (68-77°F)
- keep vial tightly closed and protect from light

Inactive ingredients

aspartame, calcium polycarbophil, flavor, maltodextrin, mannitol, sodium alginate, sodium bicarbonate, sodium stearyl fumarate, talc, xanthan gum

Questions or comments?

Call **1-800-645-2158**

DISTRIBUTED BY:

ATLANTIC BIOLOGICAL CORPS.

MIAMI, FL 33179


Package Label (2 mg lozenge)

17856-1239-01
NICOTINE (POLACRILEX)
MINI 2 MG LOZENGES



See package insert for indications and dosage schedule

Store at 20°-25°C (68°-77°F), MINT FLAVOR
** Keep this and all medication out of the reach
of children**



17856123901

17856-1239-01

Dosage 2 MG

NICOTINE (POLACRILEX)

Qty: 50 LOZENGES




GTIN: 00117856123910

S/N: 01840701

Exp: 03/07/23

Lot: 018407



Packaged by: Unit Dose Solutions
Morrisville, NC 27560

Distributed by: Atlantic Biologicals Corp,
Miami FL 33179

ev.08/21

Call to Reorder: 800.509.7592

NICOTINE MINI

nicotine polacrilex lozenge

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:17856-1239(NDC:0536-1239)
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)		NICOTINE	2 mg	
Inactive Ingredients				
Ingredient Name			Strength	
ASPARTAME (UNII: Z0H242BBR1)				
CALCIUM POLYCARBOPHIL (UNII: 8F049NKY49)				
MALTODEXTRIN (UNII: 7CVR7L4A2D)				
MANNITOL (UNII: 3OWL53L36A)				
SODIUM ALGINATE (UNII: C269C4G2ZQ)				
SODIUM BICARBONATE (UNII: 8MDF5V39QO)				
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)				
TALC (UNII: 7SEV7J4R1U)				
XANTHAN GUM (UNII: TTV12P4NEE)				
Product Characteristics				
Color	WHITE	Score	no score	
Shape	OVAL	Size	10mm	
Flavor	MINT	Imprint Code	129	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17856-1239-1	50 in 1 BOX, UNIT-DOSE	09/08/2022	
1	NDC:17856-1239-2	1 in 1 POUCH; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA207868	07/31/2019		

Labeler - ATLANTIC BIOLOGICALS CORP. (047437707)

Registrant - ATLANTIC BIOLOGICALS CORP. (047437707)

Establishment

Name	Address	ID/FEI	Business Operations
UNIT DOSE SOLUTIONS		360804194	repack(17856-1239)

