

**ANTI-DIARRHEAL- loperamide hcl tablet**  
**Walgreen Company**

-----  
**Walgreens 44-375**

***Active ingredient (in each caplet)***

Loperamide HCl 2 mg

***Purpose***

Anti-diarrheal

***Use***

controls symptoms of diarrhea, including Travelers' Diarrhea

***Warnings***

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

**Heart alert:** Taking more than directed can cause serious heart problems or death.

**Do not use**

if you have bloody or black stool.

**Ask a doctor before use if you have**

- a fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product,**

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

***Directions***

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

**Other information**

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store between 20°-25°C (68°-77°F)
- see end flap for expiration date and lot number

**Inactive ingredients**

corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

**Questions or comments?**

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

**Principal Display Panel**

**Walgreens**

Compare to Imodium®  
A-D active ingredient<sup>††</sup>

NDC 0363-0375-08

**Anti-Diarrheal**  
**LOPERAMIDE HCl 2 mg / ANTI-DIARRHEAL**

CAPLETS

- Controls the symptoms of diarrhea
- Easy-to-swallow caplets

**24**  
CAPLETS

ACTUAL SIZE

**NOT FOR HOUSEHOLDS WITH YOUNG CHILDREN**

**TAMPER EVIDENT: DO NOT  
USE IF PACKAGE IS  
OPENED OR IF BLISTER  
UNIT IS TORN, BROKEN  
OR SHOWS ANY SIGNS  
OF TAMPERING**

Walgreens Pharmacist Recommended

Walgreens Pharmacist Survey

††This product is not manufactured or distributed by Johnson  
& Johnson Corporation, owner of the registered trademark  
Imodium® A-D.

50844 REV0619B37508

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

**100% SATISFACTION GUARANTEED**

walgreens.com ©2017 Walgreen Co.

*EASY OPEN*

- 1. Tear off a section.**
- 2. Peel back paper at  
arrow and push  
caplet through foil.**

ORG1217-F  
REV1019

### Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

#### Active ingredient (in each caplet)

Loperamide HCl 2 mg

#### Purpose

Anti-diarrheal

**Use** controls symptoms of diarrhea, including Travelers' Diarrhea

#### Warnings

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

### Drug Facts (continued)

**Heart alert:** Taking more than directed can cause serious heart problems or death.

Do not use if you have bloody or black stool.

Ask a doctor before use if you have ■ a fever ■ mucus in the stool ■ a history of liver disease  
■ a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product, tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days  
■ you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions** ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

■ find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

#### Other information

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN  
■ store between 20°-25°C (68°-77°F) ■ see end flap for expiration date and lot number

**Inactive ingredients** corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions or comments? Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday



**EASY OPEN**  
1. Tear off a section.  
2. Peel back paper at arrow and push caplet through foil.


TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

50844 REV0619B37508

DISTRIBUTED BY: WALGREEN CO.  
200 WILLMOT RD., DEERFIELD, IL 60015  
**Walgreens**  
**100% SATISFACTION GUARANTEED**  
walgreens.com ©2017 Walgreen Co.

LOT/EXP are embossed  
NOT ink Jetted on

Walgreens Pharmacist Recommended  
This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Imodium® A-D.



ITEM 502887 W00000-0000-0  
1-1917-05592  
3

B-2201-375-08-H  
REV0619B37508

Walgreens

# Anti-Diarrheal

LOPERAMIDE HCl 2 mg / ANTI-DIARRHEAL

Walgreens


Compare to Imodium®  
A-D active ingredient!

NDC 0363-0375-08

# Anti-Diarrheal

LOPERAMIDE HCl 2 mg / ANTI-DIARRHEAL

CAPLETS



44-375

## ANTI-DIARRHEAL

loperamide hcl tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0375
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

### Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

### Product Characteristics

Color	GREEN (Light Green)	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	44;375
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0375-08	4 in 1 CARTON	05/03/2005	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0363-0375-22	8 in 1 CARTON	05/03/2005	07/10/2021
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:0363-0375-07	6 in 1 CARTON	05/03/2005	02/07/2021
3		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
4	NDC:0363-0375-02	2 in 1 CARTON	05/03/2005	
4		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
5	NDC:0363-0375-45	1 in 1 CARTON	05/03/2005	
5		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
6	NDC:0363-0375-90	5 in 1 CARTON	05/03/2005	02/07/2021
6		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
7	NDC:0363-0375-06	1 in 1 CARTON	05/03/2005	09/12/2020
7		200 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
8	NDC:0363-0375-46	1 in 1 CARTON	05/03/2005	07/10/2021
8		96 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076497	05/03/2005	

**Labeler** - Walgreen Company (008965063)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(0363-0375)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(0363-0375)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(0363-0375)

## Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

LNK International, Inc.		868734088	PACK(0363-0375)
-------------------------	--	-----------	-----------------

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(0363-0375)

Revised: 4/2020

Walgreen Company