

MEDI FIRST PLUS ALLERGY RELIEF- loratadine tablet, film coated
MEDIQUE LORADAMED- loratadine tablet, film coated
Unifirst First Aid Corporation

Medique Loradamed

Drug Facts

Active ingredient (in each tablet)

Loratadine 10 mg

Purpose

Antihistamine

Uses

Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver or kidney disease.

Your doctor should determine if you need a different dose.

When using this product do not take more than directed.

Taking more than directed may cause drowsiness.

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

Adults and children (6 years and over)

1 tablet daily, not more than 1 tablet in 24 hours.

Children under 6 years:

Ask a doctor

Consumer with liver or kidney disease:

Ask a doctor

Other information

- store between 68° and 77°F (20° and 25°C)
- protect from excessive moisture
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

corn starch*, lactose monohydrate, magnesium stearate, pregelatinized corn starch*, pregelatinized starch*, sodium starch glycolate*

* *may contain*

Questions or comments? Call 1-800-634-7680

Medique Loradamed Label

Medique®

Loradamed

Antihistamine • Laratadine 10 mg

Antistaminico • Loratadina 10 mg

24-Hour Allergy Relief

24-horas alivio a las alergias

Pull to Open

Tire Para Abrir

packet not child-resistant.

Paquete no resistente a los niños


*When taken as directed. See Drug Facts panel.

*Cuando se toma como se indica. Ver la información del medicamento.

Collect Medibucks

See inside flap for more details

50 Tablets (50 x 1)



Medi-First Plus Allergy Relief Label

Medi-First® Plus

24-Hour Allergy Relief

24-horas alivio a las alergias

Loratadine/ Loratadina 10 mg

Pull to Open

Tire Para Abrir

Antihistamine/Antistaminico

Packet is not child-resistant.

Paquete no resistente a los niños.

*When taken as directed. See Drug Facts panel.

*Cuando se toma como se indica. Ver la información del medicamento.

Compare active ingredient to:

Claritin®

Registered Trademark of Bayer Corp.



MEDI FIRST Plus
24-Hour Allergy Relief
24-horas alivio a las alergias

Loratadine/Loratadina 10 mg

Antihistamine / Antistaminico

"When taken as directed. See Drug Facts panel. Paquete no resistente a los niños. Ver la información del medicamento."

100 Tablets (200 x 1's)



MEDI FIRST Plus
24-Hour Allergy Relief
24-horas alivio a las alergias

Loratadine/Loratadina 10 mg

Antihistamine / Antistaminico

"When taken as directed. See Drug Facts panel. Paquete no resistente a los niños. Ver la información del medicamento."

9133380X
Rev. 3/16/23
Mfg. 23.44

Drug Facts

Active ingredient (in each tablet) Purpose
Loratadine 10 mg. Antihistamine

Uses
Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
■ runny nose ■ itchy, watery eyes ■ sneezing ■ itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
When using this product do not take more than directed. Taking more than directed may cause drowsiness.
Stop use and ask a doctor if:
■ an allergic reaction to this product occurs. Seek medical help right away.
If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
Adults and children: (6 years and over) 1 tablet daily, not more than 1 tablet in 24 hours.
Children under 6 years: Ask a doctor.
Consumers with liver or kidney disease: Ask a doctor.

Other information
■ use between 68° and 77°F (20° and 25°C)
■ protect from excessive moisture
■ larger evident sealed packets
■ do not use any opened or torn packets

Inactive ingredients com starch*, lactose monohydrate, magnesium stearate, pregelatinized com starch*, pregelatinized starch*, sodium starch glycolate*
*may contain

Questions or comments? Call 1-800-634-7680

Información del Medicamento

Ingrediente activo (en cada tableta) Propiedad
Loratadina 10 mg. Antistaminico

Usos
Para el alivio temporal de los siguientes síntomas asociados con la fiebre del heno u otras alergias del sistema respiratorio superior:
■ resaca de la nariz ■ ojos con comación o llorosos ■ estornudos ■ comción de la nariz/garganta

Advertencias
No se administre si alguna vez ha presentado alguna reacción alérgica a este producto o a cualquiera de sus ingredientes.
Consulte al médico antes de consumir este producto si usted tiene alguna enfermedad del hígado o de los riñones. Su doctor deberá determinar si usted necesita una dosis diferente.
No debe exceder la dosis recomendada. Si consume más de lo que se indica pudiera causar somnolencia.
Deje de tomar este medicamento y consulte al médico si:
■ si ocurre alguna reacción alérgica a este producto.
■ Oteenga ayuda médica de inmediato.

Información del Medicamento (continuación)

Si está embarazada o en período de lactancia, consulte con un médico antes de consumir este producto.
Mantenga este y todos los medicamentos fuera del alcance de los niños. En caso de ocurrir una sobredosis, comuníquese de inmediato con un médico o con un centro de control de envenenamiento "Poison Control Center" (1-800-222-1222).

Instrucciones
Adultos y niños de 6 años y mayores Pueden consumir 1 tableta diaria, no tome más de 1 tableta en 24 horas.
Niños menores de 6 años Consultar al médico.
Las personas con enfermedad del hígado o los riñones deben: Consultar al médico.

Otra información
■ almacenamiento: consérvese a una temperatura de entre 68° y 77° (20° y 25°C)
■ evite la humedad excesiva
■ paquetes con sellado evidente
■ no use ningún paquete abierto o roto

Información del Medicamento (continuación)

Ingredientes inactivos
almidón de maíz*, lactosa monohidratada, estearato de magnesio, almidón de maíz pregelatinizado*, almidón pregelatinizado*, amonón glicosilado*
*puede contener

¿Tiene alguna pregunta o comentario? Llame al 1-800-634-7680

Retain carton for complete product information
Tamper Evident Unit Dose Packets
Empaquetado con sellado evidente en dosis unitarias

Compare active ingredient to: **Claritin®**
Registered Trademark of Bayer Corp.

"When taken as directed. See Drug Facts panel. Cuando se toma como se indica. Ver la información del medicamento."

Conservar la caja para obtener la información completa del producto
Tamper Evident Unit Dose Packets
Empaquetado con sellado evidente en dosis unitarias

3 476821 91333 5

86461111

Manufactured by Medi First Plus
Fort Worth, TX 76102 USA
1-800-634-7680
www.medifirstproducts.com

MEDI FIRST PLUS ALLERGY RELIEF

loratadine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-913
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LORATADINE (UNII: 7AJ03BO7QN) (LORATADINE - UNII: 7AJ03BO7QN)	LORATADINE	10 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q815X)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white (white)	Score	no score
Shape	ROUND (ROUND)	Size	6mm
Flavor		Imprint Code	RX;526
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date

1	NDC:47682-913-33	100 in 1 BOX	04/23/2019	
1		1 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-913-46	1 in 1 PACKET; Type 0: Not a Combination Product	04/23/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076134	04/23/2019	

MEDIQUE LORADAMED

loratadine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-203
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LORATADINE (UNII: 7AJ03BO7QN) (LORATADINE - UNII:7AJ03BO7QN)	LORATADINE	10 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white (white)	Score	no score
Shape	ROUND (ROUND)	Size	6mm
Flavor		Imprint Code	RX;526
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-203-50	50 in 1 BOX	12/30/2008	
1	NDC:47682-203-46	1 in 1 PACKET; Type 0: Not a Combination Product		

2	NDC:47682-203-12	10 in 1 BOX	10/13/2016	
2	NDC:47682-203-46	1 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-203-46	1 in 1 PACKET; Type 0: Not a Combination Product	12/30/2008	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076134	12/30/2008	

Labeler - Unifirst First Aid Corporation (832947092)

Establishment

Name	Address	ID/FEI	Business Operations
Prestige Packaging		080667761	pack(47682-203, 47682-913)

Revised: 4/2023

Unifirst First Aid Corporation