

LIDOCAINE PAIN RELIEF GEL-PATCH- lidocaine patch
Guangzhou Liuquanmiao Import & Export Trade Co., Ltd.

Liuquanmiao, LIDOCAINE PAIN RELIEF Gel-Patch, Lidocaine 4%, 15-41 Patches

Active ingredients

Lidocaine 4%

Purpose

Topical Anesthetic

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use

- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or rashes
- do not bandage tightly

Stop use ... if

- skin reactions such as redness, swelling, blistering or other discomfort occur
- symptoms persist for more than 7 days

... ask a doctor if

- skin reactions such as redness, swelling, blistering or other discomfort occur
- symptoms persist for more than 7 days

If pregnant or breast-feeding

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN

If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old:

- Clean and dry the affected area.
- Tear off the protective film and apply the exposed part of the patch to the affected area.
- Carefully remove remaining film while pressing the patch firmly on the skin.
- Remove patch from the skin after at most 8-hour application.

Children under age of 12: Consult a doctor

Inactive Ingredients

Dihydroxyaluminum Aminoacetate, Disodium Ethylenediaminetetraacetic Acid, Glycerin, Kaolin, Methylparaben, Polysorbate 80, Propylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Water

Usage

1. Remove patch from outer package
2. Partially peel back protective film
3. Place on affected area
4. Remove remaining film while pressing patch to skin

Questions or Comments

supports@sumifunofficial.com

158-12

www.sumifunofficial.com
 Instagram: @sumifunofficial
 YouTube: @sumifun
 Facebook: @sumifunofficial
 TikTok: @sumifunofficial
 Twitter: @sumifunofficial

Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch

Numbing muscle and relieve pain:



- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	



12 PATCHES
 5.5" x 3.94"
 (14cm x 10cm)

Guangzhou Luquanmao Import & Export Trade Co., Ltd.
 417, Building A, No. 8, Dong Heng San Road,
 Zhuosen, Taishan District, Guangzhou, Guangdong, China.

Made in PRC

sumifun

Drug Facts

Active Ingredients Purpose
 Lidocaine, 4% Topical Anesthetic

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use

- on wounds or damaged skin
- with a heating pad or device
- with other ointments, creams, sprays, or liniments
- if you are allergic to any ingredients of this product

When using this product

- use only as directed
- avoid contact with the eyes, mucous membranes, or nares
- do not bandage tightly

Stop use and ask a doctor if

- skin reactions such as redness, swelling, blistering or other discomfort occur
- symptoms persist for more than 7 days

If pregnant or breast-feeding,

ask a health professional before use.
KEEP OUT OF REACH OF CHILDREN.
 If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old:
 Clean and dry the affected area.
 Tear off the protective film and apply the exposed part of the patch to the affected area.
 Carefully remove remaining film while pressing the patch firmly on the skin.
 Remove patch from the skin after at most 8-hour application.
 Children under age of 12: Consult a doctor

Inactive Ingredients

Dihydroxyaluminum Amineoctanoate,
 Disodium Ethylenediaminetetraacetate Acid, Glycerin,
 Kaolin, Methylparaben, Polysorbate 80, Propylene Glycol,
 Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben,
 Sodium Polyacrylate, Titanium Dioxide, Tartaric Acid, Water

Usage



Questions or Comments: support@sumifunofficial.com

Business Cooperation: business@sumifunofficial.com

Child Resistant Packaging



www.sumifunofficial.com
 @NSI @sumifunofficial
 YouTube: @sumifun
 Facebook: @sumifunOfficial
 TikTok: @sumifunofficial
 Twitter: @sumifunOfficial

Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch

4% LIDOCAINE

Numbing muscle and relieve pain:

- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	

13 PATCHES
 5.5" x 3.94"
 (14cm x 10cm)

Guangzhou Luquanmiao Import & Export Trade Co., Ltd.
 417 Building A, No. 8, Dong Hong Sun Road,
 Zhuwan, Taishan District, Guangzhou, Guangdong, China.

Made in PRC

unijuns

Drug Facts

Active Ingredients	Purpose
Lidocaine 4%	Topical Anesthetic

Uses
 Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbow.

Warnings
 For external use only

Do not use
 - on wounds or damaged skin
 - with a heating pad or device
 - with other ointments, creams, sprays, or liniments
 - if you are allergic to any ingredients of this product

When using this product
 use only as directed
 avoid contact with the eyes, mucous membranes, or rashes
 do not bandage tightly

Stop use and ask a doctor if
 - skin reactions such as redness, swelling, blistering or other discomfort occur
 - symptoms persist for more than 7 days

If pregnant or breast-feeding, ask a health professional before use.
KEEP OUT OF REACH OF CHILDREN.
 If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions
 Adults and Children above 12 years old:
 - Clean and dry the affected area.
 - Tear off the protective film and apply the exposed part of the patch to the affected area.
 - Carefully remove remaining film while pressing the patch firmly on the skin.
 - Remove patch from the skin after at most 8-hour application.
 Children under age of 12: Consult a doctor

Inactive Ingredients
 Dihydroxyaluminum Aminoacetate, Glycerin, Kaolin, Methylparaben, Polypropylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polycrylate, Titanium Dioxide, Tartaric Acid, Water

Usage

Questions or Comments: support@sumifunofficial.com
Business Cooperation: business@sumifunofficial.com

Child Resistant Packaging

www.sumifunofficial.com
 Instagram: @sumifunofficial
 YouTube: @sumifun
 Facebook: @sumifunOfficial
 TikTok: @sumifunofficial
 Twitter: @sumifunofficial

Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch

Numbing muscle and relieve pain:

- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	



14 PATCHES
 5.5" x 3.94"
 (14cm x 10cm)

Guangzhou Luquanmao Import & Export Trade Co., Ltd
 317, Building A, No. 8, Dong Heng San Road,
 Zhushan, Tianhe District, Guangzhou, Guangdong, China

Made in PRC

sumifun



Knee



Back



Neck



Elbow



Shoulder

Drug Facts

Active Ingredients: Lidocaine 4% Purpose: Topical Anesthetic

Uses: Temporarily relieves muscle soreness and minor joint pains in the wrist, lower back, neck, hips, shoulders, elbow

Warnings: For external use only

Do not use on wounds or damaged skin with a heating pad or device with other external creams, sprays, or treatments if you are allergic to any ingredients of this product

When using this product use only as directed avoid contact with the eyes, mucous membranes, or rashes do not bandage tightly

Stop use and ask a doctor if skin reactions such as redness, swelling, blistering or other discomfort occur symptoms persist for more than 7 days

If pregnant or breast-feeding, use a health professional before use. KEEP OUT OF REACH OF CHILDREN. If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old: Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor

Inactive Ingredients

Dihydroxyethylamine Aminocaproate, Disodium Ethylenediaminetetraacetic Acid, Glycerin, Kaolin, Methylparaben, Poly sorbate 80, Propylene Glycol, Polycrylic Acid, Polyethylene Glycol 400, Propylparaben, Sodium Polycrylate, Titanium Dioxide, Tartaric Acid, Water

Usage



Questions or Comments: support@sumifunofficial.com

Business Cooperation: business@sumifunofficial.com

Child Resistant Packaging



www.sumifunofficial.com
 @sumifunofficial
 YouTube: @sumifun
 Facebook: @sumifunOfficial
 TikTok: @sumifunofficial
 Twitter: @sumifunOfficial

Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch

Numbing muscle and relieve pain:



- Odor-Free
- Maximum Strength
- Easy to apply and remove
- Single-Use Application

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	

17 PATCHES
 5.5" x 3.94"
 (14cm x 10cm)



Guangzhou Lidoqinmao Import & Export Trade Co., Ltd.
 437, Baoding A, No. 8, Dong Heng San Road,
 Zhucun, Tianhe District, Guangzhou Guangdong, China.

Made in PRC

unijuns

<p>Drug Facts</p> <p>Active ingredients Purpose Lidocaine 4% Topical Anesthetic</p> <p>Uses Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.</p> <p>Warnings For external use only</p> <p>Do not use on wounds or damaged skin with a heating pad or device with other ointments, creams, sprays, or liniments if you are allergic to any ingredients of this product</p> <p>When using this product use only as directed avoid contact with the eyes, mucous membranes, or rashes do not bandage tightly</p> <p>Stop use and ask a doctor if skin reactions such as redness, swelling, blistering or other discomfort occur symptoms persist for more than 7 days</p> <p>If pregnant or breast-feeding, ask a health professional before use. KEEP OUT OF REACH OF CHILDREN. If swallowed accidentally, get medical help or contact with Poison Control Center immediately.</p>	<p>Directions Adults and Children above 12 years old: Clean and dry the affected area. Peel off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor</p> <p>Inactive Ingredients Dihydroxyaluminum Aminoacetate, Disodium Ethylenediaminetetraacetate Acid, Glycerin, Kaolin, Methylparaben, Polyethylene 80, Propylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polycrylate, Titanium Dioxide, Tartaric Acid, Water</p> <p>Usage</p> <p>Questions or Comments: support@sumifunofficial.com Business Cooperation: business@sumifunofficial.com</p> <p>Child Resistant Packaging</p>
--	--

Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch



Numbing muscle and relieve pain:

- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	



Knee



Back



Neck



Elbow



Shoulder

19 PATCHES
5.5" x 3.54"
(14cm x 10cm)

Guangzhou Liqunmao Import & Export Trade Co., Ltd.
337, Daibang A Road, Dongfeng San Road,
Zhuchang, Tianhe District, Guangzhou, Guangdong, China.

Made in PRC

www.sumifunofficial.com
INS @sumifunofficial

YouTube @sumifun
Facebook @sumifunofficial

TikTok @sumifunofficial
Twitter @sumifunofficial

unjinms

Drug Facts

Active Ingredients Purpose
Lidocaine 4% Topical Anesthetic

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knee, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use

on wounds or damaged skin with a heating pad or device with other ointments, creams, sprays, or liniments if you are allergic to any ingredients of the product

When using this product

use only as directed avoid contact with the eyes, mucous membranes, or mouth do not bandage tightly

Stop use and ask a doctor if

skin reactions such as redness, swelling, blistering or other discomfort occur symptoms persist for more than 7 days

If pregnant or breast-feeding,

ask a health professional before use. **KEEP OUT OF REACH OF CHILDREN.** If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old:

Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor

Inactive Ingredients

Dihydroxyaluminum Aminoacetate, Dimethylammonium Acrylate, Glycerin, Kaolin, Methylparaben, Polysorbate 80, Propylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polyacrylate, Titanium Dioxide, Tartaric Acid, Water

Usage



Remove patch from outer package



Partially peel back protective film



Place on affected area



Remove remaining film while pressing patch to skin

Questions or Comments: support@sumifunofficial.com
Business Cooperations: business@sumifunofficial.com

Child Resistant Packaging



Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch




Numbing muscle and relieve pain:

- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	

22 PATCHES
5.5" x 3.94"
(14cm x 10cm)

Guangzhou Lidoan Import & Export Trade Co., Ltd.
817, Baiming Rd, No. 8, Dong Feng East Road,
Zhuzhou, Tianhe District, Guangzhou, Guangdong, China.

Made in PRC

www.sumifunofficial.com
INS: @sumifunofficial
YouTube: @sumifun
Facebook: @sumifunofficial
TikTok: @sumifunofficial
Twitter: @sumifunofficial

Sumifun®

<p>Drug Facts</p> <p>Active ingredients Lidoanaine 4% Purpose: Topical Anesthetic</p> <p>Uses Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.</p> <p>Warnings For external use only</p> <p>Do not use on wounds or damaged skin with a heating pad or device with other electrical, chemical, sprays, or treatments if you are allergic to any ingredients of this product</p> <p>When using this product use only as directed avoid contact with the eyes, mucous membranes, or rashes do not bandage tightly</p> <p>Stop use and ask a doctor if skin reactions such as redness, swelling, blistering or other discomfort occur symptoms persist for more than 7 days</p> <p>If pregnant or breast-feeding, ask a health professional before use. KEEP OUT OF REACH OF CHILDREN. If swallowed accidentally, get medical help or contact with Poison Control Center immediately.</p>	<p>Directions Adults and Children above 12 years old: Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor</p> <p>Inactive Ingredients Dihydroaluminum Ammonioacetate, Disodium Ethylenediaminetetraacetic Acid, Glycerin, Kaolin, Methylparaben, Polypropylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polyacrylate, Titanium Dioxide, Talcum Acid, Water</p> <p>Usage</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">  <p>Remove patch from outer package</p> </div> <div style="text-align: center;">  <p>Partially peel back protective film</p> </div> </div> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">  <p>Place on affected area</p> </div> <div style="text-align: center;">  <p>Remove remaining film while pressing patch to skin</p> </div> </div> <p>Questions or Comments: support@sumifunofficial.com Business Cooperation: business@sumifunofficial.com</p> <p>Child Resistant Packaging</p> 
---	--

www.sumifunofficial.com
 INS: @sumifunofficial
 YouTube: @sumifun
 Facebook: @sumifunOfficial
 Twitter: @sumifunOfficial

Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch



Numbing muscle and relieve pain:

- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	



23 PATCHES
 3.5" x 3.94"
 (14cm x 10cm)

Guangzhou Lidaqian Import & Export Trade Co., Ltd.
 117, Building A, No. 6, Dong Heng San Road,
 Zhuzhen, Tieshe District, Guangzhou Guangdong, China

Made in PRC

Sumifun

Drug Facts

Active Ingredients _____ **Purpose**
 Lidocaine 4% _____ Typical Anesthetic

Uses
 Temporarily relieves muscle soreness and minor joint pains in the wrist, knees, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use
 - on wounds or damaged skin
 - with a heating pad or device
 - with other ointments, creams, sprays, or liniments
 - if you are allergic to any ingredients of this product

When using this product

- use only as directed
 - avoid contact with the eyes, mucous membranes, or rashes
 - do not bandage tightly

Stop use and ask a doctor if

- skin reactions such as redness, swelling, blistering or other discomfort occur
 - symptoms persist for more than 7 days

If pregnant or breast-feeding,

ask a health professional before use.
 Keep out of the reach of children.
 If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old:
 Clean and dry the affected area.
 Tear off the protective film and apply the exposed part of the patch to the affected area.
 Carefully remove remaining film while pressing the patch firmly on the skin.
 Remove patch from the skin after at most 8-hour application.
 Children under age of 12: Consult a doctor

Inactive Ingredients

Dihydroxyethylum Aminoacetate, Disodium Ethylenediaminetetraacetic Acid, Glycerin, Kaolin, Methylparaben, Polyoxybutene-26, Propylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polyacrylate, Titanium Dioxide, Tartaric Acid, Water

Usage



Questions or Comments: support@sumifunofficial.com
 Business Cooperatives: business@sumifunofficial.com

Child Resistant Packaging



www.sumifunofficial.com
 YouTube: @sumifun
 Facebook: @sumifunOfficial
 Twitter: @sumifunOfficial

Sumifun®

NDC 83602-158-24

LIDOCAINE

PAIN RELIEF Gel-Patch



Numbing muscle and relieve pain:

- Odor-Free
- Maximum Strength
- Easy to apply and remove
- Single-Use Application

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	



24 PATCHES
 5.5" x 3.54"
 (14cm x 10cm)

Guangzhou Lidaqianxin Import & Export Trade Co., Ltd.
 117, Building A, No. 6, Dong Heng San Road,
 Zhuzhen, Tiesha District, Guangzhou, Guangdong, China

Made in PRC

Sumifun

Drug Facts

Active ingredient(s) _____ Purpose

Lidocaine 4% _____ Typical Anesthetic

Uses

Temporarily relieves muscle soreness and minor joint pains in the wrist, knee, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use

on wounds or damaged skin
 with a heating pad or device
 with other ointments, creams, sprays, or liniments
 if you are allergic to any ingredients of this product

When using this product

use only as directed
 avoid contact with the eyes, mucous membranes, or noses
 do not bandage tightly

Stop use and ask a doctor if

skin reactions such as redness, swelling, blistering or other discomfort occur
 symptoms persist for more than 7 days

If pregnant or breast-feeding,

ask a health professional before use.
KEEP OUT OF REACH OF CHILDREN.
 If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old:
 Clean and dry the affected area.
 Tear off the protective film and apply the exposed part of the patch to the affected area.
 Gently remove remaining film while pressing the patch firmly on the skin.
 Remove patch from the skin after at most 8-hour application.
 Children under age of 12: Consult a doctor

Inactive ingredients

Dihydroxyethylum Ammonioacetate,
 Disodium Ethylenediaminetetraacetic Acid, Glycerin,
 Ketin, Methylparaben, Polyethylene Glycol, Propylene Glycol,
 Polystyrene, Polyvinylpyrrolidone K90, Propylene Glycol,
 Sodium Polyacrylate, Titanium Dioxide, Terephthalic Acid, Water

Usage



Questions or Comments: cs.support@sumifunofficial.com
 Business Cooperation: business@sumifunofficial.com

Child Resistant Packaging



www.sumifunofficial.com
 INS: @sumifunofficial
 YouTube: @sumifun
 Facebook: @sumifunofficial
 TikTok: @sumifunofficial
 Twitter: @sumifunofficial

Sumifun®

LIDOCAINE PAIN RELIEF Gel-Patch



Numbing muscle and relieve pain:

- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	



25 PATCHES
 5.5" x 3.94"
 (14cm x 10cm)

Guangzhou Luqianmao Import & Export Trade Co., Ltd.
 117, Guoding A, No. 3, Dongfeng East Road,
 Zhuchun, Taishan District, Guangzhou/Guangdong, China.

Made in PRC

Sumifun®

Drug Facts

Active Ingredients: Lidocaine 4% Purpose: Topical Anesthetic

Uses: Temporarily relieves muscle soreness and minor joint pains in the wrist, knee, back, neck, hips, shoulders, elbows.

Warnings

For external use only

Do not use:
 - on wounds or damaged skin
 - with a heating pad or device
 - with other ointments, creams, sprays, or liniments
 - if you are allergic to any ingredients of this product

When using this product

- use only as directed
 - avoid contact with the eyes, mucous membranes, or nostrils
 - do not bandage tightly

Stop use and ask a doctor if

- skin reactions such as redness, swelling, blistering or other discomfort occur
 - symptoms persist for more than 7 days

If pregnant or breast-feeding,

ask a health professional before use.
KEEP OUT OF REACH OF CHILDREN.
 If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 13 years old:
 - Clean and dry the affected area.
 - Tear off the protective film and apply the exposed part of the patch to the affected area.
 - Carefully remove remaining film while pressing the patch firmly on the skin.
 - Remove patch from the skin after at most 8-hour application.
 - Children under age of 12: Consult a doctor

Inactive Ingredients

Dihydroxyaluminum Amorphacetate, Disodium Ethylenediaminetetraacetate Acid, Glycerin, Kaolin, Methylparaben, Polycarbonate, Propylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polycarboxylate, Titanium Dioxide, Tartaric Acid, Water

Usage



Questions or Comments: support@sumifunofficial.com
 Business Cooperation: business@sumifunofficial.com

Child Resistant Packaging



www.sumifunofficial.com
 INS: @sumifunofficial
 YouTube: @sumifun
 Facebook: @sumifunofficial
 TikTok: @sumifunofficial
 Twitter: @sumifunofficial

Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch

Numbing muscle and relieve pain:

- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	

4% LIDOCAINE

26 PATCHES
 5.5" x 3.94"
 (14cm x 10cm)

Guangzhou Lixianmao Import & Export Trade Co., Ltd.
 117 Building A, No. 6, Dong Heng San Road,
 Zhushan, Tianhe District, Guangzhou, Guangdong, China.

Made in PRC

Sumifun

Drug Facts

Active Ingredients: LidoCAINE 4%
Purpose: Topical Anesthetic

Uses: Temporarily relieves muscle soreness and minor joint pains in the wrist, lower back, neck, hips, shoulders, elbows.

Warnings: For external use only.

Do not use on wounds or damaged skin with a heating pad or device with other ointments, creams, lotions, or liniments if you are allergic to any ingredients of this product.

When using this product use only as directed avoid contact with the eyes, mucous membranes, or surfaces do not bandage tightly.

Stop use and ask a doctor if skin reactions such as redness, swelling, blistering or other discomfort occur symptoms persist for more than 7 days.

If pregnant or breast-feeding, ask a health professional before use.
KEEP OUT OF REACH OF CHILDREN.
 If misused accidentally, get medical help or contact with Poison Control Center immediately.

Directions: Adults and Children above 12 years old: Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor.

Inactive Ingredients: Dihydroxyacetone, Aminoacetic Acid, Glycine, Kaolin, Methylcellulose, Polyacrylate B9, Propylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Polyethylene Glycol, Polyacrylamide, Titanium Dioxide, Terbutyl Alcohol, Water.

Usage:

Questions or Comments: support@sumifunofficial.com
Business Cooperation: business@sumifunofficial.com

Child Resistant Packaging

www.sumifunofficial.com
 Instagram: @sumifunofficial
 Facebook: @sumifunofficial
 YouTube: @sumifun
 TikTok: @sumifunofficial
 Twitter: @sumifunofficial

Sumifun®


LIDOCAINE PAIN RELIEF Gel-Patch

Numbing muscle and relieve pain:



- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	



27 PATCHES
 5.5" x 3.34"
 (14cm x 10cm)

Guangzhou Liguanniao Import & Export Trade Co., Ltd.
 517, Guangyuan Road, No. 3, South Hong Shan Road,
 Zhuchun, Tianhe District, Guangzhou, Guangdong, China.

Made in PRC



Drug Facts

Active Ingredients: Lidocaine 4% Purpose: Topical Anesthetic

Uses: Temporarily relieves muscle soreness and minor joint pains in the wrist, knee, back, neck, hips, shoulders, elbows.

Warnings: For external use only

Do not use on wounds or damaged skin with a heating pad or device with other ointments, creams, sprays, or lotions if you are allergic to any ingredients of this product

When using this product: use only as directed avoid contact with the eyes, mucous membranes, or nose do not bandage tightly

Stop use and ask a doctor if: skin reactions such as redness, swelling, blistering or other discomfort occur symptoms persist for more than 7 days

If pregnant or breast-feeding, ask a health professional before use. KEEP OUT OF REACH OF CHILDREN. If swallowed accidentally, get medical help or contact with Poison Control Center immediately.

Directions

Adults and Children above 12 years old: Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor

Inactive Ingredients: Glycerylpropylamine, Anisococostate, Diiodium Ethylene diamine succinate, Acids, Glycerin, Methyl Methacrylate, Polyethylene Glycol, Polyacrylic Acid, Polystyrene, Polyisobutylene, Sodium Polycrylate, Titanium Dioxide, Tartaric Acid, Water

Usage



Questions or Comments: support@sumifunofficial.com Business Cooperation: business@sumifunofficial.com

Child Resistant Packaging



www.sumifunofficial.com
 @sumifunofficial
 YouTube: @sumifun
 Facebook: @sumifunofficial
 Twitter: @sumifunofficial

Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch



Numbing muscle and relieve pain:

- Odor-Free
- Maximum Strength
- Easy to apply and remove
- Single-Use Application

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	



29 PATCHES
 5.5" x 3.56"
 (14cm x 10cm)

Guangzhou Lianqianmao Import & Export Trade Co., Ltd.
 110, Baobangqin, No. 8, Dongfeng East Road,
 Zhuoa, Tianhe District, Guangzhou, Guangdong, China.

Made in PRC



sumifun

<p>Drug Facts</p> <p>Active Ingredients Purpose Lidocaine 4% Topical Anesthetic</p> <p>Uses Temporarily relieves muscle soreness and minor joint pains in the wrist, knee, back, neck, hips, shoulders, elbows.</p> <p>Warnings For external use only</p> <p>Do not use on wounds or damaged skin with a heating pad or device with other ointments, creams, sprays, or liniments if you are allergic to any ingredients of this product</p> <p>When using this product use only as directed avoid contact with the eyes, mucous membranes, or mouth do not bandage tightly</p> <p>Stop use and ask a doctor if skin reactions such as redness, swelling, blistering or other discomfort occur symptoms persist for more than 7 days</p> <p>If pregnant or breast feeding, ask a health professional before use. KEEP OUT OF REACH OF CHILDREN. If swallowed accidentally, get medical help or contact with Poison Control Center immediately.</p>	<p>Directions Adults and Children above 12 years old Clean and dry the affected area. Peel off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8-hour application. Children (under age of 12): Consult a doctor</p> <p>Inactive Ingredients Dihydroxyethylamine Acrylate, Dimethylammonium Methacrylate, Diiodium Ethylenediaminetetraacetic Acid, Glyceryl Methacrylate, Polyethylene Glycol, Polyethylene Glycol, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylparaben, Sodium Polycrylate, Titanium Dioxide, Talcum Acid, Water</p> <p>Usage</p> <p>Questions or Comments: support@sumifunofficial.com Business Correspondence: business@sumifunofficial.com</p> <p>Child Resistant Packaging</p>
---	---

www.sumifunofficial.com
 Instagram: @sumifunofficial
 YouTube: @sumifun
 Facebook: @sumifunofficial
 TikTok: @sumifunofficial
 Twitter: @sumifunofficial

Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch

Numbing muscle and relieve pain:

- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	

33 PATCHES
 5.5" x 3.5"
 (14cm x 10cm)



Guangzhou Lida Pharmaceutical Import & Export Trade Co., Ltd.
 117, Building A, No. 8, Dadi Road, Tianhe District,
 Guangzhou, P.R. China

Made in PRC

sumifun

<p>Drug Facts</p> <p>Active ingredients Lidocaine 4% Lidocaine 4% Typical Anesthetic</p> <p>Uses Temporarily relieves muscle soreness and minor joint pain in the wrist, knees, back, neck, hips, shoulders, elbows.</p> <p>Warnings For external use only.</p> <p>Do not use on wounds or damaged skin with a healing pad or device with other ointments, creams, lotions, or liniments. If you are allergic to any ingredients of this product.</p> <p>When using this product use only as directed and avoid contact with the eyes, mucous membranes, or nearby skin. Do not bandage tightly.</p> <p>Stop use and ask a doctor if skin reactions such as redness, swelling, blistering or other discomfort occur symptoms persist for more than 7 days.</p> <p>If pregnant or breast-feeding ask a health professional before use.</p> <p>KEEP OUT OF REACH OF CHILDREN. If swallowed, accidentally get medical help or contact with Poison Control Center immediately.</p>	<p>Directions Adults and Children above 12 years old: Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor.</p> <p>Inactive ingredients Dihydroxyaluminum Aminoacetate, Croscarell, Ethylene Glycol Dimethacrylate, Acrylate Glycol, Acrylic Methacrylate, Propylene Glycol, Polyacrylic Acid, Polyethylene Glycol, Polyethylene Glycol, Polystyrene, Titanium Dioxide, Talc, Zinc Oxide, Water.</p> <p>Usage</p> <p>Questions or Comments: support@sumifun.com Business Cooperation: business@sumifun.com</p> <p>Child Resistant Packaging</p>
---	---

www.sumifunofficial.com
 Instagram: @sumifunofficial
 Facebook: @sumifunofficial
 Twitter: @sumifunofficial

Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch

Numbing muscle and relieve pain:

- **Odor-Free**
- **Maximum Strength**
- **Easy to apply and remove**
- **Single-Use Application**



34 PATCHES
 5.5" x 3.54"
 (14cm x 9cm)

▪ Knee	▪ Back	▪ Neck
▪ Elbow	▪ Shoulders	



Guangzhou Lida Import & Export Trade Co., Ltd.
 117, Building A, No. 8, Dongfeng East Road,
 Zhusheng, Tianhe District, Guangzhou, Guangdong, China.

MADE IN PRC

Sumifun

Drug Facts		Directions
Active Ingredients Purpose	Lidocaine 4% Topical Anesthetic	Adults and Children above 12 years old: Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film when peeling the patch firmly on the skin. Remove patch from the skin after at least 6-hour application. Children under age of 12: Consult a doctor.
Uses Temporarily relieve muscle soreness and minor joint pains in the neck, lower back, neck, hips, shoulders, elbow.	Warnings For external use only.	Inactive Ingredients Dihydroxyethylamine, Ammonioacetate, Diiodoethyl-2-hydroxyethylsuccinate, Glyceryl Fosters, Methylphenyls, Polyethylene Glycol, Polyethylene Glycol, Polyethylene Glycol, Polyethylene Glycol, Sodium Polycrylate, Titanium Dioxide, Talc, Water.
Do not use on wounds or damaged skin with a heating pad or device with other ointments, creams, sprays, or liniments if you are allergic to any ingredients of this product.	When using this product use only as directed avoid contact with the eyes, mucous membranes, or mucosa do not bandage tightly.	Usage 1. Remove patch from outer envelope. 2. Peel off protective film. 3. Place on affected area. 4. Remove remaining film while pressing patch to skin.
Stop use and ask a doctor if skin reactions such as redness, swelling, blistering or other discomfort occur symptoms persist for more than 7 days.	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. If swallowed accidentally, get medical help or contact with Poison Control Center immediately.	Questions or Comments? customer@sumifunofficial.com Business Cooperation: business@sumifunofficial.com
		Child Resistant Packaging

www.sumifunofficial.com
 @NS @sumifunofficial
 @sumifunofficial
 @sumifunofficial
 @sumifunofficial

Sumifun®

LIDOCAINE


PAIN RELIEF Gel-Patch



Numbing muscle and relieve pain:

- Odor-Free
- Maximum Strength
- Easy to apply and remove
- Single-Use Application

• Knee	• Back	• Neck
• Elbow	• Shoulders	



35 PATCHES
 5.5" x 3.54"
 (14cm x 9cm)

Guangzhou Lidoanmiao Import & Export Trade Co., Ltd.
 517 Baofeng Road, 510630 Guangzhou, China

Made in PRC

Sumifun®

<p>Drug Facts</p> <p>Active ingredients: LIDOCAINE Hydrochloride Purpose: Topical Anesthetic</p> <p>Uses</p> <p>Temporarily relieves muscle aches and minor joint pains in the arm, forearm, back, neck, hip, shoulder, elbow.</p> <p>Warnings</p> <p>For external use only.</p> <p>Do not use</p> <p>on wounds or damaged skin with a stinging pain or burning with other anesthetics, ointments, sprays, or lotions. If you are allergic to any ingredients of the product.</p> <p>When using this product</p> <p>Use only as directed. Avoid contact with the eyes, mucous membranes, or mucus. Do not bandage tightly.</p> <p>Stop use and ask a doctor if</p> <p>skin reactions such as redness, swelling, blistering or other discomfort occur. Symptoms persist for more than 7 days.</p> <p>If pregnant or breast-feeding, ask a health professional before use. KEEP OUT OF REACH OF CHILDREN. If swallowed or accidentally get medical help or contact with Poison Control Center immediately.</p>	<p>Directions</p> <p>Adults and Children above 12 years old: Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor.</p> <p>Inactive ingredients</p> <p>Diphenylpicrylhydrazole, Hydroxyethylcellulose, Dimethylsiloxane, Polysorbate 80, Glycerin, Polyethylene Glycol, Polyethylene Glycol 400, Polyethylene Glycol 600, Polyethylene Glycol 800, Polyethylene Glycol 1000, Polyethylene Glycol 1500, Polyethylene Glycol 2000, Polyethylene Glycol 3000, Polyethylene Glycol 4000, Polyethylene Glycol 6000, Polyethylene Glycol 8000, Polyethylene Glycol 10000, Polyethylene Glycol 15000, Polyethylene Glycol 20000, Polyethylene Glycol 30000, Polyethylene Glycol 40000, Polyethylene Glycol 60000, Polyethylene Glycol 80000, Polyethylene Glycol 100000.</p> <p>Usage</p> <p>Questions or Comments: support@sumifunofficial.com Business Development: business@sumifunofficial.com</p> <p>Child Resistant Packaging</p>
---	---



www.sumifun.com
 Instagram: @sumifunofficial
 Facebook: @sumifunofficial
 YouTube: @sumifun
 TikTok: @sumifunofficial
 Twitter: @sumifunofficial

Sumifun®

LIDOCAINE

PAIN RELIEF Gel-Patch



Numbing muscle and relieve pain:

- Odor-Free
- Maximum Strength
- Easy to apply and remove
- Single-Use Application

• Knee	• Back	• Neck
• Elbow	• Shoulders	



43 PATCHES
 5.5" x 3.5"
 (14cm x 9cm)

Guangzhou Lixunsheng Import & Export Trade Co., Ltd.
 201, Guangde 3, No. 3, Guangde Street, Zhushan, Tianhe District, Guangzhou, Guangdong, China.

Made in PRC



Knee



Back



Neck



Elbow



Shoulder

sumifun

Drug Facts

Active ingredients: LIDOCAINE HCl
Purpose: Topical Anesthetic
Uses: Temporarily relieves muscle spasms and minor joint pains in the wrist, knee, back, neck, hip, shoulders, elbows.

Warnings: For external use only.

Do not use: on wounded or damaged skin with a history of use or direct with other ointments, creams, soaps, or lotions. If you are allergic to any ingredients of this product.

When using this product: use only as directed. Avoid contact with the eyes, mucous membranes, or other sensitive areas. Do not bandage tightly.

Stop use and ask a doctor if: you experience such as nausea, vomiting, dizziness or other symptoms occur. Symptoms persist for more than 7 days.

If pregnant or breast-feeding, ask a health care provider before use. **STOP USE** if breast milk decreases. If resolved, immediately get medical help or contact with Health Canada Consumer eNotify.

Directions: Adults and Children above 12 years old: Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while pressing the patch firmly on the skin. Remove patch from the skin after at most 8 hour application. Children under age of 12: Consult a doctor.

Inactive Ingredients: Dihydroxybutyltrim Acrylate, Dimethyl Ethylammoniumacetate, Acid, Glycolic, Maleic, Methylmethacrylate, Polyethylene Glycol, Polybutyl Acrylate, Polyvinylpyrrolidone, Polyethylene Terephthalate, Titanium Dioxide, Talcum, Zinc Oxide, Water.

Usage:



Customer or Distributor: sumifun@sumifun.com
 Business Cooperation: business@sumifun.com

Child Resistant Packaging



Sumifun

www.sumifunofficial.com
NSC @sumifunofficial
@sumifunofficial
@sumifunofficial
@sumifunofficial

Sumifun®

LIDOCAINE PAIN RELIEF Gel-Patch



Numbing muscle and relieve pain:

- Odor-Free
- Maximum Strength
- Easy to apply and remove
- Single-Use Application

• Knee	• Back	• Neck
• Elbow	• Shoulders	



44 PATCHES
3.5" x 3.84"
(10cm x 10cm)

Guangzhou Lidoan Pharmaceutical & Export Trade Co., Ltd.
411, Baotai Rd, No. 8, Tangjiao, Guangzhou
Zhuolu, Lantao District, Guangzhou, Guangdong, China

Made in PRC



Drug Facts

Active Ingredients: LIDOCAINE 4%
Proprietary Topical Anesthetic

Uses: Temporarily relieves muscle soreness and minor joint pain in the wrist, knee, back, neck, hips, shoulders, elbows.

Warnings: For external use only.

Do not use: on wounds or damaged skin; with heating pad or device; with other anesthetics, creams, sprays, or lotions; if you are allergic to any ingredients of this product.

When using this product: use only as directed; avoid contact with the eyes, mucous membranes, or holes; do not bandage tightly.

Stop use and ask a doctor if: pain worsens back to severe, swelling, blistering or other discomfort occurs; symptoms persist for more than 7 days.

If pregnant or breast-feeding: ask a health professional before use.
KEEP OUT OF REACH OF CHILDREN.
If contained in container, get medical help or contact with Poison Control Center immediately.

Directions: Adults and Children 12 years old: Clean and dry the affected area. Tear off the protective film and apply the exposed part of the patch to the affected area. Carefully remove remaining film while peeling the patch from the skin. Remove patch from the skin after at most 8-hour application. Children under age of 12: Consult a doctor.

Inactive Ingredients: Ethylhexylmethacrylate, Dimethylsiloxane, Dimethylacrylate, Diethylhexylsuccinate, Acid, Glyceryl, Benzyl, Methylacrylate, Hydroxyethyl Methacrylate, Polyethyl Methacrylate, Polybutyl Methacrylate, Dimethylsiloxane, Dimethylacrylate, Terephthalic Acid, Water.

Usage: Remove patch from seal envelope. Partially peel back protective film. Place on affected area. Remove remaining film while peeling patch to skin.

Questions or Comments: nsn@sumifunofficial.com
Business Cooperation: sumifun@sumifunofficial.com

Child Resistant Packaging



LIDOCAINE PAIN RELIEF GEL-PATCH

lidocaine patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83602-158
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	0.04 g in 1 g

Inactive Ingredients

Ingredient Name	Strength
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
GLYCERIN (UNII: PDC6A3C0OX)	

POVIDONE K90 (UNII: RDH86HJV5Z)
METHYLPARABEN (UNII: A2I8C7HI9T)
PROPYLPARABEN (UNII: Z8IX2SC1OH)
KAOLIN (UNII: 24H4NWX5CO)
TARTARIC ACID (UNII: W4888I119H)
WATER (UNII: 059QF0KO0R)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
POLYACRYLIC ACID (250000 MW) (UNII: 9G2MAD7J6W)
SODIUM POLYACRYLATE (2500000 MW) (UNII: 05I15JN12J)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
DIHYDROXYALUMINUM AMINOACETATE ANHYDROUS (UNII: 1K713C615K)
EDETATE DISODIUM (UNII: 7FLD91C86K)

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83602-158-26	26 in 1 BOX	04/03/2025	
1		12 g in 1 BAG; Type 0: Not a Combination Product		
2	NDC:83602-158-23	23 in 1 BOX	04/03/2025	
2		12 g in 1 BAG; Type 0: Not a Combination Product		
3	NDC:83602-158-27	27 in 1 BOX	04/03/2025	
3		12 g in 1 BAG; Type 0: Not a Combination Product		
4	NDC:83602-158-13	13 in 1 BOX	04/03/2025	
4		12 g in 1 BAG; Type 0: Not a Combination Product		
5	NDC:83602-158-17	17 in 1 BOX	04/03/2025	
5		12 g in 1 BAG; Type 0: Not a Combination Product		
6	NDC:83602-158-19	19 in 1 BOX	04/03/2025	
6		12 g in 1 BAG; Type 0: Not a Combination Product		
7	NDC:83602-158-22	22 in 1 BOX	04/03/2025	
7		12 g in 1 BAG; Type 0: Not a Combination Product		
8	NDC:83602-158-25	25 in 1 BOX	04/03/2025	
		12 g in 1 BAG; Type 0: Not a Combination Product		

8		12 g in 1 BAG; Type 0: Not a Combination Product	
9	NDC:83602-158-24	24 in 1 BOX	04/03/2025
9		12 g in 1 BAG; Type 0: Not a Combination Product	
10	NDC:83602-158-14	14 in 1 BOX	04/03/2025
10		12 g in 1 BAG; Type 0: Not a Combination Product	
11	NDC:83602-158-12	12 in 1 BOX	04/03/2025
11		12 g in 1 BAG; Type 0: Not a Combination Product	
12	NDC:83602-158-29	29 in 1 BOX	04/03/2025
12		12 g in 1 BAG; Type 0: Not a Combination Product	
13	NDC:83602-158-33	33 in 1 BOX	04/03/2025
13		12 g in 1 BAG; Type 0: Not a Combination Product	
14	NDC:83602-158-34	34 in 1 BOX	04/03/2025
14		12 g in 1 BAG; Type 0: Not a Combination Product	
15	NDC:83602-158-35	35 in 1 BOX	04/03/2025
15		12 g in 1 BAG; Type 0: Not a Combination Product	
16	NDC:83602-158-37	37 in 1 BOX	04/03/2025
16		12 g in 1 BAG; Type 0: Not a Combination Product	
17	NDC:83602-158-42	42 in 1 BOX	04/03/2025
17		12 g in 1 BAG; Type 0: Not a Combination Product	
18	NDC:83602-158-43	43 in 1 BOX	04/03/2025
18		12 g in 1 BAG; Type 0: Not a Combination Product	
19	NDC:83602-158-44	44 in 1 BOX	04/03/2025
19		12 g in 1 BAG; Type 0: Not a Combination Product	
20	NDC:83602-158-45	45 in 1 BOX	04/03/2025
20		12 g in 1 BAG; Type 0: Not a Combination Product	
21	NDC:83602-158-46	46 in 1 BOX	04/03/2025
21		12 g in 1 BAG; Type 0: Not a Combination Product	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC Monograph Drug	M017	04/03/2025	
--------------------	------	------------	--

Labeler - Guangzhou Liuquanmiao Import & Export Trade Co., Ltd. (418399704)

Registrant - Guangzhou Liuquanmiao Import & Export Trade Co., Ltd. (418399704)

Revised: 7/2025

Guangzhou Liuquanmiao Import & Export Trade Co., Ltd.