RIGHT REMEDIES LOW DOSE ASPIRIN 81MG- aspirin tablet Strive Pharmaceuticals Inc.

RIGHT REMEDIES LOW DOSE Aspirin 81mg

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)* *nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

• for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

ask your doctor about other uses for Safety Coated 81 mg Aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

This product contains Soy.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

• hives • facial swelling • asthma (wheezing) • shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for • gout • diabetes • arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint vomit blood have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

• drink a full glass of water with each dose

adults and children 12 years and over	 take 4 to 8 tablets every 4 hours do not take more than 48 tablets in 24 hours
children under 12 years	consult a doctor

Other information

- store at room temperature 20°–25°C (68°–77°F)
- Avoid excessive heat above 40°C(104°F)

Inactive ingredients

croscarmellose sodium, D&C Yellow No.10, lactose monohydrate, methacrylic acid copolymer, microcrystalline cellulose, polacrilin potassium, polyethylene glycol, polyvinyl alcohol, precipitated silica, purified talc, silicon dioxide, sodium lauryl sulphate, soy powder, talcum, titanium dioxide, triethyl citrate

Questions or comments?

1-888-577-8033 Monday - Friday 8am - 4pm EST

Compare to the active ingredient in **BAYER**[®] Low Dose aspirin[†]

Pain reliever

Aspirin regimen

Safety coated

Small, Easy-to Swallow tablets


TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

[†]This product is not manufactured or distributed by Bayer Healthcare LLC., owner of the registered trademark Bayer[®].

Distributed by: Strive Pharmaceuticals Inc., East Brunswick, NJ 08816

PRODUCT OF INDIA

Packaging



Drug Facts Label

Drug Facts Active ingredient (in each tablet) Purpose Do not use Aspirin 81 mg (NSAID)*.. Pain reliever nonsteroidal anti-inflammatory drug Uses for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief. ask your doctor about other uses for Safety Coated 81 mg Aspirin Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness This product contains Soy. Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed

Drug Facts (continued)

if you are allergic to aspirin or any other pain reliever/fever reducer

■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ stomach bleeding warning applies to you

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- vou are taking a diuretic
- vou have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

- Stop use and ask a doctor if
- an allergic reaction occurs. Seek medical help right away.
 you experience any of the following signs of stomach bleeding:
 feel faint vomit blood have bloody or black stools
- have stomach pain that does not get better pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

drink a full glass of water with each dose ■ take 4 to 8 tablets every 4 hours ■ do not take more than 48 tablets in 24 hours adults and children 12 years and over children under 12 years consult a doctor Other information store at room temperature between 20°-25°C (68°-77°F)
 Avoid excessive heat above 40°C (104°F) Inactive ingredients

croscarmellose sodium, D&C Yellow No.10, lactose monohydrate, methacrylic acid copolymer, microcrystalline cellulose, polacrilin potassium, polyethylene glycol, polyvinyl alcohol, precipitated silica, purified talc, silicon dioxide, sodium lauryl sulphate, soy powder, talcum, titanium dioxide, triethyl citrate

Questions or comments?

1-888-577-8033 Monday - Friday 8am - 4pm EST

RIGHT REMEDIES LOW DOSE ASPIRIN 81MG

aspirin tablet

Product Inform	nation							
Product Type		HUMAN OTC DP	RUG Item	n Code (So	Code (Source) NDC:70		92-168	
Route of Adminis	stration	ORAL						
Active Ingredie	ent/Activ	ve Moiety						
Ingredient Name Basis of Strengt						rength	Strengt	
ASPIRIN (UNII: R16C05Y76E) (ACETYLSALICYLIC ACID - UNII:R16C05Y76E) ASPIRIN								
• • • • • • • • • • • • • • • • • • • •								
Inactive Ingree	dients	In a vo di	ant Nama				Strength	
Ingredient Name CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)								
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)								
LACTOSE MONOHY								
METHACRYLIC ACI	-		COPOLYMER (1:1) (UNII: 74	4G4R6TH13)			
MICROCRYSTALLIN				, (=,				
POLACRILIN POTAS								
POLYETHYLENE GL			WJQ0SDW1A)					
POLYVINYL ALCOH			-					
SILICON DIOXIDE (•					
TALC (UNII: 7SEV7J4	R1U)							
SODIUM LAURYL S	ULFATE (UI	NII: 368GB5141J)						
SOYBEAN (UNII: L7F	T8F1ZOD)							
TITANIUM DIOXIDE	(UNII: 15FI)	X9V2JP)						
TRIETHYL CITRATE	(UNII: 8Z9	6QXD6UM)						
Product Chara	cteristic	S						
Color yellow			Score no score					
Shape		ROUND	Size 7mm		7mm			
Flavor				F	C			
Contains								
contains								
Packaging								
# Item Code	F	Package Description			Marketing Start Date		Marketing End Date	
1 NDC:70692-168- 50	500 in 1 BC Product	OTTLE; Type 0: Not a Combination		04/15/2025	4/15/2025			
Marketing I	nforma	ation						
Marketing			Mark	Marketing Start Date		Marketing End Date		
Category		Citation	-				Jale	
OTC Monograph Dru	g M013	Citation		04/15/20			Jale	

Revised: 4/2025

Strive Pharmaceuticals Inc.