

**MUCINEX DAY TIME PAIN, HEADACHE AND MUCUS CONGESTION AND
MUCINEX NIGHT TIME HEADACHE, COUGH AND FEVER MAXIMUM STRENGTH-
acetaminophen, dextromethorphan hydrobromide, guaifenesin, triprolidine
hydrochloride
RB Health (US) LLC**

**MUCINEX® Rapid Clear® Pain, Headache & Mucus Congestion and Nighttime
Caplets**

Drug Facts

Acetaminophen 325 mg
Guaifenesin 200 mg

Mucinex Nightshift Night Sinus Purposes
Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough suppressant
Triprolidine HCl 1.25 mg Antihistamine

Active ingredients (in each caplet)

<i>Mucinex Sinus-Max Day Pressure, Pain & Cough</i>	<i>Purposes</i>
Acetaminophen 325 mg	Pain reliever
Guaifenesin 200 mg	Expectorant

Active ingredients (in each caplet)

<i>Mucinex Nightshift Night Sinus</i>	<i>Purposes</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Triprolidine HCl 1.25 mg	Antihistamine

Uses

Mucinex Sinus-Max Day Pressure, Pain & Cough

- temporarily relieves:
 - nasal congestion
 - headache
 - minor aches and pains
 - cough
 - sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Mucinex Nightshift Night Sinus

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - sneezing
 - sinus congestion and pressure
 - runny nose
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning

This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning (Nightshift Night Sinus only)

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription

drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma (**Nightshift Night Sinus only**)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis (**Nightshift Night Sinus only**)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (**Nightshift Night Sinus only**)

When using this product

- **do not use more than directed**
- excitability may occur, especially in children (**Nightshift Night Sinus only**)
- marked drowsiness may occur (**Nightshift Night Sinus only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**Nightshift Night Sinus only**)
- avoid alcoholic drinks (**Nightshift Night Sinus only**)
- use caution when driving a motor vehicle or operating machinery (**Nightshift Night Sinus only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Mucinex Sinus-Max Day Pressure, Pain & Cough

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Mucinex Nightshift Night Sinus

- **do not take more than directed (see Overdose warning)**
- do not take more than 8 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

Mucinex Sinus-Max Day Pressure, Pain & Cough

croscarmellose sodium, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

Mucinex Nightshift Night Sinus

croscarmellose sodium, crospovidone, hypromellose, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224

PRINCIPAL DISPLAY PANEL - Kit Carton

NDC 72854-210

MAXIMUM STRENGTH

Mucinex[®]

RAPID+CLEAR[™]

DAYTIME
PAIN, HEADACHE &
MUCUS CONGESTION

NIGHT-TIME
HEADACHE, COUGH & FEVER

Maximum Strength per 4-hour dose
Take only as directed.
Keep carton for full information.

reckitt

Dist. by: RB Health (US)
Paramparany, NJ 07654-0224
©2025 RB Health
Patents: www.reckitt.com/patents
3320664 112524

MAXIMUM STRENGTH

NDC 72854-210-20

Mucinex[®]

RAPID+CLEAR[™]

DAYTIME
PAIN, HEADACHE &
MUCUS CONGESTION

Acetaminophen »
Pain Reliever/Fever Reducer
Guaifenesin » Expectorant

AGES 12+

- ✓ Pain & Headache
- ✓ Fever
- ✓ Thins & Loosens Mucus

12 CAPLETS

NIGHT-TIME
HEADACHE,
COUGH & FEVER

Acetaminophen » Pain Reliever/Fever Reducer
Dextromethorphan HBr » Cough Suppressant
Triprolidine HCl » Antihistamine

AGES 12+

- ✓ Pain & Headache
- ✓ Fever + Itchy Throat
- ✓ Cough

8 CAPLETS

RAPID RELEASE

TOTAL 20 CAPLETS

Tamper evident:
Do not use if carton
is damaged or if
printed seal on
blister is broken or
missing.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

how2recycle.info

MULTI-LAYER
PAPER
BOX
TRAY

3 63824 99938 8

Do not take the Rapid+Clear[™] Daytime Pain, Headache & Mucus Congestion and Rapid+Clear[™] Nighttime Headache, Cough & Fever caplets at the same time. Always wait at least 4 hours before taking another dose of Mucinex[®] caplets. Keep carton for complete product information. Do not take more than directed (see Overdose warning).

DAYTIME PAIN, HEADACHE & MUCUS CONGESTION	NIGHTTIME HEADACHE, COUGH & FEVER
Drug Facts Active ingredients (in each caplet) Acetaminophen 325 mg, Pain reliever/fever reducer Guaifenesin 200 mg, Expectorant	Drug Facts Active ingredients (in each caplet) Acetaminophen 325 mg, Pain reliever/fever reducer Dextromethorphan HBr 10 mg, Cough suppressant Triprolidine HCl 1.25 mg, Antihistamine
Uses temporarily relieves these common cold and flu symptoms: minor aches and pains headache sore throat helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of both mucus and mucus made more productive temporarily reduces fever	Uses temporarily relieves these common cold and flu symptoms: minor aches and pains headache sore throat helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of both mucus and mucus made more productive temporarily reduces fever
Warnings Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take: more than 12 caplets in 24 hours, which is the maximum daily amount for this product with other drugs containing acetaminophen 3 or more alcoholic drinks daily while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness skin rash if a skin reaction occurs, stop use and seek medical help right away. If severe throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use: with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.	Warnings Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take: more than 8 caplets in 24 hours, which is the maximum daily amount for this product with other drugs containing acetaminophen 3 or more alcoholic drinks daily while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness skin rash if a skin reaction occurs, stop use and seek medical help right away. If severe throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use: with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

DAYTIME PAIN, HEADACHE & MUCUS CONGESTION (cont'd)	NIGHTTIME HEADACHE, COUGH & FEVER (cont'd)
Drug Facts (continued) Ask a doctor before use if you have: liver disease persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema cough that occurs with too much phlegm (mucus) Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. When using this product do not use more than directed. Stop use and ask a doctor if: pain or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days nausea or vomiting is present cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions Do not take more than directed (see Overdose warning).	Drug Facts (continued) Do not use: with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. Ask a doctor or pharmacist before use if you are: taking the blood thinning drug warfarin taking sedatives or tranquilizers When using this product: do not use more than directed excitability may occur, especially in children marked drowsiness may occur alcohol, sedatives, and tranquilizers may increase drowsiness avoid alcoholic drinks. Stop use and ask a doctor if: pain or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days nausea or vomiting is present cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions Do not take more than directed (see Overdose warning).

DAYTIME PAIN, HEADACHE & MUCUS CONGESTION (cont'd)	NIGHTTIME HEADACHE, COUGH & FEVER (cont'd)
Drug Facts (continued) Do not use: with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. Ask a doctor or pharmacist before use if you are: taking the blood thinning drug warfarin taking sedatives or tranquilizers When using this product: do not use more than directed excitability may occur, especially in children marked drowsiness may occur alcohol, sedatives, and tranquilizers may increase drowsiness avoid alcoholic drinks. Stop use and ask a doctor if: pain or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days nausea or vomiting is present cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions Do not take more than directed (see Overdose warning).	Drug Facts (continued) Do not use: with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. Ask a doctor or pharmacist before use if you are: taking the blood thinning drug warfarin taking sedatives or tranquilizers When using this product: do not use more than directed excitability may occur, especially in children marked drowsiness may occur alcohol, sedatives, and tranquilizers may increase drowsiness avoid alcoholic drinks. Stop use and ask a doctor if: pain or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days nausea or vomiting is present cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions Do not take more than directed (see Overdose warning).

MUCINEX DAY TIME PAIN, HEADACHE AND MUCUS CONGESTION

AND MUCINEX NIGHT TIME HEADACHE, COUGH AND FEVER MAXIMUM STRENGTH

acetaminophen, dextromethorphan hydrobromide, guaifenesin, triprolidine hydrochloride kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72854-210
--------------	----------------	--------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-210-10	1 in 1 CARTON	07/01/2025	
1		1 in 1 KIT		
2	NDC:72854-210-20	2 in 1 CARTON	07/01/2025	
2		1 in 1 KIT		
3	NDC:72854-210-40	4 in 1 CARTON	07/01/2025	
3		1 in 1 KIT		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BLISTER PACK	6
Part 2	1 BLISTER PACK	4

Part 1 of 2

MAXIMUM STRENGTH MUCINEX SINUS-MAX PRESSURE, PAIN AND COUGH

acetaminophen, guaifenesin tablet, film coated

Product Information

Item Code (Source)	NDC:72854-208
--------------------	---------------

Route of Administration	ORAL
-------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	

FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	VVV;MSC
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/01/2025	

Part 2 of 2

MUCINEX NIGHTSHIFT SINUS MAXIMUM STRENGTH

acetaminophen, dextromethorphan hydrobromide, and triprolidine hydrochloride tablet, coated

Product Information

Item Code (Source)	NDC:72854-209
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE -	TRIPROLIDINE	1.25 mg

UNII:2L8T9S52QM)	HYDROCHLORIDE	1.25 mg
------------------	---------------	---------

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (UNII: 2S7830E561)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	

Product Characteristics

Color	yellow	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	VW;LOGOcrescentmoonplus
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/01/2025	

Labeler - RB Health (US) LLC (081049410)