

# **BENZETHONIUM CHLORIDE AND DYCLONINE HYDROCHLORIDE- liquid bandage liquid**

**Target Corporation**

-----

## **Target Liquid Bandage**

### ***Active ingredients***

Benzethonium chloride 0.2%  
Dyclonine hydrochloride 0.75%

### ***Purpose***

First aid antiseptic  
Topical analgesic

### ***Uses***

- first aid to help prevent infection in minor cuts, scrapes and burns
- first aid for the temporary relief of pain in minor cuts, scrapes, and burns

### ***Warnings***

**For external use only.**

#### **Flammable:**

Keep away from fire or flame.

#### **Do not use**

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor

#### **Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

#### **Stop use and ask a doctor if**

- conditions persists or gets worse
- infection occurs
- symptoms persist for more than 7 days or clear up and occur again within a few days

#### **Keep out of reach of children.**

If swallowed, get medical help or call a Poison Control Center right away.

## **When using this product**

- do not store at temperature above 120°F

Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal.

## **Directions**

- for adults and children 2 years of age and older
- clean the affected area
- apply a small amount of this product on the area 1-3 times daily and let dry
- may be covered with a sterile bandage
- if bandaged, let dry first
- children under 2 years of age: do not use, consult doctor

## **Other information**

- **cap bottle tightly**
- wipe top of bottle clean after each use
- store between 20 °C to 25 °C ( 68 ° to 77 °F)
- may temporarily sting upon application
- to remove, apply more Liquid Bandage and quickly wipe off
- do not allow to come into contact with fabrics, floors, countertops, or other surfaces
  - will stain

## **Inactive ingredients**

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

## **Questions**

Call 1-866-964-0939


## **Principal Display Panel**

Liquid Bandage

Benzethonium chloride/First Aid Antiseptic  
Dyclonine hydrochloride 0.75%/ Topical Analgesic

Eliminates 99.9% of the three most common  
germs within an hour of contact

1 FL. OZ. (29.6 mL)





# Liquid Bandage

Benzethonium Chloride/  
First Aid Antiseptic  
Dyclonine Hydrochloride 0.75%/  
Topical Analgesic



53-033TG-02

# Liquid Bandage



up&up™

Benzethonium Chloride/First Aid Antiseptic  
Dyclonine Hydrochloride 0.75%/Topical Analgesic  
Eliminates 99.9% of the three most common  
germs within an hour of contact\*

1 FL. OZ. (29.6 mL)



245 07 0553 R00 C-002262-01-035-0000



Distributed by Target Corporation  
Minneapolis, MN 55403  
Made in USA with U.S. and  
imported parts  
TM & ©2024 Target Brands, Inc.

\*Three most common germs used are  
Target.com/comments or 1-800-910-6874.  
We welcome any questions you may have at  
k. pneumoniae, E. coli, P. aeruginosa.

**Satisfaction guaranteed –  
Love it or your money back.**

### Questions? Call 1-866-964-0939

Inactive ingredients: acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

**Other information** ■ cap bottle tightly ■ wipe top of bottle clean after each use ■ store between 20° to 25°C (68° to 77°F) ■ may temporarily sting upon application ■ to remove, apply more Liquid Bandage and quickly wipe off ■ do not allow to come into contact with fabrics, floors, countertops or other surfaces -- will stain.

**Directions** ■ for adults and children 2 years of age and older ■ clean the affected area ■ apply a small amount of this product on the area 1 to 3 times daily and let dry ■ may be covered with a sterile bandage ■ if bandaged, let dry first ■ children under 2 years of age: do not use, consult a doctor.


**Keep out of reach of children.** If swallowed, get medical help or call a Poison Control Center right away.

**Stop use and ask a doctor if** ■ condition persists or gets worse ■ infection occurs ■ symptoms persist for more than 7 days or clear up and occur again within a few days

**Ask a doctor before use if you have** ■ deep or puncture wounds ■ animal bites ■ serious burns

**Do not use** ■ in the eyes ■ over large areas of the body ■ longer than 1 week unless directed by a doctor

**Flammable:** Keep away from fire or flame.  
For external use only.  
**Warnings**



## BENZETHONIUM CHLORIDE AND DYCLONINE HYDROCHLORIDE

liquid bandage liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:82442-019
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BENZETHONIUM CHLORIDE</b> (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	2 mg in 1 mL
<b>DYCLONINE HYDROCHLORIDE</b> (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q300)	DYCLONINE HYDROCHLORIDE	7.5 mg in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>AMYL ACETATE</b> (UNII: 92Q24NH7AS)	
<b>ACETONE</b> (UNII: 1364PS73AF)	
<b>CASTOR OIL</b> (UNII: D5340Y2I9G)	
<b>ETHYL ACETATE</b> (UNII: 76845O8NMZ)	
<b>PYROXYLIN</b> (UNII: KYR8BR2X6O)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82442-019-01	1 in 1 CARTON	04/09/2025	
1		29.6 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	04/09/2025	

**Labeler** - Target Corporation (006961700)

Revised: 10/2025

Target Corporation