

VICKS DAYQUIL INTENSE FLU- acetaminophen, dextromethorphan hbr liquid
The Procter & Gamble Manufacturing Company

Vicks[®] DayQuil[™] INTENSE FLU MULTI-SYMPTOM Liquid

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 1000 mg

Dextromethorphan HBr 30 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 3 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain or cough get worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 3 doses (30 mL each) per day (24 hrs)

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL contains:** sodium 24 mg
- store at no greater than 25°C (77°F) and do not refrigerate

Inactive ingredients

alcohol, anhydrous citric acid, FD&C Red No. 40, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, water

Questions?

1-800-362-1683

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND ON BOTTLE IS BROKEN OR MISSING.

DIST. BY:

PROCTER & GAMBLE

CINCINNATI, OH 45202

PRINCIPAL DISPLAY PANEL - 354 ml Bottle Label

VICKS®

DayQuil™

INTENSE FLU

MULTI-SYMPTOM

Acetaminophen, Dextromethorphan HBr

Pain reliever/Fever reducer, Cough suppressant

BODY ACHES

FEVER

HEADACHE

SORE THROAT

COUGH

NEW FORMULA

Daytime Relief

Alcohol 10%

12 FL OZ (354 mL)



Drug Facts (continued)
Other information <ul style="list-style-type: none">• each 30 mL contains: sodium 24 mg• store at no greater than 25°C (77°F) and do not refrigerate
Inactive ingredients alcohol, anhydrous citric acid, FD&C Red No. 40, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, water
Questions? 1-800-362-1683

Drug Facts (continued)
acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have <ul style="list-style-type: none">• liver disease• cough that occurs with too much phlegm (mucus)• persistent or chronic cough such as occurs with smoking, asthma, or emphysema• a sodium-restricted diet
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
Stop use and ask a doctor if <ul style="list-style-type: none">• pain or cough get worse or last more than 7 days• fever gets worse or lasts more than 3 days• redness or swelling is present• new symptoms occur• cough comes back, or occurs with rash or headache that lasts.
These could be signs of a serious condition.
If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions <ul style="list-style-type: none">• take only as directed• only use the dose cup provided• do not exceed 3 doses (30 mL each) per day (24 hrs)
adults & children 12 yrs & over 30 mL every 6 hrs
children 4 to under 12 yrs ask a doctor
children under 4 yrs do not use

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND IS BROKEN OR MISSING. ← PEEL BACK FOR DRUG FACTS	
Drug Facts	
Active ingredients (in each 30 mL) Acetaminophen 1000 mg Dextromethorphan HBr 30 mg	Purpose Pain reliever/fever reducer Cough suppressant
Uses temporarily relieves common cold/flu symptoms: <ul style="list-style-type: none">• cough due to minor throat & bronchial irritation• sore throat• headache• minor aches & pains• fever	
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take <ul style="list-style-type: none">• more than 3 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product• with other drugs containing acetaminophen• 3 or more alcoholic drinks every day while using this product Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none">• skin reddening• blisters• rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
Do not use <ul style="list-style-type: none">• with any other drug containing	
PARENTS: Learn about teen medicine abuse www.StopMedicineAbuse.org	
P&G DIST. BY: PROCTER & GAMBLE, CINCINNATI, OH 45202 Patents: www.pg.com/patents 21268816	23900-041875

VICKS DAYQUIL INTENSE FLU

acetaminophen, dextromethorphan hbr liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84126-356
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	1000 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	

SUCRALOSE (UNII: 96K6UQ3ZD4)				
ALCOHOL (UNII: 3K9958V90M)				
Product Characteristics				
Color	red		Score	
Shape			Size	
Flavor	CHERRY (Black Cherry)		Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84126-356-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/01/2025	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	07/01/2025	

Labeler - The Procter & Gamble Manufacturing Company (004238200)