

**VICKS DAYQUIL AND VICKS NYQUIL INTENSE FLU- acetaminophen,
dextromethorphan hbr
The Procter & Gamble Manufacturing Company**

**Vicks[®] DayQuil[™] and NyQuil[™] INTENSE FLU MULTI-SYMPTOM Liquid
Convenience Pack**

Vicks[®] NyQuil[™] INTENSE FLU MULTI-SYMPTOM Liquid

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 1000 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purpose

Pain reliever/ fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- cough to help you sleep
- sore throat
- headache
- minor aches & pains
- fever
- runny nose & sneezing

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 3 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL contains:** sodium 24 mg
- store at no greater than 25°C and do not refrigerate

Inactive ingredients

alcohol, anhydrous citric acid, FD&C Red No. 40, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, water

Questions?

1-800-362-1683

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND ON BOTTLE IS BROKEN OR MISSING.

DIST. BY: PROCTER & GAMBLE,

CINCINNATI, OH 45202

Vicks[®] DayQuil[™] INTENSE FLU MULTI-SYMPTOM Liquid

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 1000 mg

Dextromethorphan HBr 30 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 3 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain or cough get worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 3 doses (30 mL each) per day (24 hrs)

adults & children 12 yrs & over

children 4 to under 6 yrs

children under 4 yrs

30 mL every 6 hrs

ask a doctor

do not use

Other information

- **each 30 mL contains:** sodium 24 mg
- store at no greater than 25°C (77°F) and do not refrigerate

Inactive ingredients

alcohol, anhydrous citric acid, FD&C Red No. 40, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, water

Questions?

1-800-362-1683

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND ON BOTTLE IS BROKEN OR MISSING.

**DIST. BY: PROCTER & GAMBLE,
CINCINNATI, OH 45202**

PRINCIPAL DISPLAY PANEL - Convenience Pack

VALUE PACK

VICKS ® DayQuil™ INTENSE FLU

MULTI-SYMPTOM

Acetaminophen, Dextromethorphan HBr

Pain reliever/Fever reducer, Cough suppressant

BODY ACHES

FEVER

HEADACHE

SORE THROAT

COUGH

Daytime Relief

Alcohol 10%

VICKS ® NyQuil™ INTENSE FLU

MULTI-SYMPTOM

Acetaminophen- Pain reliever/fever reducer

Doxylamine Succinate - Antihistamine

Dextromethorphan HBr - Cough suppressant

BODY ACHES

FEVER

HEADACHE

SORE THROAT

2 BOTTLES - 12 FL OZ (354 mL) EACH; TOTAL 24 FL OZ (708 mL)



1	NDC:84126-358-24	1 in 1 PACKAGE; Type 0: Not a Combination Product	07/01/2025	
---	------------------	---	------------	--

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE, PLASTIC	354 mL
Part 2	1 BOTTLE, PLASTIC	354 mL

Part 1 of 2

VICKS DAYQUIL INTENSE FLU
acetaminophen, dextromethorphan hbr liquid

Product Information

Item Code (Source)	NDC:84126-356
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	1000 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
ALCOHOL (UNII: 3K9958V90M)	

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	CHERRY (Black Cherry)	Imprint Code	

Contains

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84126-356-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/01/2025	

Part 2 of 2

VICKS NYQUIL INTENSE FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid

Product Information

Item Code (Source)	NDC:84126-357
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	1000 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ALCOHOL (UNII: 3K9958V90M)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SORBITOL (UNII: 506T60A25R)	
GLYCERIN (UNII: PDC6A3C0OX)	

TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

Product Characteristics			
Color	red	Score	
Shape		Size	
Flavor	CHERRY (Black Cherry)	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84126-357-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/01/2025	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/01/2025	

Labeler - The Procter & Gamble Manufacturing Company (004238200)