

**VICKS DAYQUIL AND NYQUIL VAPOCOOL COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate  
The Procter & Gamble Manufacturing Company**

---

**Vicks® DayQuil™ & Nyquil™ VapoCOOL COLD & FLU, Coated Caplets**

VICKS DayQuil VapoCOOL COLD & FLU, Coated Caplets

**Active ingredients (in each caplet)**

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg

**Purpose**

Pain reliever/Fever reducer

Cough suppressant

**Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

**Warnings**

**Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by

fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- take only as directed
- do not exceed 8 caplets per 24 hrs

---

adults & children 12 yrs & over

2 caplets with water every 4 hrs

children 4 to under 12 yrs  
children under 4 yrs

ask a doctor  
do not use

---

## **Other information**

- store at no greater than 25°C (77°F)

## **Inactive ingredients**

corn starch, crospovidone, FD&C Yellow No. 6 Aluminum Lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

## **Questions?**

**1-800-362-1683**

Vicks NyQuil VapoCOOL COLD & FLU, Coated Caplets

## **Active ingredients (in each caplet)**

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg  
Doxylamine succinate 6.25 mg

## **Purpose**

Pain reliever/Fever reducer  
Cough suppressant  
Antihistamine

## **Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- cough to help you sleep
- sore throat
- headache
- minor aches & pains
- fever
- runny nose & sneezing

## **Warnings**

### **Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount for this product

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

### **Allergy Alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

### **Sore throat warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may cause drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- do not exceed 8 caplets per 24 hrs

adults & children 12 yrs & over	2 caplets with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- store at no greater than 25°C (77°F)

**Inactive ingredients**

corn starch, D&C Yellow #10 Aluminum Lake, FD&C Blue #2 Aluminum Lake, FD&C Yellow #6 Aluminum Lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

**Questions?**

**1-800-362-1683**

**TAMPER EVIDENT: THIS PACKAGE IS SAFETY-SEALED & CHILD RESISTANT. USE ONLY IF BLISTERS ARE INTACT. IF DIFFICULT TO OPEN, USE SCISSORS.**

**DIST. BY PROCTER & GAMBLE,  
CINCINNATI, OH 45202.**

**PRINCIPAL DISPLAY PANEL - Kit Carton**

Vicks®

DayQuil™ VapoCOOL™ COLD & FLU + CONGESTION

**Acetaminophen**, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

NyQuil™ VapoCOOL™ COLD & FLU + CONGESTION

**Acetaminophen**, Doxylamine Succinate, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

16 DAYQUIL COATED CAPLETS, 8 NYQUIL COATED CAPLETS: 24 TOTAL COATED CAPLETS

**Drug Facts (continued)**

contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

• take only as directed	• do not exceed 8 caplets per 24 hrs
adults & children 12 yrs & over	2 caplets with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- store at no greater than 25°C (77°F)

**Inactive ingredients** corn starch, crospovidone, D&C Yellow #10, Aluminum Lake, FD&C Blue #2 Aluminum Lake, FD&C Yellow #6, Aluminum Lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, stearic acid, sucrose, talc, titanium dioxide

**Questions?** 1-800-362-1683

**Drug Facts (continued)****Directions**

adults & children 12 yrs & over	2 caplets with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- store at no greater than 25°C (77°F)

**Inactive ingredients** corn starch, crospovidone, FD&C Yellow No. 6 Aluminum Lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, stearic acid, sucrose, talc, titanium dioxide

**Questions?** 1-800-362-1683

**PARENTS:**

Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

TAMPER EVIDENT: THIS PACKAGE IS SAFETY-SEALED & CHILD RESISTANT. USE ONLY IF BLISTERS ARE INTACT. IF DIFFICULT TO OPEN, USE SCISSORS.

ACTUAL SIZE

**DayQuil™****VapoCOOL™****COLD & FLU****Acetaminophen.**

Dextromethorphan HBr

Headache, Fever,  
Sore Throat, Minor  
Aches & Pains

Cough

VapoCOOL  
Coated

Daytime Relief

16 DAYQUIL COATED CAPLETS 24 TOTAL COATED CAPLETS

**VALUE PACK****NyQuil™****VapoCOOL™****COLD & FLU****Acetaminophen, Doxylamine**

Succinate, Dextromethorphan HBr

Headache, Fever,  
Sore Throat, Minor  
Aches & Pains

Sneezing, Runny Nose

Cough

VapoCOOL  
Coated

Nighttime Relief

8 NYQUIL COATED CAPLETS

21349759



ACTUAL SIZE

DIST. BY: PROCTER & GAMBLE,  
CINCINNATI, OH 45202

**P&G**

Patents:  
[www.pg.com/patents](http://www.pg.com/patents)

**Drug Facts | NyQuil + VapoCool Cold & Flu****Active ingredients (in each caplet)**

Acetaminophen 325 mg Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg Cough suppressant  
Doxylamine succinate 6.25 mg Antihistamine

**Purpose**

Acetaminophen 325 mg Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg Cough suppressant

**Uses** temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- cough to help you sleep • sore throat • headache
- minor aches & pains • fever • runny nose & sneezing

**Warnings**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 8 caplets in 24 hours, which is the maximum daily amount for this product.

With other drugs containing acetaminophen  
• 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash

**Drug Facts | DayQuil + VapoCool Cold & Flu****Active ingredients (in each caplet)**

Acetaminophen 325 mg Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg Cough suppressant

**Purpose**

Acetaminophen 325 mg Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg Cough suppressant

**Uses** temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

**Warnings**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

more than 8 caplets in 24 hours, which is the maximum daily amount for this product.

With other drugs containing acetaminophen

• 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash

**Drug Facts (continued)**

If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease • glaucoma • cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers • taking the blood thinning drug warfarin
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present • new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease • cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.****Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**VICKS DAYQUIL AND NYQUIL VAPOCOOL COLD AND FLU**

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate kit

## Product Information

### Product Type

HUMAN OTC DRUG

### Item Code (Source)

NDC:84126-354

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84126-354-24	12 in 1 CARTON	05/06/2025	
1		1 in 1 BLISTER PACK; Type 1: Convenience Kit of Co-Package		
2	NDC:84126-354-48	24 in 1 CARTON	05/06/2025	
2		1 in 1 BLISTER PACK; Type 1: Convenience Kit of Co-Package		

## Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 POUCH	2
Part 2	8 BLISTER PACK	8

## Part 1 of 2

### VICKS DAYQUIL VAPOCOOL COLD AND FLU

acetaminophen, dextromethorphan hydrobromide tablet, coated

## Product Information

### Item Code (Source)

NDC:84126-352

### Route of Administration

ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

## Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: 08232NY3SJ)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)	

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
POVIDONE (UNII: FZ989GH94E)	
CROSPovidone (UNII: 2S7830E561)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

## Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	DQV
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84126-352-02	2 in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/06/2025	

## Part 2 of 2

### VICKS NYQUIL VAPOCOOL COLD AND FLU

acetaminophen, doxylamine succinate, dextromethorphan hydrobromide tablet, coated

## Product Information

Item Code (Source)	NDC:84126-353
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH)	DEXTROMETHORPHAN	10 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>FD&amp;C YELLOW NO. 6 ALUMINUM LAKE</b> (UNII: GYP6Z2JR6Q)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYVINYL ALCOHOL</b> (UNII: 532B59J990)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE</b> (UNII: CQ3XH3DET6)	
<b>FD&amp;C BLUE NO. 2 ALUMINUM LAKE</b> (UNII: 4AQJ3LG584)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPovidone</b> (UNII: 2S7830E561)	

**Product Characteristics**

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	NQV
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/06/2025	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/06/2025	

**Labeler -** The Procter & Gamble Manufacturing Company (004238200)

Revised: 1/2026

The Procter & Gamble Manufacturing Company