

**WALGREENS NIGHT TIME WAL-FLU SEVERE COLD AND COUGH HONEY LEMON FLAVOR INFUSED WITH CHAMOMILE AND WHITE TEA FLAVORS- acetaminophen, diphenhydramine hcl, and phenylephrine hcl. granule, for solution  
Walgreen Co.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

**Walgreens Nighttime Wal-flu severe cold and cough honey lemon flavor infused with chamomile and white tea flavors**

**Active ingredients (in each packet) Purpose**

Acetaminophen 650 mg .....Pain reliever/Fever reducer  
Diphenhydramine hydrochloride 25 mg .....Antihistamine/Cough suppressant  
Phenylephrine hydrochloride 10 mg .....Nasal decongestant.

**Uses**

- temporarily relieves:
- minor aches and pains - minor sore throat pain
- headache - nasal and sinus congestion
- runny nose - sneezing
- itchy nose and throat - itchy, watery eyes due to hay fever
- cough due to minor throat and bronchial irritation
- temporarily reduces fever.

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take - more than 6 packets in 24 hours, which is the maximum daily amount - with other drugs containing acetaminophen - 3 or more alcoholic drinks every day while using this product.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription and nonprescription). Ask a doctor or pharmacist before using with other drugs if you are not sure.
- with any other product containing diphenhydramine, even one used on the skin
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease - heart disease - high blood pressure
- thyroid disease - diabetes - glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema, asthma, or chronic bronchitis

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema.

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin.

**When using this product**

- do not exceed recommended dosage
- avoid alcoholic drinks - marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children.

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days - redness or swelling is present
- new symptoms occur - symptoms do not get better or worsen
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health care professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not use more than directed**
- take every 4 hours; not to exceed 6 packets in 24 hours or as directed by a doctor.
- adults and children 12 years of age and over: dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- children under 12 years of age: consult a doctor
- If using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

**Other information**

- each packet contains: potassium 7 mg,  
sodium 46 mg
- phenylketonurics: contains phenylalanine  
13 mg per packet
- store at controlled room temperature  
20-25C (68-77F). Protect from  
excessive heat and moisture.

Acesulfame K, aspartame, caramel color, citric acid, D and C Yellow 10, maltodextrin, natural flavors, sodium citrate, starch and sugar.

**Drug Facts**

**Active ingredients (in each packet)**

Acetaminophen 650 mg ..... Pain reliever/Fever reducer  
 Diphenhydramine hydrochloride 25 mg...Antihistamine/Cough suppressant  
 Phenylephrine hydrochloride 10 mg ..... Nasal decongestant

**Purpose**

**Uses**

- Temporarily relieves:
  - minor aches and pains • headache • minor sore throat pain
  - nasal and sinus congestion • cough due to minor throat and bronchial irritation • runny nose • sneezing • itchy, watery eyes due to hay fever • itchy nose and throat
- Temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take • more than 6 packets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription and nonprescription).
- Ask a doctor or pharmacist before using with other drugs if you are not sure.
- with any other product containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease • heart disease • high blood pressure • thyroid disease
- glaucoma • diabetes • trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema, asthma, or chronic bronchitis
- cough that occurs with too much phlegm (mucus) • cough that lasts or is chronic such as occurs with smoking, asthma or emphysema.

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers • taking the blood thinning drug warfarin

**When using this product**

- do not exceed recommended dosage
- avoid alcoholic drinks • marked drowsiness may occur • alcohol, sedatives and tranquilizers may increase drowsiness • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children.

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occurs • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur
- symptoms do not get better or worsen • pain, cough or nasal congestion gets worse or lasts more than 7 days • cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health care professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not use more than directed
- take every 4 hours; not to exceed 6 packets in 24 hours or as directed by a doctor.
- adults and children 12 years of age and over: dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- children under 12 years of age: consult a doctor.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

**Other information**

- each packet contains: potassium 7 mg, sodium 46 mg
- phenylephrine tartrate: contains phenylalanine 13 mg per packet
- store at controlled room temperature 20-25°C (68-77°F). Protect from excessive heat and moisture.

**Inactive ingredients:**

Acesulfame K, aspartame, caramel color, citric acid, D&C Yellow 10, maltodextrin, natural flavors, sodium citrate, starch and sugar.

**Questions or comments?**  
1-800-925-4733

CONVENIENT RECLOSING TAB

NIGHTTIME  
**Wal-Flu®**  
Severe Cold & Cough

Well at  
*Walgreens*

WALGREENS PHARMACIST RECOMMENDED†

Well at  
*Walgreens*

WALGREENS PHARMACIST RECOMMENDED†

NIGHTTIME  
**Wal-Flu®**  
Severe Cold & Cough

Acetaminophen / Pain Reliever / Fever Reducer  
 Diphenhydramine HCl / Antihistamine / Cough Suppressant  
 Phenylephrine HCl / Nasal Decongestant

**Relieves:**

- Nasal congestion • Cough
- Body ache • Sore throat pain
- Runny nose • Sneezing
- Headache • Fever

HONEY LEMON FLAVOR  
Infused with Chamomile & White Tea Flavors

Compare to Theraflu® Nighttime Severe Cold & Cough active ingredients††

**6 PACKETS**    SEE NEW WARNINGS INFORMATION

†Walgreens Pharmacist Survey Study, November 2012.  
 ††This product is not affiliated with Novartis Consumer Health, Inc., owner of the registered trademark Theraflu®.

DO NOT USE IF SEALED PACKETS ARE TORN OR BROKEN

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015    ITEM 499403

100% SATISFACTION GUARANTEED

walgreens.com © 2013 Walgreen Co. MADE IN CANADA

3 11917 10758 5

C-3109-30508-6W-2    REV0813-RF

**WALGREENS NIGHT TIME WAL-FLU SEVERE COLD AND COUGH HONEY LEMON FLAVOR INFUSED WITH CHAMOMILE AND WHITE TEA FLAVORS**  
 acetaminophen, diphenhydramine hcl, and phenylephrine hcl. granule, for solution

**Product Information**

|                                |                |                           |                |
|--------------------------------|----------------|---------------------------|----------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:0363-110 1 |
| <b>Route of Administration</b> | ORAL           |                           |                |

**Active Ingredient/Active Moiety**

| Ingredient Name  | Basis of Strength             | Strength |
|--|-------------------------------|----------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)                   | ACETAMINOPHEN                 | 650 mg   |
| DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M) | DIPHENHYDRAMINE HYDROCHLORIDE | 25 mg    |

|   |                                    |       |
|---|------------------------------------|-------|
| <b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | <b>PHENYLEPHRINE HYDROCHLORIDE</b> | 10 mg |
|---|------------------------------------|-------|

### Inactive Ingredients

| Ingredient Name                                   | Strength |
|---|----------|
| <b>ACESULFAME</b> (UNII: MA3UYZ6K1H)              |          |
| <b>ASPARTAME</b> (UNII: Z0H242BBR1)               |          |
| <b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP) |          |
| <b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)            |          |
| <b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)          |          |
| <b>SUCROSE</b> (UNII: C151H8M554)                 |          |

### Product Characteristics

|                 |   |                     |  |
|-----------------|---|---------------------|--|
| <b>Color</b>    | yellow (Caramel Color) , yellow (D & C Yellow 10)                           | <b>Score</b>        |  |
| <b>Shape</b>    |   | <b>Size</b>         |  |
| <b>Flavor</b>   | HONEY (Natural Honey & Lemon Flavor) , LEMON (Natural Honey & Lemon Flavor) | <b>Imprint Code</b> |  |
| <b>Contains</b> |   |                     |  |

### Packaging

| # | Item Code        | Package Description                           | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:0363-1101-06 | 6 in 1 BOX; Type 0: Not a Combination Product |                      |                    |

### Marketing Information

| Marketing Category  | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part341                                  | 10/08/2013           |                    |

**Labeler** - Walgreen Co. (008965063)

Revised: 12/2015

Walgreen Co.