# LOPERAMIDE HYDROCHLORIDE- loperamide hydrochloride tablet Valu Merchandisers Company

-----

# Loperamide Hydrochloride

**Drug Facts** 

# **Active ingredient (in each caplet)**

Loperamide Hydrochloride USP, 2 mg

# **Purpose**

Anti-diarrheal

#### Use

controls symptoms of diarrhea, including Travelers' Diarrhea

## Warnings

# Allergy alert

Do not use if you have ever had a rash or other allergic reaction to loperamide hydrochloride

#### Heart alert

Taking more than directed can cause serious heart problems or death

**Do not use** if you have bloody or black stool

## Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product** tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

## Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

#### Directions

# • drink plenty of clear fluids to help prevent dehydration caused by diarrhea

• find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

#### Other information

- store between 20° 25°C (68° 77°F)
- see side panel for lot number and expiration date
- TAMPER EVIDENT: THIS PRODUCT PROTECTED WITH SEALED BLISTER UNITS. DO NOT USE IF ANY ARE TORN OR BROKEN.

# **Inactive ingredients**

anhydrous lactose, croscarmellose sodium, crospovidone, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

#### Questions?

Call toll-free Monday-Friday 8:30 am to 5 pm EST at **1800-406-7984**.

PROUDLY DISTRIBUTED BY: VALU MERCHANDISERS, CO. 5000 KANSAS AVE KANSAS CITY, KS 66106

#### PRINCIPAL DISPLAY PANEL - 12 Caplet Blister Pack Carton

COMPARE TO THE ACTIVE INGREDIENT IN IMODIUM® A-D<sup>†</sup>

Best Choice®

See New Warnings and Directions

Anti-Diarrheal Loperamide Hydrochloride Tablets USP, 2 mg

Anti-Diarrheal Controls the symptoms of diarrhea 12 CAPLETS\*

EACH CAPLET (\*CAPSULE-SHAPED TABLET)
CONTAINS LOPERAMIDE HYDROCHLORIDE USP, 2 mg

.**1867-801-0081** 18 T23 mq 3 of Questions? Call tdl-fee Morday-Friday 8:30 am

cellulose, pregelafinized starch vegetable oil, magne sium steara e, powde red FD&C blue rp. 1 aluminum lake, hydrogenated crospovidone, D&Cyellow no. 10 aluminum lake, amydrous lactose, crosc am elose so dium, staelbergal entre

ANY ARETORN OR BROKEN.

WITH SEALED BLISTER UNITS, DO NOT USE IF TAMPER EN DENT: THIS PRODUCT IS PROTECTED

 see sidep anel for lot number and expiration date ■ store between 20° - 25° C(68° - 77°F)

_		-				•	
ио	де	ш	OĮ.	Щ	IJЭ	47	0

71 110	
children und er2 years (up to 33 lbs)	esn pou op
children 2-5 years (34 10 47 lbs)	ask a doctor
children 6-8 years (49-59 lbs)	1 captetatierthe first loose stoot; ½ captetatier each subsequent loose stoot; but no more han 2 captetsin 24 hours
sheay ff-e nenblido (ad 1 ae-coa)	1 captetathethefinatioose atoot; ½ captetathereach subsequentioosestoot; but no more han 3 captetsin 24 hours
Drug Facts (	couri unea)

S t ponts uo wore than teaplets in s ubsequent looses tool; but

stool; 1 caplet after each 12 years and over adults and children | 2 caplets after the first I cose

to do se; otherwise, use age.

 drink planty of clear theids to help prevent dehydration caused by diarmics
 dehydration caused by diarmics Directions

ñght away (1-800-222-1222) probesional before use Keep out of reach of children, in case of overdose, get medical help or con bed. a Poison Control Certer If pregnant or breast-feeding, æk a health

you get abdominal swelling or bulging. The se may be signs of a serious condition.

ayab ≤ narit enom not atsal serimaib

Stop use and sak a doctor if symptoms getworse

operating machinery diz ziness m ay occur. Be careful when driving or When using this product fredness, drowsinessor

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain pre scription drugs.

■ a history of abnormal heart rhythm

■ a history of live rdise a se

lode and ni subum Aska dod or b d'ore use if y ou have

Do not use it you have bloody or black stool

(couring **Facts** (continued)

Heart alent Taking more than directed can cause serious heart problems or death Allergy alert. Do not use if you have ever had a rash or other aller gic rescript to peramide hydrochloride Warnings

USE controlssymptom sof diarrhea, including Travelers' Diarrhea

Lesetth sib-ifnA əsodınd

Lope ramide Hydrochloride USR, 2 mg .... Active ingredient (in each caplet)

Drug Facts

Loperamide Hydrochloride Tablets

COMPARE TO THE ACTIVE INGREDIENT IN IMODIUM® A-D†

See New Warnings and Directions

Best

# **Anti-Diarrheal**

**Loperamide Hydrochloride Tablets** USP, 2 mg

**Anti-Diarrheal** Controls the symptoms of diarrhea

12 CAPLETS

EACH CAPLET (\* CAPSULE-SHAPED TABLET) CONTAINS LOPERAMIDE HYDROCHLORIDE USP, 2 mg All trademarks are property of their respective owners. This product is not affiliated the makers/owners of Imodium®A-D

Lot No.:



# Keep the carton. It contains important information.

R0419



PROUDLY DISTRIBUTED BY: VALU MERCHANDISERS, CO. 5000 KANSAS AVE KANSAS CITY, KS 66106



# LOPERAMIDE HYDROCHLORIDE

loperamide hydrochloride tablet

# **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:63941-123

Route of Administration ORAL

# **Active Ingredient/Active Moiety**

reave ingreated visitely				
Ingredient Name	Basis of Strength	Strength		
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII: 6 X9 OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg		

Inactive Ingredients				
Ingredient Name	Strength			
ANHYDROUS LACTOSE (UNII: 3S Y5LH9 PMK)				
CROSCARMELLOSE SODIUM (UNII: M28 OL1HH48)				
CROSPO VIDO NE (15 MPA.S AT 5%) (UNII: 68401960 MK)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
HYDRO GENATED COTTONSEED OIL (UNII: Z82Y2C65EA)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
POWDERED CELLULOSE (UNII: SMD1X3XO9M)				
STARCH, CORN (UNII: O8232NY3SJ)				

Product Characteristics				
Color	green	Score	2 pieces	
Shape	CAPSULE	Size	9 mm	
Flavor		Imprint Code	123	
Contains				

]	Packaging					
#	Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>		
1	NDC:63941-123-12	12 in 1 BLISTER PACK; Type 0: Not a Combination Product	02/01/1993			
2	NDC:63941-123-24	24 in 1 BLISTER PACK; Type 0: Not a Combination Product	02/01/1993			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA074091	02/01/1993		

# Labeler - Valu Merchandisers Company (868703513)

# Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment					
Name	Address	ID/FEI	Business Operations		
Ohm Laboratories Inc.		051565745	manufacture(63941-123)		

Revised: 5/2019 Valu Merchandisers Company