

VICKS PAINQUIL AND VICKS PAINQUIL PM PAIN RELIEVER- acetaminophen, diphenhydramine hcl

The Procter & Gamble Manufacturing Company

**Vicks ® PainQuil™ / Vicks ® PainQuil™ PM
PAIN RELIEVER, Honey Flavored Convenience Pack**

Vicks ® PainQuil™ PM PAIN RELIEVER + NIGHTTIME SLEEP-AID

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 1000 mg

Diphenhydramine HCl 50 mg

Purpose

Pain reliever

Nighttime sleep-aid

Uses

- for the temporary relief of occasional minor aches and pains with accompanying sleeplessness.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 3 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- for children under 12 years of age
- with any other product containing diphenhydramine, even one used on skin

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness. These could be signs of a serious condition.

When using this product

- avoid alcoholic beverages
- drowsiness will occur
- do not drive a motor vehicle or operate machinery

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only one dose (30 mL) per day (24 hours)
- only use the dose cup provided
- only use as directed

adults & children 12 yrs & over	30 mL at bedtime
children under 12 yrs	do not use

Other information

- **each 30 mL contains:** sodium 21 mg
- do not exceed 25°C (77°F) and do not refrigerate.

Inactive ingredients

alcohol, anhydrous citric acid, D&C Yellow No.10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No.6, flavor, glycerin, polysorbate 20, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water

Questions?

1-877-881-5813

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND IS BROKEN OR MISSING.

**DIST. BY: PROCTER & GAMBLE,
CINCINNATI, OH 45202**

Vicks ® PainQuil™ PAIN RELIEVER

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 1000 mg

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains associated with:

- sore throat
- headache
- muscular aches
- backache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 3 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- liver disease
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur. These could be signs of a serious condition.

If pregnant or breastfeeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 3 doses (30 mL each) per day (24 hours)
- only use the dose cup provided

adults & children 12 yrs & over	30 mL every 6 hours
children under 12 yrs	do not use

Other information

- **each 30 mL contains:** sodium 21 mg
- do not exceed 25°C (77°F) and do not refrigerate.

Inactive ingredients

alcohol, anhydrous citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water

Questions?

1-877-881-5813

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND IS BROKEN OR MISSING.

**DIST. BY: PROCTER & GAMBLE,
CINCINNATI, OH 45202**

PRINCIPAL DISPLAY PANEL - Convenience Pack

Vicks [®]PainQuil[™] PAIN RELIEVER / Vicks [®]PainQuil[™] PM PAIN RELIEVER + NIGHTTIME SLEEP-AID
MAX STRENGTH‡

VALUE PACK

‡Maximum strength dose of active ingredients per dosing interval, only use as directed.

Vicks [®]PainQuil[™] PAIN RELIEVER

STARTS WORKING FAST FOR RELIEF OF:

ACHES | PAINS

HEADACHE

SORE THROAT

HONEY FLAVORED

Alcohol 10%

Vicks [®]PainQuil[™] PM PAIN RELIEVER + NIGHTTIME SLEEP-AID

STARTS WORKING FAST FOR RELIEF OF:

ACHES | PAINS

HEADACHE
SORE THROAT
+

OCCASSIONAL SLEEPLESSNESS
HONEY FLAVORED

Alcohol 10%

2 BOTTLES - 1 PAINQUIL/1 PAINQUIL PM 12 FL OZ (354 mL) EACH; TOTAL 24 FL OZ (708 mL)



VICKS PAINQUIL AND VICKS PAINQUIL PM PAIN RELIEVER			
acetaminophen, diphenhydramine hcl kit			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84126-351
Packaging			

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84126-351-24	1 in 1 PACKAGE; Type 0: Not a Combination Product	05/06/2025	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	354 mL
Part 2	1 BOTTLE	354 mL

Part 1 of 2

VICKS PAINQUIL PAIN RELIEVER

acetaminophen liquid

Product Information

Item Code (Source)	NDC:84126-349
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	1000 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
ALCOHOL (UNII: 3K9958V90M)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
WATER (UNII: 059QF0KO0R)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Product Characteristics				
Color		brown	Score	
Shape			Size	
Flavor		HONEY	Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84126-349-12	354 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M013	05/07/2024	
Part 2 of 2				
VICKS PAINQUIL PM PAIN RELIEVER PLUS NIGHTTIME SLEEP-AID				
acetaminophen, diphenhydramine hcl liquid				
Product Information				
Item Code (Source)		NDC:84126-350		
Route of Administration		ORAL		
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)			ACETAMINOPHEN	1000 mg in 30 mL
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)			DIPHENHYDRAMINE HYDROCHLORIDE	50 mg in 30 mL
Inactive Ingredients				
Ingredient Name				Strength
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
POLYSORBATE 20 (UNII: 7T1F30V5YH)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
WATER (UNII: 059QF0KO0R)				

FD&C RED NO. 40 (UNII: WZB9127XOA)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
ALCOHOL (UNII: 3K9958V90M)	
GLYCERIN (UNII: PDC6A3C0OX)	

Product Characteristics

Color	brown	Score	
Shape		Size	
Flavor	HONEY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84126-350-12	354 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/07/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/06/2025	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 4/2025

The Procter & Gamble Manufacturing Company