

MAXIMUM STRENGTH 4% LIDOCAINE PATCH- lidocaine patch
InterMed Laboratories Private Limited

Maximum Strength 4% Lidocaine Patch

Active Ingredient

Lidocaine 4%

Purpose

Topical Anesthetic

Uses

Temporarily relieves minor pain

Warnings

For external use only

Do Not Use

- more than 1 patch on your body at a time or on cut, irritated or swollen skin
- on puncture wounds
- for more than one week without consulting a doctor
- over raw surfaces or blistered areas
- if you are allergic to any active or inactive ingredients
- if pouch is damaged or opened

When Using This Product

- use only as directed. Read and follow all directions and warnings on this label.
- do not allow contact with the eyes
- do not bandage tightly or apply local heat (such as heating pads) to the area of use
- do not use at the same time as other topical analgesics
- dispose of used patch in manner that always keeps product away from children or pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop Use and Ask a Doctor if

- condition worsens
- redness is present
- irritation develops
- symptoms persist for more than 7 days or clear up and occur again within a few

days

- you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied

If Pregnant or Breast Feeding

ask a health professional before use.

Keep Out of Reach of Children and Pets

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and children over 12 years:

- clean and dry affected area free of lotions, ointments and creams.
- carefully remove backing from patch starting at a corner. Apply sticky side of patch to affected area.
- Do not use more than one patch in a 12 hour period. Maximum 2 patches per day. Discard patch after single use.

children 12 years or younger:ask a physician.

Directions

Adults and children over 12 years:

- clean and dry affected area
- remove backing from patch by firmly grasping both ends and gently pulling until backing separates in middle
- carefully remove smaller portion of backing from patch and apply exposed portion of patch to affected area
- once exposed portion of patch is positioned, carefully remove remaining backing to completely apply patch to affected area
- use 1 patch for up to 12 hours

children 12 years or younger:ask a doctor

Inactive Ingredients

dihydroxy aluminium aminoacetate, disodium EDTA, glycerin, methyl paraben, polyvinyl alcohol, propyl paraben, propylene glycol, sodium carboxy methyl cellulose, sodium polyacrylate, tartaric acid, titanium dioxide, water

Bulk Package Label


CUT OPEN POUCH AND REMOVE PATCH



by Veridian Healthcare

*Compare to active ingredient in
Aspercreme® Lidocaine Pain Relief Patch

Maximum OTC Strength
4% Lidocaine
Pain Relief Patch


Temporary relief
of minor pain

TOPICAL ANESTHETIC


- Provides Temporary Pain Relief
- Flexible Adhesive Patch
- Single-Use Application
- Unscented Formula





1 PATCH
3.93 IN. X 5.51 IN. (10 cm x 14 cm)



CUT OPEN POUCH AND REMOVE PATCH


Model 24-911


PULL TO SEPARATE FILM


PEEL OFF ONE SIDE OF FILM


PLACE PATCH ON SKIN


PEEL OFF OTHER SIDE OF FILM


PRESS FIRMLY TO SKIN

Drug Facts

Active ingredient
Lidocaine 4.0%

Purpose
..... Topical anesthetic

Use For temporary relief of pain.

Warnings
For external use only

Do not use ■ More than one patch on your body at a time ■ On cut, irritated or swollen skin ■ On puncture wounds ■ Over raw surfaces or blistered areas ■ For more than one week without consulting a doctor ■ If you are allergic to any active or inactive ingredients ■ If pouch is damaged or opened.

When using this product ■ Use only as directed ■ Read and follow all directions and warnings on this carton ■ Do not allow contact with the eyes ■ Do not use at the same time as other topical anesthetics ■ Do not bandage tightly or apply local heat (such as heating pads) to the area of use ■ Do not microwave ■ Dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop use and consult a doctor if ■ Condition worsens ■ Redness is present ■ Irritation develops ■ Symptoms persist for more than 7 days or clear up and occur again within a few days ■ You experience signs of skin injury, such as pain, swelling or blistering where the product was applied.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children and pets. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
Adults and children 12 years of age and over: ■ Clean and dry affected area, free of lotions, ointments and creams. Carefully remove backing from patch starting at a corner. Apply sticky side of patch to affected area. Do not use more than one patch in a 12 hour period. Maximum 2 patches per day. Discard patch after single use.
Children under 12 years of age: ■ consult a physician.

Other information Store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.

Inactive ingredients dihydroxy aluminum aminoacetate, disodium EDTA, glycerin, methyl paraben, polyvinyl alcohol, propyl paraben, propylene glycol, sodium carboxy methyl cellulose, sodium polyacrylate, tartaric acid, titanium dioxide, water

*This product is not manufactured or distributed by Chatterm, Inc., owner of the registered trademark Aspercreme®

Manufactured for:
Veridian Healthcare, LLC
11751 Lakeside Drive
Gurnee, IL 60031 USA
www.VeridianHealthcare.com
Made in India

Child-Resistant Packaging

TOLL-FREE
Customer Care Help Line
866-326-1313
Mon-Fri 8:30 am-4:30 pm CST

84-B-W0225/PC14901-00

Code No: TN/DRUGS/22/1999
LOT
EXP

MAXIMUM STRENGTH 4% LIDOCAINE PATCH

lidocaine patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71811-004
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	40 mg

Inactive Ingredients

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE CALCIUM (UNII: UTY7PDF93L)	
TARTARIC ACID (UNII: W4888I119H)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
DIHYDROXYALUMINUM AMINOACETATE (UNII: DO250MG0W6)	
GLYCERIN (UNII: PDC6A3C0OX)	
EDETATE DISODIUM ANHYDROUS (UNII: 8NLQ36F6MM)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
SODIUM POLYACRYLATE (2500000 MW) (UNII: 05I15JNI2J)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
WATER (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71811-004-05	5 in 1 CARTON	04/29/2025	
1		1 in 1 POUCH; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M017	04/29/2025	

Labeler - InterMed Laboratories Private Limited (676244169)