MUCINEX RAPID CLEAR PAIN, HEADACHE, AND MUCUS CONGESTION AND SINUS MAX CLEAR AND COOL- oxymetazoline hydrochloride, acetaminophen and guaifenesin RB Health (US) LLC

MUCINEX® RAPID+CLEAR Pain Headache & Mucus Congestion + MUCINEX® SINUS-MAX® Severe Nasal Congestion Relief CLEAR & COOL Nasal Spray

Active ingredient Purpose Oxymetazoline hydrochloride 0.05%.....Nasal decongestant

Active ingredients (in each capsule) Purposes Acetaminophen 325 mg......Pain reliever/fever reducer Guaifenesin 200 mg.....Expectorant

Uses

- temporarily relieves nasal congestion due to:
- \blacksquare a cold \blacksquare hay fever or other upper respiratory allergies
- promotes nasal and/or sinus drainage; temporarily relieves sinus congestion and pressure
- helps clear nasal passages; shrinks swollen membranes

Uses

- temporarily relieves these common cold and flu symptoms:
- \blacksquare minor aches and pains \blacksquare headache \blacksquare sore throat
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways
- of bothersome mucus and make coughs more productive **I** temporarily reduces fever

Warnings

Ask a doctor before use if you have

- heart disease high blood pressure
- thyroid disease diabetes
- difficulty in urination due to enlargement of the prostate gland

When using this product

- do not exceed recommended dosage
- do not use this product for more than 3 days. Use only as directed. Frequent or prolonged use

may cause nasal congestion to recur or worsen.

■ this product may cause temporary discomfort such as burning, stinging, sneezing, or an

increase in nasal discharge

■ the use of this container by more than one person may spread infection

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 capsules in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or

followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use

 \blacksquare with any other drug containing acetaminophen (prescription or nonprescription). If you are not

sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Directions

■ adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each

nostril, not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.

■ children under 6 years of age: consult a doctor.

Shake well before use.

To open, hold by the white grips then squeeze, push down firmly and turn cap counterclockwise.

Before using for the first time, remove the protective cap from the tip and prime metered pump

by depressing firmly several times. To spray, hold bottle with thumb at the base and nozzle

between first and second fingers.Without tilting head, insert nozzle into nostril. Fully depress

pump all the way down with a firm even stroke and sniff deeply. Wipe nozzle clean after use.

To close, turn cap clockwise. DO NOT DISCARD CAP.

Directions

■ do not take more than directed (see Overdose warning)

■ do not take more than 12 capsules in any 24-hour period

■ adults and children 12 years of age and over: take 2 capsules every 4 hours

■ children under 12 years of age: do not use

Other information

■ store at 20-25°C (68-77°F)

avoid excessive heat

Inactive ingredients

benzalkonium chloride, benzododecinium chloride, camphor, cetalkonium chloride, colloidal

silicon dioxide, edetate disodium, eucalyptol, glycine, linoleic acid, linolenic acid, menthol, myristalkonium chloride, myristic acid, oleic acid, palmitic acid, palmitoleic acid, polyethylene

glycol, polysorbate 80, propylene glycol, purified water, sodium carbonate, sodium chloride,

sodium hydroxide, stearic acid

Inactive ingredients

FD&C blue no.1, gelatin, glycerin, hypromellose, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

Keep out of reach of Children

Purpose

Oxymetazoline hydrochloride 0.05%.....Nasal decongestant

Acetaminophen 325 mg.....Pain reliever/fever reducer Guaifenesin 200 mg.....Expectorant



Severe Nasal Congestion Relief Clear & Cool Nasal Spray Drug Facts	Pain, Headache & Mucus Congestion			
-				
Active ingredient Purpose	Active ingredients (in each capsule) Purpos			
Oxymetazoline hydrochloride 0.05%Nasal decongestant	Acetaminophen 325 mgPain reliever/fever reduc Guaifenesin 200 mgExpectora			
Uses temporarily relieves nasal congestion due to: a cold hay fever or other upper respiratory allergies promotes nasal and/or sinus drainage; temporarily relieves sinus congestion and pressure helps clear nasal passages; shrinks swollen membranes	Uses temporarily relieves these common cold and flu symptoms: minor aches and pains headache sore throat helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageway of bothersome mucus and make couchs more productive temporarily reduces fever			
Warnings				
Ask a doctor before use if you have heart disease high blood pressure thyroid disease diabetes difficulty in urination due to enlargement of the prostate gland	Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you tai more than 12 capsules in 24 hours, which is the maximum daily amount for this product with other drugs containing acetaminophen 2 or over a clockeling data data with which which the product			
When using this product do not exceed recommended dosage do not use this product for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen. this product may cause temporary discomfort such as burning, stinging, sneezing, or an increase in nasal discharge	 3 or more alcoholic drinks daily while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening blisters rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warming: If sore throat is severe, persists for more than 2 days, is accompanied o followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use 			
the use of this container by more than one person may spread infection Stop use and ask a doctor if symptoms persist	with any other drug containing acetaminophen (prescription or nonprescription). If you are n sure whether a drug contains acetaminophen, ask a doctor or pharmacist.			
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	Ask a doctor before use if you have liver disease persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphyser cough that occurs with too much phlegm (mucus)			
Directions	Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin			
adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril, not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.	When using this product do not use more than directed			
children under 6 years of age: consult a doctor. Shake well before use. To open, hold by the white grips then squeeze, push down firmly and turn cap counterclockwise. Before using for the first time, remove the protective cap from the tip and prime metered pump	Stop use and ask a doctor if pain or cough gets worse or lasts more than 7 days If fever gets worse or lasts more than 3 days redness or swelling is present Icough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. Image: Comes back or occurs			
by depressing firmly several times. To spray, hold bottle with thumb at the base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress pump all the way down with a firm even stroke and sniff deeply. Wipe nozzle clean after use. To close, turn cap clockwise. DO NOT DISCARD CAP.	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cau liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.			
Other information ■ store at 20-25°C (68-77°F)	Directions ■ do not take more than directed (see Overdose warning)			
Inactive ingredients benzalkonium chloride, benzododecinium chloride, camphor, cetalkonium chloride, colloidal silicon dioxide, edetate disodium, eucalvotol, olycine, linoleic acid, linolenic acid, menthol,	do not take more than 12 capsules in any 24-hour period adults and children 12 years of age and over: take 2 capsules every 4 hours children under 12 years of age: do not use			
myristalkonium chloride, myristic acid, ôleic acid, palmitic acid, palmitoleic acid, polyethylene glycol, polysorbate 80, propylene glycol, purified water, sodium carbonate, sodium chloride,	Other information ■ store at 20-25°C (68-77°F) ■ avoid excessive heat			
sódium hydroxide, stearic acid Questions? 1-866-MUCINEX (1-866-682-4639)	<i>Inactive ingredients</i> FD&C blue no.1, gelatin, glycerin, hypromellose, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide			
	Questions? 1-866-MUCINEX (1-866-682-4639)			

MUCINEX RAPID CLEAR PAIN, HEADACHE, AND MUCUS CONGESTION AND SINUS MAX CLEAR AND COOL

oxymetazoline hydrochloride, acetaminophen and guaifenesin kit

Ρ	Product Information							
P	roduct Type	HUMAN OTC DRUG	ltem Cod	e (Source)	NDC:72854-216			
D								
Packaging								
#	ltem Code	Package Description	Package Description		Marketing End Date			
1	NDC:72854-216- 02	1 in 1 CARTON	n 1 CARTON					
1		1 in 1 CARTON; Type 0: Not a Com	bination					

Ŧ			Product						
Q	uant	ity of P	arts						
Part # Package Quantity			Total Product Quantity						
Pa				1 mL					
Pa	rt 2	0 BLISTER	R PACK		1				
Ρ	art	1 of 2							
			SINUS-M NASAL	AX SEVERE N	ASAL C	ON	GESTION RE	LIEF	CLEAR
0>	ymet	azoline l	hydrochlorid	e solution					
Ρ	rodu	ct Info	rmation						
lte	em Co	ode (Sou	irce)	NDC:63824-129					
Ro	oute	of Admin	istration	NASAL					
A	tive	Ingred	lient/Active	e Moiety					
			Ingre	dient Name			Basis of Stre	ngth	Strength
			HYDROCHLO UNII:8VLN5B4	RIDE (UNII: K89MJ0S5VY))		OXYMETAZ OLINE HYDROCHLORIDE		0.05 g in 100 mL
(0,		AZ ULINE -		+2 1)			HIDROCHLORIDE		
In	activ	ve ingre	edients						
				Ingredient Nam	е			9	Strength
				III: F5UM2KM3W7)					
		-	RAL) (UNII: N2						
BE	NZOC	DODECINI	UM CHLORIDI	(UNII: Y5A751G47H)					
Pı	odu	ct Char	acteristics						
Co	lor				Score				
Sh	ape				Size				
	vor			MENTHOL	Imprin	t Cod	de		
Co	ntair	าร							
Pa	acka	ging							
#	lten	n Code	F	ackage Descriptio	n	М	larketing Start	Mar	keting End
	NDC:6	3824-	1 in 1 CARTON				Date		Date
_	129-1	7			ta				
1			22 mL in 1 BOTTLE, PUMP; Type 0: Not a Combination Product						

Marketing In	format	ion					
Marketing			er or Monograph	Marke	eting Start	Marke	ting End
Category	Аррпси	Citat			Date		Date
OTC Monograph Drug	M012			05/01/20	25		
Part 2 of 2							
MUCINEX SI							
acetaminophen an							
acetainiinophen an	iu gualierie	esin capsui					
Product Inform	ation						
ltem Code (Source	e)	NDC:72854-2	211				
Route of Administ	ration	ORAL					
Active Ingredier	t/Activo	Maiaty					
······	IL/ACTIVE	Molety					
		redient Nar	ne		Basis of St	rength	Strengt
GUAIFENESIN (UNII: 4	Ingr	edient Nar			Basis of St GUAIFENESIN	rength	Strength 200 mg
GUAIFENESIN (UNII: 4	ing r 95W7451VQ	edient Nar) (GUAIFENESI	N - UNII:495W7451VQ)	ГL9D)		-	
-	ing r 95W7451VQ	edient Nar) (GUAIFENESI	N - UNII:495W7451VQ)	ГL9D)	GUAIFENESIN	-	200 mg
GUAIFENESIN (UNII: 4	ing r 95W7451VQ NII: 36209ITI	edient Nar) (GUAIFENESI	N - UNII:495W7451VQ)	FL9D)	GUAIFENESIN	-	200 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UN	ing r 95W7451VQ NII: 36209ITI	edient Nar) (GUAIFENESI L9D) (ACETAMI	N - UNII:495W7451VQ)	FL9D)	GUAIFENESIN	EN	200 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UN	Ing n 95W7451VQ NII: 36209ITI ents	edient Nar) (GUAIFENESI L9D) (ACETAMI	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ⁻	FL9D)	GUAIFENESIN	EN	200 mg 325 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UNII: 1 Inactive Ingredi	Ing r 95W7451VQ NII: 36209ITL ents	edient Nar) (GUAIFENESI L9D) (ACETAMI	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ⁻	FL9D)	GUAIFENESIN	EN	200 mg 325 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UN Inactive Ingredi GELATIN (UNII: 2G86Q	Ing r 95W7451VQ NII: 36209ITL ents N327L) 5A3C0OX)	edient Nan) (GUAIFENESI L9D) (ACETAMI Ingredia	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ⁻ ent Name	ΓL9D)	GUAIFENESIN	EN	200 mg 325 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UNII: 4 Inactive Ingredi GELATIN (UNII: 2G86Q GLYCERIN (UNII: PDC6 HYPROMELLOSE, UN POLYETHYLENE GLYC	Ing r 95W7451VQ NII: 36209ITL ents N327L) SA3C0OX) SPECIFIED COL, UNSPI	edient Nan) (GUAIFENESI L9D) (ACETAMI Ingredia (UNII: 3NXW29 ECIFIED (UNII	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ⁻ ent Name IV3WO)	FL9D)	GUAIFENESIN	EN	200 mg 325 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UNII: 4 Inactive Ingredi GELATIN (UNII: 2G86Q GLYCERIN (UNII: PDC6 HYPROMELLOSE, UN POLYETHYLENE GLYCO	Ing r 95W7451VQ NII: 362O9ITI ents N327L) 5A3C0OX) SPECIFIED COL, UNSPI	edient Nar) (GUAIFENESI L9D) (ACETAMI Ingredie (UNII: 3NXW29 ECIFIED (UNII Q167V3)	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ⁻ ent Name IV3WO)	ΓL9D)	GUAIFENESIN	EN	200 mg 325 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UNII: 4 Inactive Ingredi GELATIN (UNII: 2G86Q GLYCERIN (UNII: PDC6 HYPROMELLOSE, UN POLYETHYLENE GLYCOL FD&C BLUE NO. 1 (U	Ing r 95W7451VQ NII: 36209ITL ents N327L) 5A3C00X) SPECIFIED COL, UNSPI (UNII: 6DC9 NII: H3R47K3	edient Nan) (GUAIFENESI L9D) (ACETAMI Ingredie (UNII: 3NXW29 ECIFIED (UNII Q167V3) 3TBD)	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ⁻ ent Name IV3WO)	ΓL9D)	GUAIFENESIN	EN	200 mg 325 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UNII: 4 Inactive Ingredi GELATIN (UNII: 2G86Q GLYCERIN (UNII: PDC6 HYPROMELLOSE, UN POLYETHYLENE GLYCO FD&C BLUE NO. 1 (U LECITHIN, SUNFLOW	Ing r 95W7451VQ NII: 362O9ITI ents N327L) 5A3C0OX) SPECIFIED COL, UNSPI (UNII: 6DC9 NII: H3R47K3 ZER (UNII: 83	edient Nan) (GUAIFENESI L9D) (ACETAMI Ingredie (UNII: 3NXW29 ECIFIED (UNII Q167V3) 3TBD)	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ⁻ ent Name IV3WO)	FL9D)	GUAIFENESIN	EN	200 mg 325 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UNII: 4 Inactive Ingredi GELATIN (UNII: 2G86Q GLYCERIN (UNII: PDC6 HYPROMELLOSE, UN POLYETHYLENE GLYCOL FD&C BLUE NO. 1 (U LECITHIN, SUNFLOW POVIDONE (UNII: FZ98	Ing r 95W7451VQ NII: 362O9ITI ents N327L) 5A3C0OX) SPECIFIED COL, UNSPI (UNII: 6DC9 NII: H3R47K3 (ER (UNII: 83 89GH94E)	edient Nan) (GUAIFENESI L9D) (ACETAMI Ingredie (UNII: 3NXW29 ECIFIED (UNII Q167V3) 3TBD) 94K0WOS5G)	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ⁻ ent Name IV3WO)	FL9D)	GUAIFENESIN	EN	200 mg 325 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UNII: 4 Inactive Ingredi GELATIN (UNII: 2G86Q GLYCERIN (UNII: PDC6 HYPROMELLOSE, UN POLYETHYLENE GLYCOL FD&C BLUE NO. 1 (U LECITHIN, SUNFLOW POVIDONE (UNII: FZ 98 SORBITOL SOLUTION	Ing r 95W7451VQ NII: 362O9ITL ents N327L) 5A3C0OX) SPECIFIED COL, UNSPI (UNII: 6DC9 NII: H3R47K3 7ER (UNII: 83 89GH94E) N (UNII: 8KW3	edient Nan) (GUAIFENESI L9D) (ACETAMI Ingredia (UNII: 3NXW29 ECIFIED (UNII Q167V3) 3TBD) 34K0WOS5G) 3E20702)	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ⁻ ent Name IV3WO)	ΓL9D)	GUAIFENESIN	EN	200 mg 325 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UNII: 4 Inactive Ingredi GELATIN (UNII: 2G86Q GLYCERIN (UNII: PDC6 HYPROMELLOSE, UN POLYETHYLENE GLYCOL FD&C BLUE NO. 1 (U LECITHIN, SUNFLOW POVIDONE (UNII: FZ98 SORBITOL SOLUTION LIGHT MINERAL OIL (Ing r 95W7451VQ NII: 362O9ITI ents N327L) 6A3C0OX) SPECIFIED COL, UNSPI (UNII: 6DC9 NII: H3R47K3 (ER (UNII: 83 89GH94E) N (UNII: 8KW3 (UNII: N6K57	edient Nan) (GUAIFENESI L9D) (ACETAMI Ingredia (UNII: 3NXW29 ECIFIED (UNII Q167V3) 3TBD) 34K0WOS5G) 3E207O2) 87QVP)	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ⁻ ent Name IV3WO)	FL9D)	GUAIFENESIN	EN	200 mg 325 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UNII: 4 Inactive Ingredi GELATIN (UNII: 2G86Q GLYCERIN (UNII: PDC6 HYPROMELLOSE, UN POLYETHYLENE GLYCOL FD&C BLUE NO. 1 (U LECITHIN, SUNFLOW POVIDONE (UNII: FZ98 SORBITOL SOLUTION LIGHT MINERAL OIL (TITANIUM DIOXIDE (U	Ing r 95W7451VQ NII: 362O9ITL ents N327L) 5A3C0OX) SPECIFIED COL, UNSPI (UNII: 6DC94 NII: H3R47K3 (ER (UNII: 838 89GH94E) N (UNII: 8KW3 (UNII: 15FIX9V	edient Nan) (GUAIFENESI L9D) (ACETAMI Ingredie (UNII: 3NXW29 ECIFIED (UNII Q167V3) 3TBD) 34K0WOS5G) 3E207O2) 87QVP)	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ⁻ ent Name IV3WO)	ΓL9D)	GUAIFENESIN	EN	200 mg 325 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UNII: 4 Inactive Ingredi GELATIN (UNII: 2G86Q GLYCERIN (UNII: PDC6 HYPROMELLOSE, UN POLYETHYLENE GLYCOL FD&C BLUE NO. 1 (U LECITHIN, SUNFLOW POVIDONE (UNII: FZ98 SORBITOL SOLUTION LIGHT MINERAL OIL (Ing r 95W7451VQ NII: 362O9ITL ents N327L) 5A3C0OX) SPECIFIED COL, UNSPI (UNII: 6DC94 NII: H3R47K3 (ER (UNII: 838 89GH94E) N (UNII: 8KW3 (UNII: 15FIX9V	edient Nan) (GUAIFENESI L9D) (ACETAMI Ingredie (UNII: 3NXW29 ECIFIED (UNII Q167V3) 3TBD) 34K0WOS5G) 3E207O2) 87QVP)	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ⁻ ent Name IV3WO)		GUAIFENESIN	EN	200 mg 325 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UNII: 4 Inactive Ingredi GELATIN (UNII: 2G86Q GLYCERIN (UNII: PDC6 HYPROMELLOSE, UN POLYETHYLENE GLYCOL FD&C BLUE NO. 1 (U LECITHIN, SUNFLOW POVIDONE (UNII: FZ98 SORBITOL SOLUTION LIGHT MINERAL OIL (TITANIUM DIOXIDE (U	Ing r 95W7451VQ NII: 362O9ITL ents N327L) 5A3C0OX) SPECIFIED COL, UNSPI (UNII: 6DC94 NII: H3R47K3 (ER (UNII: 838 89GH94E) N (UNII: 8KW3 (UNII: 15FIX9V	edient Nan) (GUAIFENESI L9D) (ACETAMI Ingredie (UNII: 3NXW29 ECIFIED (UNII Q167V3) 3TBD) 34K0WOS5G) 3E207O2) 87QVP)	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ⁻ ent Name IV3WO)	ΓL9D)	GUAIFENESIN	EN	200 mg 325 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UNII: 4 Inactive Ingredi GELATIN (UNII: 2G86Q GLYCERIN (UNII: PDC6 HYPROMELLOSE, UN POLYETHYLENE GLYCOL FD&C BLUE NO. 1 (U LECITHIN, SUNFLOW POVIDONE (UNII: FZ98 SORBITOL SOLUTION LIGHT MINERAL OIL (TITANIUM DIOXIDE (U	Ing r 95W7451VQ NII: 362O9ITI ents N327L) 5A3C0OX) SPECIFIED COL, UNSPI (UNII: 6DC9 NII: H3R47K3 (UNII: 6DC9 NII: H3R47K3 (UNII: 8573 89GH94E) (UNII: 86K573 UNII: 15FIX9V KOOR)	edient Nan) (GUAIFENESI L9D) (ACETAMI Ingredie (UNII: 3NXW29 ECIFIED (UNII Q167V3) 3TBD) 34K0WOS5G) 3E207O2) 87QVP)	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ⁻ ent Name IV3WO)		GUAIFENESIN	EN	200 mg 325 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UNII: 4 ACETAMINOPHEN (UNII: 2G86Q GLYCERIN (UNII: 2G86Q GLYCERIN (UNII: 2G86Q GLYCERIN (UNII: PDC6 HYPROMELLOSE, UN POLYETHYLENE GLYCOL FD&C BLUE NO. 1 (U LECITHIN, SUNFLOW POVIDONE (UNII: FZ98 SORBITOL SOLUTION LIGHT MINERAL OIL (UNII: FZ98) SORBITOL SOLUTION LIGHT MINERAL OIL (UNII: 059QF0)	Ing r 95W7451VQ NII: 362O9ITI ents N327L) 5A3C0OX) SPECIFIED COL, UNSPI (UNII: 6DC9 NII: H3R47K3 (UNII: 6DC9 NII: H3R47K3 (UNII: 8573 89GH94E) (UNII: 86K573 UNII: 15FIX9V KOOR)	edient Nar) (GUAIFENESI L9D) (ACETAMI Ingredia (UNII: 3NXW29 ECIFIED (UNII Q167V3) 3TBD) 34K0WOS5G) 3E207O2) 87QVP) (2JP)	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ⁻ ent Name IV3WO)		GUAIFENES IN ACETAMINOPHI	EN	200 mg 325 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UNII: 4 ACETAMINOPHEN (UNII: 2G86Q GLYCERIN (UNII: 2G86Q GLYCERIN (UNII: PDC6 HYPROMELLOSE, UN POLYETHYLENE GLYCOL FD&C BLUE NO. 1 (U LECITHIN, SUNFLOW POVIDONE (UNII: FZ98 SORBITOL SOLUTION LIGHT MINERAL OIL (UNII: FZ98 SORBITOL SOLUTION LIGHT MINERAL OIL (UNII: 059QF01) PRODUCT Charact	Ing r 95W7451VQ NII: 362O9ITI ents N327L) 5A3C0OX) SPECIFIED COL, UNSPI (UNII: 6DC9 NII: H3R47K3 (UNII: 6DC9 NII: H3R47K3 (UNII: 6DC9 NII: H3R47K3 (UNII: 6DC9 NII: 15FIX9V (UNII: 8KW3 (UNII: 8KW3 (UNII: 8KW3 (UNII: 15FIX9V KOOR) teristics	edient Nar) (GUAIFENESI L9D) (ACETAMI Ingredia (UNII: 3NXW29 ECIFIED (UNII Q167V3) 3TBD) 34K0WOS5G) 3E207O2) 87QVP) (2JP)	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ent Name NV3WO) : 3WJQ0SDW1A)		GUAIFENES IN ACETAMINOPHI	EN S S S S S S S S S S S S S S S S S S S	200 mg 325 mg
GUAIFENESIN (UNII: 4 ACETAMINOPHEN (UNII: 4 ACETAMINOPHEN (UNII: 2G86Q GLYCERIN (UNII: 2G86Q GLYCERIN (UNII: PDC6 HYPROMELLOSE, UN POLYETHYLENE GLYCOL FD&C BLUE NO. 1 (U LECITHIN, SUNFLOW POVIDONE (UNII: FZ98 SORBITOL SOLUTION LIGHT MINERAL OIL (TITANIUM DIOXIDE (U WATER (UNII: 059QF0) Product Charact Color	Ing r 95W7451VQ NII: 362O9ITI ents N327L) 5A3C0OX) SPECIFIED COL, UNSPI (UNII: 6DC9 NII: H3R47K3 (UNII: 6DC9 NII: H3R47K3 (UNII: 6DC9 NII: H3R47K3 (UNII: 6DC9 NII: 15FIX9V (UNII: 8KW3 (UNII: 8KW3 (UNII: 8KW3 (UNII: 15FIX9V KOOR) teristics	edient Nan) (GUAIFENESI L9D) (ACETAMI Ingredia (UNII: 3NXW29 ECIFIED (UNII Q167V3) 3TBD) 34K0WOS5G) 3E207O2) 87QVP) /2JP)	N - UNII:495W7451VQ) NOPHEN - UNII:362O9I ent Name V/3WO) : 3WJQ0SDW1A) Score		GUAIFENES IN ACETAMINOPHI	EN Socre	200 mg 325 mg

Packaging						
# Item Code	Package Description	Marketing Start Date	Marketing End Date			
1 NDC:72854-211-	1 in 1 CARTON					
	3 in 1 BLISTER PACK; Type 0: Not a Combination Product					
Marketing Information						
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
OTC Monograph Drug	M012	05/01/2025				
Marketing Information						
	Application Number or Monograph	Marketing Start	Marketing End			
Marketing Category	Citation	Date	Date			

Labeler - RB Health (US) LLC (081049410)

Revised: 4/2025

RB Health (US) LLC