

ACETAMINOPHEN- acetaminophen tablet
AARNA USA INC.

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain Reliever/Fever Reducer

Uses

To reduce fever and for the temporary relief of minor aches and pains due to:

- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps.
- Temporarily reduces fever.

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if the user has ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if the user has

- has liver disease

- is a child with pain of arthritis

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if:

- pain gets worse or lasts more than 10 days (for adults) or 5 days (for children)
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose Warning:

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away.(1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

- **do not take more than directed (see overdose warning)**

adults & children 12 years and over	<ul style="list-style-type: none"> • take 2 tablets every 4 to 6 hours while symptoms last • do not take more than 10 tablets in 24 hours • do not use for more than 10 days unless directed by a doctor
children 6-11 years	<ul style="list-style-type: none"> • take 1 tablet every 4 to 6 hours while symptoms last • do not take more than 5 tablets in 24 hours • do not use for more than 5 days unless directed by a doctor
child under 6 years	ask a doctor

Other information

- store at 20° to 25°C (68° to 77°F)

Inactive Ingredients:

- Gelatinized starch, magnesium stearate, povidone


Questions or comments?

call toll-free 1-877-225-6999.

Manufactured for:

AARNA USA INC.

* This product is not manufactured or distributed by Johnson and Johnson, consumer inc., distributor of regular Tylenol Tablets.



NDC 82568-0001-1


Acetaminophen

325MG TABLETS

REGULAR STRENGTH

ASPIRIN FREE
PAIN RELIEVER &
FEVER RELIEVER

*Compare to the active ingredient in Regular Strength Tylenol® Tablets



100
TABLETS

Drug Facts	<p>Active Ingredient (in each tablet) Acetaminophen 325 mg</p> <p>Purpose Pain relief/Fever reducer</p> <p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> ■ muscular aches ■ headache ■ minor pain of arthritis ■ toothache ■ the common cold ■ premenstrual and menstrual cramps ■ temporarily reduces fever <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if: <ul style="list-style-type: none"> ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product </p> <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> ■ skin redness ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. </p> <p>Do not use</p> <ul style="list-style-type: none"> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if the user has ever had an allergic reaction to this product or any of its ingredients. <p>Ask a doctor before use if the user</p> <ul style="list-style-type: none"> ■ has liver disease ■ is a child with pain of arthritis <p>Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin</p>
-------------------	---

DO NOT USE WITH
OTHER MEDICINES
CONTAINING
ACETAMINOPHEN

No Varnish

Child-Resistant Packaging

Drug Facts (Continued under label)

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Regular Strength

**PEEL HERE
FOR MORE
FACTS**

Aarna USA
Manufactured for:
ARNA USA Inc.
Leland, NC-28451

<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if ■ pain gets worse or lasts more than 10 days (for adults) or 5 days (for children) ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present.</p> <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.</p> <p>Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>	
<p>Directions ■ do not take more than directed (see overdose warning)</p>	
adults & children 12 years and over	<p>■ take 2 tablets every 4 to 6 hours while symptoms last</p> <p>■ do not take more than 10 tablets in 24 hours</p> <p>■ do not use for more than 10 days unless directed by a doctor</p>
children 6-11 years	<p>■ take 1 tablet every 4 to 6 hours while symptoms last</p> <p>■ do not take more than 5 tablets in 24 hours</p> <p>■ do not use for more than 5 days unless directed by a doctor</p>
child under 6 years	ask a doctor
<p>Other Information: store at 20° to 25°C (68° to 77°F)</p>	
<p>Inactive ingredients: Gelatinized starch, magnesium stearate, povidone</p>	
<p>Questions or comments? Call toll-free 1-877-225-6999</p>	

NDC 82568-00012-9

NDC 82568-0001-2

Acetaminophen 325MG TABLETS

REGULAR STRENGTH

ASPIRIN FREE
PAIN RELIEVER &
FEVER RELIEVER

*Compare to the active ingredient
in Regular Strength Tylenol® Tablets

DO NOT USE WITH
OTHER MEDICINES
CONTAINING
ACETAMINOPHEN

1000
TABLETS

Drug Facts

Active ingredient (in each tablet)
Acetaminophen 325 mg Pain reliever/ Fever reducer

Purpose

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - backache
 - minor pain of arthritis
 - the common cold
 - toothache
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

 If a skin reaction occurs, stop use and seek medical help right away.

Do not use

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If the user has ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if the user

- has liver disease
- is a child with pain of arthritis

Ask a doctor or pharmacist before use if the user is

taking the blood thinning drug warfarin

Child-Resistant Packaging

Drug Facts (continued)

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days (for adults) or 5 days (for children)
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present.

 These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- adults & children 12 years and over
 - take 2 tablets every 4 to 6 hours while symptoms last
 - do not take more than 10 tablets in 24 hours
 - do not use for more than 10 days unless directed by a doctor
- children 6-11 years
 - take 1 tablet every 4 to 6 hours while symptoms last
 - do not take more than 5 tablets in 24 hours
 - do not use for more than 5 days unless directed by a doctor
- child under 6 years
 - ask a doctor

Do to at 20° to 25° C (68° to 77° F)

Other information: store at 20° to 25° C (68° to 77° F)

Inactive ingredients: gelatinized starch, magnesium stearate, povidone

Questions or comments? Call toll-free 1-877-225-6999

Do not use if imprinted safety seal under cap is broken or missing
*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Regular Strength

No Varnish

Manufactured for:
AARNA USA Inc.
Lenoir, NC-28645

100
TABLETS
REGULAR STRENGTH

			
---	---	--	---

ACETAMINOPHEN			
acetaminophen tablet			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82568-0001
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	325 mg
Inactive Ingredients			
Ingredient Name			Strength
STARCH, CORN (UNII: O8232NY3SJ)			
POVIDONE (UNII: FZ989GH94E)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	4mm
Flavor		Imprint Code	A325
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82568-0001-1	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/15/2025	
2	NDC:82568-0001-2	1000 in 1 BOTTLE; Type 0: Not a Combination Product	05/15/2025	
3	NDC:82568-0001-3	100 in 1 PACKAGE; Type 0: Not a Combination Product	05/15/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/15/2025	

Labeler - AARNA USA INC. (118515992)

Revised: 4/2025

AARNA USA INC.