

ACID RELIEF- famotidine tablet, film coated
Rite Aid Corporation

Rite Aid Corporation Acid Relief Drug Facts

Active ingredient (in each tablet)

Famotidine 20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20° -25°C (68° -77°F)
- protect from moisture

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, FD&C blue #1 aluminum lake, hypromellose, lactose (monohydrate), magnesium stearate, maltodextrin, microcrystalline cellulose, modified food starch, natural and artificial flavor, sucralose, titanium dioxide, triacetin

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

FREE FROM | GLUTEN FREE

SOY FREE

Compare to the active ingredient of Maximum Strength Pepcid® AC

MAXIMUM STRENGTH

ACID RELIEF

Famotidine Tablets, 20 mg

Acid Reducer

Just one tablet prevents & relieves heartburn due to acid indigestion

COOL MINT FLAVOR

Releases a cooling sensation in mouth & throat

ACTUAL SIZE

50 TABLETS

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.



ACTUAL SIZE

**MAXIMUM STRENGTH
ACID RELIEF**
Famotidine Tablets, 20 mg
Acid Reducer
Just one tablet prevents and relieves heartburn due to acid indigestion. Releases a cooling sensation in mouth & throat.
COOL MINT FLAVOR
50 TABLETS

NDC 11822-1014-0

FREE FROM GLUTEN FREE SOY FREE

Compare to the active ingredient of Maximum Strength Pepcid® AC*

MAXIMUM STRENGTH ACID RELIEF

Famotidine Tablets, 20 mg

Acid Reducer

Just one tablet prevents & relieves heartburn due to acid indigestion. Releases a cooling sensation in mouth & throat.



COOL MINT FLAVOR



ACTUAL SIZE

50 TABLETS



DISTRIBUTED BY:
RHEINOLD, 30 HUNTER LANE
CAMDEN, NJ 07911
www.rheinald.com

OUR PROMISE:
Gluten Free
Soy Free

SATISFACTION GUARANTEE
If you're not satisfied, we'll happily refund your money.

32FT1 89 C1



TIPS FOR MANAGING HEARTBURN


Do not lie flat or bend over after eating.
Do not wear tight-fitting clothing around the stomach.
Do not eat before bedtime.
Raise the head of your bed.

Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables.
Eat slowly and avoid big meals.
If overweight, lose weight.
Quit smoking.

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

Drug Facts	Drug Facts (continued)
<p>Active ingredient (in each tablet) Famotidine 20 mg..... Acid reducer</p> <p>Purpose</p> <p>Uses <ul style="list-style-type: none"> relieves heartburn associated with acid indigestion and sores in the stomach prevents heartburn associated with acid indigestion and sores in the stomach brought on by eating or drinking certain foods and beverages </p> <p>Warnings <p>Allergy alert: Do not use if you are allergic to famotidine or other acid reducers.</p> <p>Do not use <ul style="list-style-type: none"> if you have trouble swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. with other acid reducers </p> <p>Ask a doctor before use if you have <ul style="list-style-type: none"> had heartburn over 3 months. This may be a sign of a more serious condition. </p> </p>	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> heartburn with lightheadedness, sweating, or dizziness chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck, or shoulders; or lightheadedness frequent chest pain frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain kidney disease <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if <ul style="list-style-type: none"> your heartburn continues or worsens you need to take this product for more than 14 days </p> <p>If pregnant or breastfeeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p>

<p>Drug Facts (continued)</p> <p>Directions <ul style="list-style-type: none"> adults and children 12 years and over: <ul style="list-style-type: none"> to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food and drinking beverages that cause heartburn. </p>	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> store at 20°-25°C (68°-77°F) protect from moisture <p>Inactive ingredients: colloid silicon dioxide, croscarmellose sodium, FD&C blue #1 aluminum lake, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin, microcrystalline cellulose, and food-grade starch, natural and artificial flavors.</p>
--	--



ACID RELIEF

famotidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-1014
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	BLUE	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	32F
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-1014-0	1 in 1 CARTON	03/31/2022	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077351	03/31/2022	

Labeler - Rite Aid Corporation (014578892)

Revised: 4/2022

Rite Aid Corporation