

**MAXI-OTIC- gentamicin sulfate, mometasone furoate monohydrate, and clotrimazole suspension**  
**Butler Animal Health Supply**

-----  
**Maxi-Otic™**

**(GENTAMICIN SULFATE, USP; MOMETASONE FUROATE MONOHYDRATE; AND CLOTRIMAZOLE, USP, OTIC SUSPENSION)**

F-27078915

NADA #141-177, Approved by FDA.

**PRODUCT INFORMATION**

**VETERINARY**

**For Otic Use in Dogs Only**

**CAUTION Federal law restricts this drug to use by or on the order of a licensed veterinarian.**

**Keep this and all drugs out of the reach of children.**

**DESCRIPTION** Each gram of Maxi-Otic™ Otic Suspension contains gentamicin sulfate, USP equivalent to 3 mg gentamicin base; mometasone furoate monohydrate equivalent to 1 mg mometasone; and 10 mg clotrimazole, USP in a mineral oilbased system containing a plasticized hydrocarbon gel.

**PHARMACOLOGY**

**Gentamicin:** Gentamicin sulfate is an aminoglycoside antibiotic active against a wide variety of gram-negative and grampositive bacteria. *In vitro* tests have determined that gentamicin is bactericidal and acts by inhibiting normal protein synthesis in susceptible microorganisms. In clinical trials, gentamicin was shown to have a range of activity against the following organisms commonly isolated from infected canine ears:

*Pseudomonas* spp. (including *P. aeruginosa*), coagulasepositive staphylococci, *Enterococcus faecalis*, *Proteus mirabilis* and beta-hemolytic streptococci.

**Mometasone:** Mometasone furoate monohydrate is a synthetic adrenocorticoid characterized by a novel (2') furoate 17-ester having chlorine at the 9 and 21 positions, which have shown to possess high topical potency.

Systemic absorption of mometasone furoate ointment was found to be minimal (2%) over 1 week when applied topically to dogs with intact skin. In a 6-month dermal toxicity study using 0.1% mometasone ointment on healthy intact skin in dogs, systemic effects typical of corticosteroid therapy were noted.

The extent of percutaneous absorption of topical corticosteroids is determined by many factors including the integrity of the epidermal barrier. Topical corticosteroids can be absorbed from normal, intact skin. Inflammation can increase percutaneous absorption. Once absorbed through the skin, topical corticosteroids are handled through pharmacokinetic pathways similar to systemically administered corticosteroids.

**Clotrimazole:** Clotrimazole is a broad-spectrum antifungal agent that is used for the treatment of dermal infections caused by various species of dermatophytes and yeast. The primary action of clotrimazole is against dividing and growing organisms.

*In vitro*, clotrimazole exhibits fungistatic and fungicidal activity against isolates of *Trichophyton rubrum*, *Trichophyton mentagrophytes*, *Epidermophyton floccosum*, *Microsporum canis*, *Candida* spp., and *Malassezia pachydermatis*. Resistance to clotrimazole is very rare among the fungi that cause superficial mycoses. In an induced otitis externa study using dogs infected with *Malassezia pachydermatis*, 1% clotrimazole in the vehicle formulation was effective both microbiologically and clinically in terms of

reduction of exudate, odor, and swelling.

In studies of the mechanism of action, the minimum fungicidal concentration of clotrimazole caused leakage of intracellular phosphorus compounds into the ambient medium with concomitant breakdown of cellular nucleic acids and accelerated potassium efflux. These events began rapidly and extensively after addition of the drug. Clotrimazole is very poorly absorbed following dermal application.

**Gentamicin-Mometasone-Clotrimazole:** By virtue of its three active ingredients, Maxi-Otic Otic Suspension has antibacterial, anti-inflammatory, and antifungal activity. In clinical field trials, Maxi-Otic Otic Suspension was effective in the treatment of otitis externa associated with bacteria and *Malassezia pachydermatis*. Maxi-Otic Otic Suspension reduced discomfort, redness, swelling, exudate, and odor.

**INDICATIONS** Maxi-Otic Otic Suspension is indicated for the treatment of otitis externa in dogs caused by susceptible strains of yeast (*Malassezia pachydermatis*) and bacteria (*Pseudomonas* spp. [including *P. aeruginosa*], coagulase-positive staphylococci, *Enterococcus faecalis*, *Proteus mirabilis*, and beta-hemolytic streptococci).

**CONTRAINDICATIONS** If hypersensitivity to any of the components occurs, treatment should be discontinued and appropriate therapy instituted. Concomitant use of drugs known to induce ototoxicity should be avoided. Do not use in dogs with known perforation of eardrums.

**WARNINGS** The use of these components has been associated with deafness or partial hearing loss in a small number of sensitive dogs (eg, geriatric). The hearing deficit is usually temporary. If hearing or vestibular dysfunction is noted during the course of treatment, discontinue use of Maxi-Otic Otic Suspension immediately and flush the ear canal thoroughly with a nonototoxic solution.

Corticosteroids administered to dogs, rabbits, and rodents during pregnancy have resulted in cleft palate in offspring. Other congenital anomalies including deformed forelegs, phocomelia, and anasarca have been reported in offspring of dogs that received corticosteroids during pregnancy.

Field and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition if used during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

**PRECAUTIONS** Before instilling any medication into the ear, examine the external ear canal thoroughly to be certain the tympanic membrane is not ruptured in order to avoid the possibility of transmitting infection to the middle ear as well as damaging the cochlea or vestibular apparatus from prolonged contact.

Administration of recommended doses of Maxi-Otic Otic Suspension beyond 7 days may result in delayed wound healing. If overgrowth of nonsusceptible bacteria or fungi occurs, treatment should be discontinued and appropriate therapy instituted.

Avoid ingestion. Adverse systemic reactions have been observed following the oral ingestion of some topical corticosteroid preparations. Patients should be closely observed for the usual signs of adrenocorticoid overdosage which include sodium retention, potassium loss, fluid retention, weight gain, polydipsia, and/or polyuria. Prolonged use or overdosage may produce adverse immunosuppressive effects.

Use of corticosteroids, depending on dose, duration, and specific steroid, may result in endogenous steroid production inhibition following drug withdrawal. In patients presently receiving or recently withdrawn from corticosteroid treatments, therapy with a rapidly acting corticosteroid should be considered in especially stressful situations.

**TOXICOLOGY** Field and safety studies with Maxi-Otic Otic Suspension have shown a wide safety margin at the recommended dose level in dogs (see **PRECAUTIONS/ADVERSE REACTIONS**).

## **ADVERSE REACTIONS**

**Gentamicin:** While aminoglycosides are absorbed poorly from skin, intoxication may occur when aminoglycosides are applied topically for prolonged periods of time to large wounds, burns, or any denuded skin, particularly if there is renal insufficiency. All aminoglycosides have the potential to produce reversible and irreversible vestibular, cochlear, and renal toxicity.

**Mometasone:** ALP (SAP) and ALT (SGPT) enzyme elevations, weight loss, anorexia, polydipsia, polyuria, neutrophilia, and lymphopenia have occurred following the use of parenteral, high-dose, and/or prolonged or systemic synthetic corticosteroids in dogs. Cushing's syndrome in dogs has been reported in association with prolonged or repeated steroid therapy.

**Clotrimazole:** The following have been reported occasionally in humans in connection with the use of clotrimazole: erythema, stinging, blistering, peeling, edema, pruritus, urticaria, and general irritation of the skin not present before therapy.

**Maxi-Otic Otic Suspension:** In field studies following once daily treatment with Maxi-Otic Otic Suspension, ataxia, proprioceptive deficits, and increased water consumption were observed in less than 1% of 164 dogs. In a field study following twice-daily treatment with Maxi-Otic Otic Suspension, inflammation of the pinna and diarrhea were observed in less than 1% of 141 dogs.

## **DOSAGE AND ADMINISTRATION**

The external ear canal should be thoroughly cleaned and dried before treatment. Verify that the eardrum is intact. For dogs weighing less than 30 lbs, instill 4 drops from the 7.5 g, 15 g, and 30 g bottles (2 drops from the 215 g bottle) of Maxi-Otic Otic Suspension once daily into the ear canal. For dogs weighing 30 lbs or more, instill 8 drops from the 7.5 g, 15 g, and 30 g bottles (4 drops from the 215 g bottle) once daily into the ear canal. Therapy should continue for 7 consecutive days.

**HOW SUPPLIED** Maxi-Otic Otic Suspension is available in 7.5 g (NDC 11695-7025-7), 15 g (NDC 11695-7025-1), 30 g (NDC 11695-7025-3), and 215 g (NDC 11695-7025-2) plastic bottles.

**Store between 2° and 25°C (36° and 77°F). Shake well before use.**

U.S. Patent No. 6,127,353.

Distributed by  
Henry Schein Animal Health  
Dublin, OH 43017

Made in Canada.  
11/14

81-151901

## **PRINCIPAL DISPLAY PANEL - 215 g Bottle Carton**

**HENRY SCHEIN®**  
ANIMAL HEALTH

NDC #11695-7025-2

**MAXI-OTIC™**  
(Gentamicin Sulfate, USP;  
Mometasone Furoate  
Monohydrate; and  
Clotrimazole, USP, Otic  
Suspension)

**For otic use in dogs only.**

**Keep Out of Reach of Children.**

**CAUTION:** Federal law restricts  
this drug to use by or on the order  
of a licensed veterinarian.

NADA #141-177, Approved by FDA

**Reorder No. 054570**

**Net Contents: 215 g**

Distributed Exclusively by  
Henry Schein Animal Health  
Dublin, OH 43017

**[www.henryscheinvet.com](http://www.henryscheinvet.com)**

**HENRY SCHEIN**

**SEAL OF EXCELLENCE®**



## Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:11695-7025
<b>Route of Administration</b>	AURICULAR (OTIC)		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>Gentamicin Sulfate</b> (UNII: 8X7386QRLV) (Gentamicin - UNII:T6Z9V48IKG)	Gentamicin	3 mg in 1 g
<b>Mometasone Furoate Monohydrate</b> (UNII: MTW0WEG809) (Mometasone - UNII:8HR4QJ6DW8)	Mometasone Furoate	1 mg in 1 g
<b>Clotrimazole</b> (UNII: G07GZ97H65) (Clotrimazole - UNII:G07GZ97H65)	Clotrimazole	10 mg in 1 g

## Packaging

#	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:11695-7025-7	1 in 1 CARTON		
1		7.5 g in 1 BOTTLE, DISPENSING		
2	NDC:11695-7025-1	1 in 1 CARTON		
2		15 g in 1 BOTTLE, DISPENSING		
3	NDC:11695-7025-3	1 in 1 CARTON		
3		30 g in 1 BOTTLE, DISPENSING		
4	NDC:11695-7025-2	1 in 1 CARTON		
4		215 g in 1 BOTTLE, PLASTIC		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
NADA	NADA141177	03/15/2015	

**Labeler** - Butler Animal Health Supply (017880659)