

COLD PLUS HEAD CONGESTION SEVERE- acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated
DISCOUNT DRUG MART

DDM-1175-2020-0810

Drug Facts

| <i>Active ingredients (in each caplet)</i> | <i>Purpose</i> |
|---|-----------------------------|
| Acetaminophen 325 mg | Pain reliever/fever reducer |
| Guaifenesin 200 mg | Expectorant |
| Phenylephrine HCl 5 mg | Nasal decongestant |

Uses

- for the temporary relief of the following cold symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

| | |
|---------------------------------------|---|
| adults and children 12 years and over | <ul style="list-style-type: none">▪ take 2 caplets every 4 hours▪ swallow whole - do not crush, chew, or dissolve▪ do not take more than 10 caplets in 24 hours |
| children under 12 years | <ul style="list-style-type: none">▪ ask a doctor |

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, crospovidone, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

PRINCIPAL DISPLAY PANEL

DISCOUNT DRUG MART FOOD FAIR

NDC 53943-175-02

†Compare to the active ingredients in Tylenol® Cold + Head Congestion Severe
FOR ADULTS

COLD + HEAD CONGESTION SEVERE

Acetaminophen, Phenylephrine HCl, Guaifenesin


Pain Reliever / Fever Reducer, Nasal Decongestant, Expectorant

RELIEF OF:

- HEADACHE + BODY ACHES
- FEVER + SORE THROAT
- NASAL CONGESTION
- MUCUS + CHEST CONGESTION

24 COOL TASTE CAPLETS

Actual Size



Drug Facts (continued)

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, crospovidone, >

Drug Facts (continued)

flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours.

■ 3 or more alcoholic drinks every day while using this product

■ With other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains acetaminophen, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease ■ diabetes
- heart disease ■ high blood pressure
- thyroid disease

■ trouble urinating due to an enlarged prostate gland

Drug Facts (continued)

Directions

- do not take more than directed (see overdose warning)
- take 2 caplets every 4 hours
- swallow whole – do not crush, chew, or dissolve
- do not take more than 10 caplets in 24 hours

adults and children 12 years and over

children under 12 years

- ask a doctor

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Keep out of reach of children.

If pregnant or breast-feeding, ask a health professional before use. These could be signs of a serious condition.

- cough comes back or occurs with rash or headache that lasts
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur

Stop use and ask a doctor if

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Drug Facts

Active ingredients (in each caplet)

- Acetaminophen 325 mg.....Pain reliever/fever reducer
- Guafenesin 200 mg.....Expectorant
- Phenylephrine HCl 5 mg.....Nasal decongestant

Purpose

Relief of the following cold symptoms:

- minor aches and pains ■ headache ■ sore throat
- nasal congestion
- sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Drug Facts (continued)

Uses

- for the temporary relief of the following cold symptoms:

This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Cold + Head Congestion Severe.

DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN

Distributed by: Drug Mart - Food Fair
211 Commerce Drive • Medina, OH 44256

SATISFACTION GUARANTEED

IF ASSURED BY THIS RETURN FROM TOWN AND COUNTRY TO STORE SEND RETURN TO ASSURANCE MAIL ADDRESS AND RETURN PACKAGE TO: ASSURANCE MAIL, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

† Compare to the active ingredients in Tylenol® Cold + Head Congestion Severe

FOR ADULTS

COLD + HEAD CONGESTION SEVERE

Acetaminophen, Phenylephrine HCl, Guaifenesin
Pain Reliever / Fever Reducer, Nasal Decongestant, Expectorant

RELIEF OF:

- HEADACHE + BODY ACHES
- NASAL CONGESTION
- FEVER + SORE THROAT
- MUCUS + CHEST CONGESTION



24 COOL TASTE CAPLETS

NC

NC

NC

COLD PLUS HEAD CONGESTION SEVERE

acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:53943-175 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-----------------------------|----------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg |
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) | GUAIFENESIN | 200 mg |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 5 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| ACESULFAME POTASSIUM (UNII: 23OV73Q5G9) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| CROSCARMELLOSE SODIUM (UNII: M28OL1HH48) | |
| CROSPVIDONE (UNII: 2S7830E561) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| MALTODEXTRIN (UNII: 7CVR7L4A2D) | |
| CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| POLYVINYL ALCOHOL (UNII: 532B59J990) | |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E) | |
| STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|----------|
| Color | white | Score | no score |
| Shape | OVAL | Size | 19mm |
| Flavor | MINT | Imprint Code | AAA;1119 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:53943-175-02 | 2 in 1 CARTON | 05/16/2014 | |
| 1 | | 12 in 1 BLISTER PACK; Type 1: Convenience Kit of Co-Package | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 05/16/2014 | |

Labeler - DISCOUNT DRUG MART (047741335)

Revised: 10/2024

DISCOUNT DRUG MART