

MAPAP EXTRA STRENGTH- acetaminophen capsule

Major Pharmaceuticals

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

0617 - Major

Drug Facts

Active Ingredient (in each banded capsule)

Acetaminophen 500 mg

Purpose

Purpose

Pain reliever/ Fever reducer

Uses

Temporarily reduces fever and relieves minor aches and pains due to:

- headache
- muscular aches
- common cold
- toothache
- backache
- minor pain of arthritis
- premenstrual and menstrual cramps

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drugs containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning:

In the case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed (see overdose warning).

adults and children 12 years and over:

- take 2 capsules every 6 hours while symptoms last
- Do not take more than 6 capsules in 24 hours, unless directed by a doctor
- Do not take for more than 10 days unless directed by a doctor

children under 12 years: ask a doctor

Other information

- store at room temperature

Inactive Ingredients

FD&C Blue #1, FD&C Red #40, Gelatin, Polyvinylpyrrolidone, Pregelatinized Starch, Sodium Starch Glycolate, Stearic Acid and Titanium Dioxide

Questions?

Call: (800) 616-2471

Distributed by: **Major® Pharmaceuticals**

17177 N Laurel Park Drive, Suite 233, Livonia, MI 48152 USA

Re-order No. 700691 Rev.05/19 M-29 4050-05-19

Major

NDC 0904-1987-60

EXTRA STRENGTH

Mapap

Acetaminophen 500 mg Each

CAPSULES SAFETY SEALED FOR YOUR PROTECTION

SEE NEW WARNINGS INFORMATION AND DIRECTIONS

ASPIRIN FREE

Strong Pain & Fever Relief

100 CAPSULES

NDC 0904-1987-60



MAJOR
EXTRA STRENGTH
Māpap
Acetaminophen 500 mg Each

SEE NEW WARNINGS INFORMATION AND DIRECTIONS

ASPIRIN FREE
STRONG PAIN & FEVER RELIEF

100 CAPSULES

TAMPER EVIDENT: DO NOT USE IF CAPSULE BAND OR IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each banded capsule) Purpose
Acetaminophen 500 mg Pain reliever/Fever reducer

Uses
Temporarily reduces fever and relieves minor aches and pains due to: • headache • muscular aches • common cold • toothache • backache • minor pain of arthritis • premenstrual and menstrual cramps

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: • skin redness • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
• If you are allergic to acetaminophen or any of the inactive ingredients in this product

Distributed By: **MAJOR® PHARMACEUTICALS**
17177 N Laurel Park Drive, Suite 233, Livonia, MI 48152 USA
Re-order No. 700691 Rev.05/19 M-29 4050-05-19



0904198760 9

3 Lot # & Exp. Date

Drug Facts (continued)

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if
• pain gets worse or lasts more than 10 days
• fever gets worse or lasts more than 3 days
• new symptoms occur
• redness or swelling is present
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Overdose warning: In the case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
• do not take more than directed (see overdose warning)
adults and children 12 years and over:
• take 2 capsules every 6 hours while symptoms last
• do not take more than 6 capsules in 24 hours, unless directed by a doctor
• do not take for more than 10 days unless directed by a doctor
children under 12 years: ask a doctor

Other information store at room temperature

Inactive ingredients FD&C Blue #1, FD&C Red #40, Gelatin, Polyvinylpyrrolidone, Pregelatinized Starch, Sodium Starch Glycolate, Stearic Acid and Titanium Dioxide

Questions or comments?
Call (800) 616-2471

MAPAP EXTRA STRENGTH

acetaminophen capsule

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-1987
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN (UNII: 2G86QN327L)	
POVIDONE K30 (UNII: U725QWY32X)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white, red	Score	no score
Shape	CAPSULE	Size	22mm
Flavor		Imprint Code	CPC;617
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-1987-60	100 in 1 BOTTLE; Type 0: Not a Combination Product	12/22/1997	
2	NDC:0904-1987-80	1000 in 1 BOTTLE; Type 0: Not a Combination Product	01/29/2001	04/30/2017

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	12/22/1997	

Labeler - Major Pharmaceuticals (191427277)

Revised: 6/2019

Major Pharmaceuticals