

**EQUATE INFANTS PAIN AND FEVER- acetaminophen suspension  
WALMART INC.**

-----  
**Wal-Mart Infants Pain and Fever Drug Facts**

**Active ingredient (in each 5 mL)**

Acetaminophen 160 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

temporarily:

- reduces fever
- relieves minor aches and pains due to:
  - the common cold
  - flu
  - headache
  - sore throat
  - toothache

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if your child has**

liver disease

**Ask a doctor or pharmacist before use if your child is**

taking the blood thinning drug warfarin

**When using this product**

**do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise, use age.
- push air out of syringe. Insert syringe tip into bottle opening.
- flip bottle upside down. Pull yellow part of syringe to the first dose line and then push product back into bottle.
- pull yellow part of syringe until it reaches and stays at the correct dose
- dispense liquid slowly into child’s mouth, toward inner cheek
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- replace cap tightly to maintain child resistance

**Dosing Chart**

<b>Weight (lb)</b>	<b>Age (yr)</b>	<b>Dose (mL)*</b>
under 24	under 2 years	ask a doctor

24-35	2-3 years	5 mL
-------	-----------	------

\* or as directed by a doctor

**Attention:** use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

### **Other information**

- store at 20-25°C (68-77°F)
- **do not use if printed neckband is broken or missing**

### **Inactive ingredients**

anhydrous citric acid, calcium sulfate, carrageenan, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, tribasic sodium phosphate

### **Questions or comments?**

**1-888-287-1915**

### **Package/Label Principal Display Panel**

equate™

Compare to Infants' Tylenol® Oral Suspension active ingredient

infants

Dye-Free

PAIN & FEVER

Acetaminophen

160 mg per 5 mL

Suspension Liquid

Pain Reliever/Fever Reducer

AGES 2 to 3 YEARS

Pediatrician preferred dosing system

- Alcohol-Free
- Aspirin-Free
- Ibuprofen-Free

PARABEN FREE

Grape Flavored

Use only with enclosed syringe. See side panel for more information.

2 FL OZ (59 mL)



**Drug Facts**  
**Active ingredient (in each 5 mL)**    **Purpose**  
 Acetaminophen 160 mg.....Pain reliever/fever reducer

**Uses** temporarily:    ■ reduces fever  
 ■ relieves minor aches and pains due to:    ■ the common cold  
 ■ flu    ■ headache    ■ sore throat    ■ toothache

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes  
 ■ more than 5 doses in 24 hours, which is the maximum daily amount  
 ■ with other drugs containing acetaminophen  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:    ■ skin reddening    ■ blisters    ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.  
**Do not use**    ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if your child has ever had an allergic reaction to this product or any of its ingredients  
**Ask a doctor before use if your child has liver disease**  
**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin**  
**When using this product do not exceed recommended dose (see overdose warning)**  
**Stop use and ask a doctor if**    ■ new symptoms occur  
 ■ pain gets worse or lasts more than 5 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ redness or swelling is present. These could be signs of a serious condition.  
**Keep out of reach of children.**  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**  
 ■ this product does not contain directions or complete warnings for adult use  
 ■ do not give more than directed (see overdose warning)  
 ■ shake well before using    ■ mL = milliliter  
 ■ find right dose on chart. If possible, use weight to dose; otherwise, use age.  
 ■ push air out of syringe. Insert syringe tip into bottle opening.  
 ■ flip bottle upside down. Pull yellow part of syringe to the first dose line and then push product back into bottle. ▶



**Drug Facts (continued)**

- pull yellow part of syringe until it reaches and stays at the correct dose
- dispense liquid slowly into child's mouth, toward inner cheek
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- replace cap tightly to maintain child resistance

**Dosing Chart**

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL

\* or as directed by a doctor  
**Attention:** use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

**Other information**  
 ■ store at 20-25°C (68-77°F)  
 ■ do not use if printed neckband is broken or missing

**Inactive ingredients** anhydrous citric acid, calcium sulfate, carrageenan, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, tribasic sodium phosphate

**Questions or comments?**  
 1-888-287-1915

**DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716**  
 \*\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Infants' Tylenol® Oral Suspension.

NDC 79903-225-02

**equate™**  
**infants**  
 Dye-Free  
**PAIN & FEVER**  
 AGES 2 to 3 YEARS  
 Grape Flavored

Compare to Infants' Tylenol® Oral Suspension active ingredients™

**equate™**  
**infants**  
 Dye-Free  
**PAIN & FEVER**  
 Acetaminophen  
 160 mg per 5 mL  
 Suspension Liquid  
 Pain Reliever/Fever Reducer  
 AGES 2 to 3 YEARS

■ Pediatrician preferred dosing system

- Alcohol-Free
- Aspirin-Free
- Ibuprofen-Free

**PARABEN FREE**

Grape Flavored  
 2 FL OZ (59 mL)

Use only with enclosed syringe. See side panel for more information.

\* 164152  
  
 1 94346 11230 2

76616 2E C1

Compare to Infants' Tylenol® Oral Suspension active ingredients™

**IMPORTANT:**  
 Please see dosing chart or ask a doctor for your child's correct dose.

STEP 1. See *Directions* for the correct dose for your child.  
 STEP 2. Shake well before using.  
 STEP 3. Push air out of syringe. Insert syringe tip into bottle opening.



STEP 4. Flip bottle upside down. Pull yellow part of syringe to the first dose line and then push product back into bottle.  
 STEP 5. Pull yellow part of syringe until it reaches and stays at the correct dose.



STEP 6. Dispense liquid slowly into child's mouth.

Recycled Seal & Cap

  
 PAPER BOX    PLASTIC BOTTLE    PLASTIC WRAPPER    PLASTIC SYRINGE

100% Recycled Content

Satisfaction guaranteed - Or we'll replace it or give you your money back. For questions or comments or to report an undesired reaction or side effect, please call 1-888-287-1915.



## EQUATE INFANTS PAIN AND FEVER

acetaminophen suspension

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:79903-225
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WAT0DDB505)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)	

**Product Characteristics**

Color	WHITE	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-225-02	1 in 1 CARTON	02/20/2024	
1		59 mL in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/20/2024	

**Labeler** - WALMART INC. (051957769)