

**ZILACTIN-B- benzocaine gel**  
**Blairex Laboratories, Inc.**

-----  
**Zilactin B Blairex**

***Drug Facts***

**Active ingredient**

Benzocaine 10%

**Purpose**

Oral pain reliever

**Use**

temporarily relieves pain caused by:

- canker sores
- minor mouth sores and gum irritations
- denture and brace pain

**Warnings**

**Flammable:**

- keep away from fire or flame

**Allergy Alert:**

- do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other “caine” anesthetics

**Stop use and ask a physician if**

- sore mouth symptoms do not improve in 7 days
- swelling, rash or fever develops
- irritation, pain or redness persists or worsens
- apply only to affected area
- do not exceed recommended dosage
- avoid contact with the eyes
- do not use for more than 7 days unless directed by a physician or dentist

**Keep out of reach of children.** If swallowed, get medical help or contact a poison control center immediately.

**Directions**

---

adults and children 2 years and older	Dry affected area. Apply a thin coat of gel with cotton swab or clean finger up to 4 times daily. Allow to dry 30-60 seconds.
children under 12 years	adult supervision should be given in the use of this product
children under 2 years	do not use, consult a physician or dentist

### Other information

- do not peel off protective film. Attempting to peel off film may result in skin irritation or tenderness. To remove film, first apply another coat of Zilactin-B Long Lasting Mouth Sore Gel to film, and immediately wipe the area with a moist gauze pad or tissue.
- contains alcohol 65% by volume
- store at 15-30°C (59-86°F)

### Inactive ingredients

Boric acid, hydroxypropylcellulose, propylene glycol, purified water, salicylic acid, SD alcohol 38-B, tannic acid

Manufactured for: Blairex Laboratories, Inc. P.O. Box 2127 Columbus, IN 47202 USA

 Questions or Comments?

Call TOLL FREE 1-800-252-4739

[www.zilactin.com](http://www.zilactin.com)

PROMOTES HEALING

Oral Pain Reliever

Zilactin<sup>®</sup>-B 6 Hour Canker & Mouth Sore Relief

- Bioadhesive Promotes Healing
- Eat and Drink Pain-Free for up to 6 Hours

Net wt. 0.25 oz (7g)

PROMOTES  
HEALING

Oral Pain  
Reliever

# zilactin-B<sup>®</sup>

## 6 Hour Canker & Mouth Sore Relief

- **Bioadhesive  
Promotes Healing**
- **Eat and Drink Pain-Free  
for up to 6 Hours**

Net wt. 0.25 oz (7g)

Relieves Pain Caused by  
Canker Sores, Dentures or Braces

[www.zilactin.com](http://www.zilactin.com)

# zilactin-B<sup>®</sup>

## Long Lasting Mouth Sore Gel



3 50486 55032 3



Manufactured for:  
**Blair Laboratories, Inc.**  
P.O. Box 2127 Columbus, IN 47202 USA  
Questions or Comments?  
Call TOLL FREE 1-800-252-4739  
[www.zilactin.com](http://www.zilactin.com)

000-500028-04

**Tamper Evident Feature:** Do not use if glued carton has been opened or tampered with.

**Inactive ingredients** boric acid, hydroxypropylcellulose, propylene glycol, purified water, salicylic acid, SD alcohol 38-B, tannic acid

■ store at 15-30°C (59-86°F)  
film, and immediately wipe the area with a moist gauze pad or tissue. ■ contains alcohol 65% by volume  
**Other information** ■ do not peel off protective film. Attempting to peel off film may result in skin irritation or tenderness. To remove film, first apply another coat of Zilactin-B Long Lasting Mouth Sore Gel to clean finger up to 4 times daily. Allow to dry 30-60 seconds.


adults and children 2 years and older	Dry affected area. Apply a thin coat of gel with cotton swab or clean finger up to 4 times daily. Allow to dry 30-60 seconds.
children under 12 years	adult supervision should be given in the use of this product
children under 2 years	do not use, consult a physician or dentist

### Directions

**Keep out of reach of children.** If swallowed, get medical help or contact a poison control center immediately.

■ apply only to affected area ■ do not exceed recommended dosage ■ avoid contact with the eyes ■ do not use for more than 7 days unless directed by a physician or dentist

**Stop use and ask a physician if** ■ sore mouth symptoms do not improve in 7 days ■ swelling, rash or fever develops ■ irritation, pain or redness persists or worsens



## ZILACTIN-B

benzocaine gel

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50486-550
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BENZOCAINE</b> (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	0.1 g in 1 g

### Inactive Ingredients

Ingredient Name	Strength
<b>BORIC ACID</b> (UNII: R57ZHV85D4)	
<b>HYDROXYPROPYL CELLULOSE (TYPE M)</b> (UNII: U3JF91U133)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>PEPPERMINT OIL</b> (UNII: AV092KU4JH)	
<b>TANNIC ACID</b> (UNII: 28F9E0DJY6)	
<b>WATER</b> (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50486-550-32	1 in 1 CARTON	06/29/2005	
1		7 g in 1 TUBE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	06/29/2005	

**Labeler** - Blairex Laboratories, Inc. (092575133)

Revised: 11/2024

Blairex Laboratories, Inc.