# COLD FLU RELIEF DAYTIME- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled AmerisourceBergen (Good Neighbor Pharmacy) 46122

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

## **Drug Facts**

# Active ingredients (in each softgel)

# Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

# **Purpose**

#### Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

#### Uses

- temporarily relieves common cold and flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - o nasal congestion
  - fever
  - cough due to minor throat and bronchial irritation

# Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert**: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure

- whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

# Ask a doctor before use if you have

- liver disease
- diabetes
- heart disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

# Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

### When using this product,

do not exceed recommended dos age.

#### Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

# If pregnant or breast-feeding,

ask a health professional before use.

## Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

## Other information

- store between 15-30°C (59-86°F)
- avoid excessive heat

# **Inactive ingredients**

butylated hydroxyanisole, butylated hydroxytoluene, carminic acid\*, edible white ink, D&C yellow #10\*, FD&C red #40\*, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sodium metabisulfite\*, sorbitan, sorbitol

# Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

# **Principal Display Panel**

\*\*Compare to Vicks® DayQuil® LiquiCaps® active ingredients

Non-Drowsy

**DAYTIME Liquid Caps** 

ACETAMINOPHEN, DEXTROMETHORPHAN HBr, PHENYLEPHRINE HCl

Cold & Flu

Pain Reliever-Fever Reducer

Cough Suppressant-Nasal Decongestant

Relieves Major Cold and Flu Symptoms

Through Your Day

Alcohol free, Antihistamine free

**SOFTGELS** 

\*\*This product is not manufactured or distributed by The Procter & Gamble Company, Vicks®, DayQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

## KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Distributed By AmerisourceBergen

1300 Morris Drive, Chesterbrook, PA 19087

Questions or Concerns? Visit us at www.mygnp.com

# **Product Label**

<sup>\*</sup>contains one or more of these ingredients

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST Unestions of comments?

contains one or more of these ingredients propylene glycol, purified water, sodium metabisulfite\*, sorbitan\*, sorbitol #40", FD&C yellow #6, gelatin, glycerin, polyethylene glycol", povidone, hydroxytoluene\*, carminic acid\*, D&C yellow #10\*, edible white ink, FD&C red INSCINCE INGLEGIEUIS DUTYISTED INJURION PRINTED PAY INDUSTED

■ store between 15-30°C (59-86°F)
■ avoid excessive heat

Uther Intormation

label to ensure correct dosing

- when using other Daytime or Mighttime products, carefully read each
  - children under 12 years: do not use
  - swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 4 hours
  - do not take more than 4 doses in 24 hours

DILECTIONS 

a do not take more than directed (see Overdose warning)

even if you do not notice any signs or symptoms. right away. Quick medical attention is critical for adults as well as for children damage. In case of overdose, get medical help or contact a Poison Control Center Overdose warning: Taking more than the recommended dose can cause liver keep out of reach of children.

If pregnant or breast-feeding, ask a health professional before use.

- signs of a serious condition. ■ condu comes pack or occurs with rash or headache that lasts. These could be
  - redness or swelling is present
     new symptoms occur
    - fever gets worse or lasts more than 3 days
    - UGLAORZUGSZ OLZSIUGZZ OLZGEBIGSZUGSZ OCCOL
  - pain, cough, or nasal congestion gets worse or lasts more than 7 days Stop use and ask a doctor if

nund racts (continued)

When using this product, do not exceed recommended dosage.

thinning drug warrarin. yak a doctor or pharmacist betore use it you are taking the blood

- condp that occurs with too much phiegm (mucus)
- ешылугеша ■ becalagent or chronic cough such as occurs with smoking, asthma, or
- trouble unnating due to an enlarged prostate gland
- high blood pressure ■ heart disease
  ■ thyroid disease Ask a doctor before use if you have ■ liver disease ■ diabetes

pharmacist before taking this product.

- not know if your prescription drug contains an MAOI, ask a doctor or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do (certain drugs for depression, psychiatric, or emotional conditions, or
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) sak a doctor or pharmacist.
- nonprescription). If you are not sure whether a drug contains acetaminophen, ■ with any other drug containing acetaminophen (prescription or

cousnit a doctor promptly. accompanied or followed by fever, headache, rash, nausea, or vomiting, Sore throat warning: If sore throat is severe, persists for more than 2 days, is It a skin reaction occurs, stop use and seek medical help right away. may include: ■ skin reddening ■ blisters ■ rash

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms

- 3 or more alcoholic drinks every day while using this product
  - with other drugs containing acetaminophen
  - more than 4,000 mg of acetaminophen in 24 hours

occnr if you take:

Liver warning: This product contains acetaminophen. Severe liver damage may Warnings

nund racts (continued)

■ cough due to minor throat and bronchial irritation ■ usasi congestion

19v9f ■ ■ µезqэсµе ■ winor aches and pains

■ temporarily relieves common cold and flu symptoms

(couruned)

Nasal decongestant Phenylephrine HCl 5 mg. Dextromethorphan HBr 10 mg. Cough suppressant Acetaminophen 325 mg. Pain reliever/fever reducer PUTPOSE

Active ingredients (in each softgel)

Drug Facts

\*Compare to Vicks\* DayQuil\* LiquiCaps\* active ingredients



NDC 46122-339-73

NON-DROWSY

# Day Time Liquid Caps

acetaminophen, dextromethorphan HBr, phenylephrine HCl

Cold & Flu

Pain Reliever-Fever Reducer Cough Suppressant-Nasal Decongestant Relieves Major Cold and Flu Symptoms **Through Your Day** 

Alcohol Free Antihistamine Free



PER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING. KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

"This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.



PLD-E40P FC002934

Distributed By AmerisourceBergen 1300 Morris Drive, Chesterbrook, PA 19087 Questions or Concerns? Visit us at www.mygnp.com





## Good Neighbor Pharmacy DayTime Cold & Flu

# **COLD FLU RELIEF DAYTIME**

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

<b>Product Information</b>
----------------------------

Product Type HUMAN OTC DRUG Item Code (Source) NDC:46 122-339

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	325 mg	
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9 D2RTI9 KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
<b>PHENYLEPHRINE HYDRO CHLO RIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients		
Ingredient Name	Strength	
BUTYLATED HYDRO XYANISOLE (UNII: REK4960K2U)		
BUTYLATED HYDRO XYTO LUENE (UNII: 1P9 D0 Z171K)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		
GELATIN (UNII: 2G86QN327L)		
GLYCERIN (UNII: PDC6A3C0OX)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
PO VIDO NE (UNII: FZ989 GH9 4E)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SORBITAN (UNII: 6O92ICV9RU)		
SORBITOL (UNII: 506T60A25R)		
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)		
CARMINIC ACID (UNII: CID8 Z8 N9 5N)		
SODIUM METABISULFITE (UNII: 4VON5FNS3C)		

Product Characteristics			
Color	ORANGE	Score	no score

Shape	CAPSULE	Size	19 mm
Flavor		Imprint Code	P19;95A;512;P119
Contains			

l	Packaging				
	# Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
	1 NDC:46122-339-73	16 in 1 CARTON	11/30/2015	12/31/2021	
	1	1 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	11/30/2015	12/31/2021

Labeler - AmerisourceBergen (Good Neighbor Pharmacy) 46122 (007914906)

Revised: 11/2019 AmerisourceBergen (Good Neighbor Pharmacy) 46122