

**MEDICHOICE OF HYDROGEN PEROXIDE 3 10 VOLUME- hydrogen peroxide liquid**  
**Owens & Minor Distribution, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**MediChoice Solution of Hydrogen Peroxide 3% USP 10 Volume**

***Drug Facts***

***Active ingredient***

Hydrogen Peroxide (stabilized) 3%

***Purpose***

First aid antiseptic/Oral debriding agent

***Uses***

- first aid to help prevent the risk of infection in minor cuts, scrapes and burns
- aids in the removal of phlegm, mucous, or other secretions associated with occasional sore mouth

***Warnings***

**For external use only**

**Do not use**

- in the eyes or apply over large areas of the body • longer than 1 week

**Ask a doctor before use if you have**

deep or puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if**

- the condition persists or gets worse
- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

***Directions***

First aid antiseptic:

- clean the affected area
- apply a small amount of product on the area 1 to 3 times daily
- may be covered with a sterile bandage
- if bandaged, let dry first Oral debriding agent (oral rinse): adults and children 2 years of age and over:
- mix with an equal amount of water
- swish around in the mouth over the affected area for at least 1 minute and then spit out
- use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor

- children under 12 years of age should be supervised in the use of this product
- children under 2 years of age: consult a dentist or doctor


**Other information**

Keep tightly closed in a cool dark place. Do not shake bottle. Hold away from face when opening.

**Inactive ingredient**

purified water


**MediChoice Solution of Hydrogen Peroxide 3% USP 10 Volume 4 Fl.Oz./118mL (39892-0700-1)**



NDC 39892-0700-1

**SOLUTION OF  
HYDROGEN PEROXIDE  
3% USP 10 VOLUME**

**3%**

**FIRST AID ANTISEPTIC**  
**Made in USA** 

SEALED FOR YOUR PROTECTION  
IF SEAL IS BROKEN OR MISSING  
DO NOT USE

4 FL. OZ. (118 ml)

Distributed by Owens & Minor  
9120 Lockwood Boulevard  
Mechanicsville, VA 23116

Drug Facts

<b>Active ingredient</b>	<b>Purpose</b>
Hydrogen Peroxide (stabilized) 3%.....	First aid antiseptic/Oral debriding agent

Uses

- first aid to help prevent the risk of infection in minor cuts, scrapes and burns
- aids in the removal of phlegm, mucous, or other secretions associated with occasional sore mouth

Warnings For external use only

**Do not use** • in the eyes or apply over large areas of the body • longer than 1 week

**Ask a doctor before use if you have** deep or puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** • the condition persists or gets worse

- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness persists or worsens
- swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

<p><b>First aid antiseptic:</b></p> <ul style="list-style-type: none"> <li>• clean the affected area</li> <li>• apply a small amount of product on the area 1 to 3 times daily</li> <li>• may be covered with a sterile bandage</li> <li>• if bandaged, let dry first</li> </ul>	<p><b>Oral debriding agent (oral rinse):</b> adults and children 2 years of age and over:</p> <ul style="list-style-type: none"> <li>• mix with an equal amount of water</li> <li>• swish around in the mouth over the affected area for at least 1 minute and then spit out</li> <li>• use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor</li> </ul>
--	---

- children under 12 years of age should be supervised in the use of this product
- children under 2 years of age: consult a dentist or doctor

Other information

Keep tightly closed in a cool dark place. Do not shake bottle. Hold away from face when opening.

Inactive ingredient

purified water

1-07-F0010029C

Reorder: HP0304

**MediChoice Solution of Hydrogen Peroxide 3% USP 10 Volume 8 Fl.Oz./237mL (39892-0700-2)**

# MEDICHOICE®

NDC 39892-0700-2

**SOLUTION OF  
HYDROGEN PEROXIDE  
3% USP 10 VOLUME**

# 3%

FIRST AID ANTISEPTIC

Made in USA 

SEALED FOR YOUR PROTECTION  
IF SEAL IS BROKEN OR MISSING  
DO NOT USE



**8 FL. OZ. (237 ml)**

Distributed by Owens & Minor  
9120 Lockwood Boulevard  
Mechanicsville, VA 23116

1-07-D0011029C

**Reorder: HP0308**

## Drug Facts

Active ingredient	Purpose
Hydrogen Peroxide (stabilized) 3%.....	First aid antiseptic/Oral debriding agent

**Uses** • first aid to help prevent the risk of infection in minor cuts, scrapes and burns  
• aids in the removal of phlegm, mucous, or other secretions associated with occasional sore mouth

### Warnings For external use only

**Do not use** • in the eyes or apply over large areas of the body • longer than 1 week

**Ask a doctor before use if you have** deep or puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** • the condition persists or gets worse  
• sore mouth symptoms do not improve in 7 days  
• irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

### Directions

#### First aid antiseptic:

- clean the affected area
- apply a small amount of product on the area 1 to 3 times daily
- may be covered with a sterile bandage
- if bandaged, let dry first

#### Oral debriding agent (oral rinse):

- adults and children 2 years of age and over:
  - mix with an equal amount of water
  - swish around in the mouth over the affected area for at least 1 minute and then spit out
  - use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor

- children under 12 years of age should be supervised in the use of this product
- children under 2 years of age: consult a dentist or doctor

**Other information** Keep tightly closed in a cool dark place. Do not shake bottle. Hold away from face when opening.

**Inactive ingredient** purified water

# MEDICHOICE®

NDC 39892-0700-2

**SOLUTION OF  
HYDROGEN PEROXIDE  
3% USP 10 VOLUME**

# 3%

FIRST AID ANTISEPTIC

Made in USA 

SEALED FOR YOUR PROTECTION  
IF SEAL IS BROKEN OR MISSING  
DO NOT USE



**8 FL. OZ. (237 ml)**

Distributed by Owens & Minor  
9120 Lockwood Boulevard  
Mechanicsville, VA 23116

1-07-D0011029C

**Reorder: HP0308**

## Drug Facts

Active ingredient	Purpose
Hydrogen Peroxide (stabilized) 3%.....	First aid antiseptic/Oral debriding agent

**Uses** • first aid to help prevent the risk of infection in minor cuts, scrapes and burns  
• aids in the removal of phlegm, mucous, or other secretions associated with occasional sore mouth

### Warnings For external use only

**Do not use** • in the eyes or apply over large areas of the body • longer than 1 week

**Ask a doctor before use if you have** deep or puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** • the condition persists or gets worse  
• sore mouth symptoms do not improve in 7 days  
• irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

### Directions

#### First aid antiseptic:

- clean the affected area
- apply a small amount of product on the area 1 to 3 times daily
- may be covered with a sterile bandage
- if bandaged, let dry first

#### Oral debriding agent (oral rinse):

- adults and children 2 years of age and over:
  - mix with an equal amount of water
  - swish around in the mouth over the affected area for at least 1 minute and then spit out
  - use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor

- children under 12 years of age should be supervised in the use of this product
- children under 2 years of age: consult a dentist or doctor

**Other information** Keep tightly closed in a cool dark place. Do not shake bottle. Hold away from face when opening.

**Inactive ingredient** purified water

MediChoice Solution of Hydrogen Peroxide 3% USP 10 Volume 16 Fl.Oz./473mL (39892-0700-3)

# MEDICHOICE®

NSN 6505-00-153-8480  
NDC 39892-0700-3

**SOLUTION OF  
HYDROGEN PEROXIDE  
3% USP 10 VOLUME**

# 3%

**FIRST AID ANTISEPTIC**

**Made in USA** 

TAMPER EVIDENT CAP FOR YOUR PROTECTION. IF RING-BAND IS DETACHED FROM CAP OR MISSING DO NOT USE.



**16 FL. OZ. (473 ml)**

Distributed by Owens & Minor  
9120 Lockwood Boulevard  
Mechanicsville, VA 23116

1-07-D0012029C

**Reorder: HP0316**

## Drug Facts

Active ingredient	Purpose
Hydrogen Peroxide (stabilized) 3%.....	First aid antiseptic/Oral debriding agent

**Uses** • first aid to help prevent the risk of infection in minor cuts, scrapes and burns  
• aids in the removal of phlegm, mucous, or other secretions associated with occasional sore mouth

### Warnings For external use only

**Do not use** • in the eyes or apply over large areas of the body • longer than 1 week

**Ask a doctor before use if you have** deep or puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** • the condition persists or gets worse  
• sore mouth symptoms do not improve in 7 days  
• irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

### Directions

**First aid antiseptic:**

- clean the affected area
- apply a small amount of product on the area 1 to 3 times daily
- may be covered with a sterile bandage
- if bandaged, let dry first

**Oral debriding agent (oral rinse):**

- adults and children 2 years of age and over:
  - mix with an equal amount of water
  - swish around in the mouth over the affected area for at least 1 minute and then spit out
  - use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor

- children under 12 years of age should be supervised in the use of this product
- children under 2 years of age: consult a dentist or doctor

**Other information** Keep tightly closed in a cool dark place. Do not shake bottle. Hold away from face when opening.

**Inactive ingredient** purified water

# MEDICHOICE®

NSN 6505-00-153-8480  
NDC 39892-0700-3

## SOLUTION OF HYDROGEN PEROXIDE 3% USP 10 VOLUME

# 3%

FIRST AID ANTISEPTIC

Made in USA 

TAMPER EVIDENT CAP FOR YOUR PROTECTION. IF RING-BAND IS DETACHED FROM CAP OR MISSING DO NOT USE.



16 FL. OZ. (473 ml)

Distributed by Owens & Minor  
9120 Lockwood Boulevard  
Mechanicsville, VA 23116

1-07-D0012029C

Reorder: HP0316

### Drug Facts

Active ingredient	Purpose
Hydrogen Peroxide (stabilized) 3%.....	First aid antiseptic/Oral debriding agent

**Uses** • first aid to help prevent the risk of infection in minor cuts, scrapes and burns  
• aids in the removal of phlegm, mucous, or other secretions associated with occasional sore mouth

### Warnings For external use only

**Do not use** • in the eyes or apply over large areas of the body • longer than 1 week

**Ask a doctor before use if you have** deep or puncture wounds, animal bites or serious burns

**Stop use and ask a doctor if** • the condition persists or gets worse  
• sore mouth symptoms do not improve in 7 days  
• irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

### Directions

**First aid antiseptic:**

- clean the affected area
- apply a small amount of product on the area 1 to 3 times daily
- may be covered with a sterile bandage
- if bandaged, let dry first

**Oral debriding agent (oral rinse):**

- adults and children 2 years of age and over:
  - mix with an equal amount of water
  - swish around in the mouth over the affected area for at least 1 minute and then spit out
  - use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor

- children under 12 years of age should be supervised in the use of this product
- children under 2 years of age: consult a dentist or doctor

**Other information** Keep tightly closed in a cool dark place. Do not shake bottle. Hold away from face when opening.

**Inactive ingredient** purified water

## MEDICHOICE OF HYDROGEN PEROXIDE 3 10 VOLUME

hydrogen peroxide liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:39892-0700
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>HYDROGEN PEROXIDE</b> (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V)	HYDROGEN PEROXIDE	30 mg in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:39892-0700-1	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/20/2019	
2	NDC:39892-0700-2	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/20/2019	
3	NDC:39892-0700-3	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/20/2019	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	03/20/2019	

**Labeler** - Owens & Minor Distribution, Inc. (847412269)

Revised: 12/2019

Owens &amp; Minor Distribution, Inc.