

**MUCINEX FAST-MAX KICKSTART SEVERE CONGESTION AND COUGH AND
MUCINEX FAST-MAX NIGHTTIME COLD AND FLU- dextromethorphan
hydrobromide, guaifenesin, acetaminophen,triprolidine hydrochloride
RB Health (US) LLC**

Active ingredients

(in each 20 mL)

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Triprolidine HCl 2.5 mg

Dextromethorphan HBr » Cough Suppressant

Guaifenesin » Expectorant

Acetaminophen » Pain reliever/fever reducer

Dextromethorphan » Cough suppressant

Triprolidine HCl » Antihistamine

Uses

■ helps loosen phlegm (mucus) and thin
bronchial secretions to rid the bronchial
passageways of bothersome mucus and
make coughs more productive

■ temporarily relieves:

■ cough due to minor throat and
bronchial irritation as may occur with
the common cold or inhaled irritants

■ the intensity of coughing

■ the impulse to cough to help you get
to sleep

■ temporarily relieves these common cold and flu symptoms:

■ cough ■ minor aches and pains ■ sore throat

■ headache ■ runny nose ■ sneezing ■ itching of the nose or throat

■ itchy, watery eyes due to hay fever

■ temporarily reduces fever

■ controls cough to help you get to sleep

Do not use

■ if you are now taking a prescription
monoamine oxidase inhibitor (MAOI) (certain
drugs for depression, psychiatric, or
emotional conditions, or Parkinson's disease),
or for 2 weeks after stopping the MAOI drug.
If you do not know if your prescription drug
contains an MAOI, ask a doctor or
pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product do not use more than directed

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts.
- These could be signs of a serious condition.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease ■ glaucoma

- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product ■ do not use more than directed

- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 9 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Other information

- each 20 mL contains: sodium 16 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Inactive ingredients

ammonium, glycyrrhizate, anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C blue no. 1, flavor, glycerin (soy), propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum
*may contain this ingredient

Inactive ingredients ammonium glycyrrhizate, anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no.

40, flavor, glycerin (soy), propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions?

1-866-MUCINEX (1-866-682-4639)

Mucinex
POWERFUL MULTI-SYMPTOM RELIEF

POWERFUL COUGH & COLD RELIEF » DAY & NIGHT

DAY & NIGHT PACK

POWERFUL RELIEF FOR YOUR DAY
+
NIGHTTIME RELIEF FOR BETTER MORNINGS


TAKE ONLY AS DIRECTED.
PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org
Do not take MUCINEX® FAST-MAX® KICKSTART SEVERE CONGESTION & COUGH, and MUCINEX® FAST-MAX® NIGHTTIME COLD & FLU at the same time.
Always wait at least 4 hours before taking another dose of Mucinex® Liquid.
Maximum Strength per 4-hour dose


63824 99934

COME BACK WITH A KICK™
✓ Relieves Chest Congestion
✓ Controls Cough
✓ Thins & Loosens Mucus
MENTHOL FLAVORED LIQUID

⌚ Nighttime Relief For Better Mornings
✓ Fever, Headache & Body Pain
✓ Cough + Sore Throat
✓ Runny Nose & Sneezing

AGES 12+
TWO - 6 FL OZ (180 mL) bottles TOTAL - 12 FL OZ (360 mL)
AGES 12+

NIGHTTIME COLD & FLU (cont'd)
Drug Facts (continued)
■ redness or swelling is present
■ new symptoms occur
■ cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
■ do not take more than directed (see Overdose warning)
■ do not take more than 4 doses in any 24-hour period
■ measure only with dosing cup provided
■ do not use dosing cup with other products
■ dose as follows or as directed by a doctor
■ adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
■ children under 12 years of age: do not use
Other information
■ each 20 mL contains: sodium 16 mg
■ store at 20-25°C (68-77°F)
■ do not refrigerate
Inactive ingredients ammonium glycyrrhizate, anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin (soy), propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum
Questions? 1-866-MUCINEX (1-866-682-4639)
Please visit our website
www.mucinex.com
Patents: www.reckitt.com/patents

reckitt
Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
© 2025 RB Health
3318836 121224

NIGHTTIME COLD & FLU

KICKSTART SEVERE CONGESTION & COUGH

Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Triprolidine HCl 2.5 mg	Antihistamine

Purposes

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - sore throat
 - headache
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product do not use more than directed

- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

Drug Facts

Active ingredients (in each 20 mL)

Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant

Purposes

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product do not use more than directed

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 9 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients

ammonium glycyrrhizate, anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C blue no. 1, flavor, glycerin (soy), propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum
*may contain this ingredient

Questions? 1-866-MUCINEX (1-866-682-4639)

MUCINEX FAST-MAX KICKSTART SEVERE CONGESTION AND COUGH AND MUCINEX FAST-MAX NIGHTTIME COLD AND FLU

dextromethorphan hydrobromide, guaifenesin, acetaminophen, triprolidine hydrochloride kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72854-160
--------------	----------------	--------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-160-26	1 in 1 CARTON; Type 1: Convenience Kit of Co-Package	04/02/2025	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	0 BOTTLE	1 mL
Part 2	0 BOTTLE	1

Part 1 of 2

MUCINEX FAST-MAX NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, triprolidine hydrochloride solution

Product Information

Item Code (Source)	NDC:72854-143
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	2.5 mg in 20 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
GLYCERIN (UNII: PDC6A3C0OX)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color	blue	Score	
Shape		Size	
Flavor	FRUIT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-143-66	180 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/02/2025	

Part 2 of 2

MUCINEX FAST-MAX KICKSTART SEVERE CONGESTION AND COUGH MAXIMUM STRENGTH

dextromethorphan hydrobromide, guaifenesin liquid

Product Information

Item Code (Source)	NDC:72854-159
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	

EDETATE DISODIUM (UNII: 7FLD91C86K)	
WATER (UNII: 059QF0KO0R)	
AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	green	Score	
Shape		Size	
Flavor	MENTHOL	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-159-66	180 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/02/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/02/2025	

Labeler - RB Health (US) LLC (081049410)