

**CORICIDIN HBP MAXIMUM STRENGTH COLD, COUGH AND FLU-
acetaminophen, dextromethorphan hydrobromide, doxylamine succinate
Bayer HealthCare LLC.**

**Coricidin HBP Maximum Strength Cold , Cough & Flu Day/Cold & Flu Night
Bundle UIs 1615257/1614785**

Product Name

Coricidin HBP Max Strength Cold, Cough, and Flu

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan hydrobromide 20 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Uses

- temporarily relieves these symptoms due to a cold or flu:
 - minor aches and pains
 - headache
 - cough
 - sore throat
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions.

Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that is accompanied with excessive phlegm (mucus)
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than recommended dose
- adults and children 12 years and over: take 30 mL in the dosing cup provided every 4 hours. Do not exceed 150 mL (5 doses) in 24 hours or as directed by a doctor.
- children under 12 years of age: do not use

Other information

- **each 30 mL contains:** sodium 38 mg
- store between 20° to 25°C (68° to 77°F)

Inactive ingredients anhydrous citric acid, FD&C blue #1, FD&C red # 40, flavor, glycerin, maltitol solution, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments? 1-800-317-2165 (Mon – Fri 9AM – 5PM EST)

Product Name

Coricidin HBP Max Strength Night Cold and Flu

Active ingredients (in each 30mL)

Acetaminophen 650mg
Dextromethorphan hydrobromide 20mg
Doxylamine succinate 12.5mg

Purposes

Pain reliever/fever reducer
Cough suppressant
Antihistamine

- temporarily relieves these symptoms due to a cold or flu:
- minor aches and pains
- headache
- cough
- sore throat
- runny nose
- sneezing

- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions.

Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease),

or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- glaucoma
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with excessive phlegm (mucus)
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than recommended dose
- adults and children 12 years and over: take 30 mL in the dosing cup provided every 4 hours. Do not exceed 150 mL (5 doses) in 24 hours or as directed by a doctor.
- children under 12 years of age: do not use

Other information

- **each 30 mL contains:** sodium 38 mg
- store between 20° to 25°C (68° to 77°F)

Inactive ingredients anhydrous citric acid, FD&C blue #1, FD&C red#40, flavor, glycerin, maltitol solution, polyethylene glycol, propylene glycol, purified water, saccharin

CORICIDIN HBP MAXIMUM STRENGTH COLD, COUGH AND FLU

acetaminophen, dextromethorphan hydrobromide liquid

Product Information

Item Code (Source) NDC:11523-0093

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
MALTITOL (UNII: D65DG142WK)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
GLYCERIN (UNII: PDC6A3C0OX)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
WATER (UNII: 059QF0KO0R)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11523-0093-1	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/27/2022	

Part 2 of 2

CORICIDIN HBP MAXIMUM STRENGTH NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid

Product Information

Item Code (Source) NDC:11523-0077

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
MALTITOL (UNII: D65DG142WK)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
GLYCERIN (UNII: PDC6A3C0OX)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
WATER (UNII: 059QF0KO0R)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11523-0077-1	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/27/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC Monograph Drug M012

05/12/2025

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 12/2025

Bayer HealthCare LLC.