

**MUCINEX NIGHTSHIFT SINUS MAXIMUM STRENGTH- acetaminophen,
dextromethorphan hydrobromide, and triprolidine hydrochloride tablet,
coated**
RB Health (US) LLC

Mucinex Rapid+Clear Nighttime Headache, Cough & Fever

Active ingredients

(in each caplet)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Triprolidine HCl 1.25 mg

Uses

- temporarily relieves these common cold and flu symptoms:
- cough ■ minor aches and pains
- sore throat ■ headache
- runny nose ■ sneezing
- itching of the nose or throat
- itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 8 caplets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase

inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease ■ glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 8 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)

Questions? 1-866-MUCINEX
(1-866-682-4639)

Inactive ingredients

croscarmellose sodium, crospovidone, hypromellose,
microcrystalline cellulose, polyethylene glycol,
polysorbate 80, povidone, titanium dioxide

Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....Cough suppressant
Triprolidine HCl 1.25 mg.....Antihistamine

Keep out of reach of children.

The image displays the packaging for Mucinex Rapid+Clear Night+Time Maximum Strength capsules. The box is primarily blue and red. The top section features the brand name 'Mucinex' in large white letters, followed by 'RAPID+CLEAR' in red and 'NIGHT+TIME' in yellow. Below this, it states 'HEADACHE, COUGH & FEVER'. The active ingredients are listed: Acetaminophen (Pain Reliever/Fever Reducer), Dextromethorphan HBr (Cough Suppressant), and Triprolidine HCl (Antihistamine). The box also includes a 'Drug Facts' panel with detailed information on uses, warnings, and directions. A QR code is visible on the side of the box, and a barcode is at the bottom right. The text '20 CAPLETS' and 'RAPID RELEASE' are prominently displayed.

Drug Facts
Active ingredients (in each caplet)
Acetaminophen 325 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....Cough suppressant
Triprolidine HCl 1.25 mg.....Antihistamine

Purposes
■ temporarily relieves these common cold and flu symptoms:
■ cough ■ sore throat ■ headache ■ sneezing
■ itchy, watery eyes due to hay fever
■ temporarily reduces fever
■ controls cough to help you get to sleep

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 8 caplets in 24 hours, which is the maximum daily amount for this product
■ with other drugs containing acetaminophen
■ 5 or more alcoholic drinks daily while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin redness ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have
■ liver disease ■ glaucoma
■ trouble urinating due to an enlarged prostate gland
■ a breathing problem such as emphysema or chronic bronchitis
■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Drug Facts (continued)
■ cough that occurs with too much phlegm (mucus)
Ask a doctor or pharmacist before use if you are
■ taking the blood thinning drug warfarin
■ taking sedatives or tranquilizers
When using this product
■ do not use more than directed
■ excitability may occur, especially in children
■ marked drowsiness may occur
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ avoid alcoholic drinks
■ use caution when driving a motor vehicle or operating machinery
Stop use and ask a doctor if
■ pain or cough gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ do not take more than directed (see Overdose warning)
■ do not take more than 8 caplets in any 24-hour period
■ adults and children 12 years of age and over: take 2 caplets every 4 hours
■ children under 12 years of age: do not use

Other information
■ store at 20-25°C (68-77°F)

Inactive ingredients
croscarmellose sodium, crospovidone, hypromellose, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639)

Parental Information: Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

PARENTS: Learn about teen medicine abuse. www.StopMedicineAbuse.org

how2recycle.info
PAPER RECYCLED 100%
BOX RECYCLED 100%

20 CAPLETS
RAPID RELEASE
AGES 12+

MUCINEX NIGHTSHIFT SINUS MAXIMUM STRENGTH

acetaminophen, dextromethorphan hydrobromide, and triprolidine hydrochloride tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72854-209
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH)	DEXTROMETHORPHAN	10 mg

(DEXTROMETHORPHAN - UNII:7355X3ROTS)	HYDROBROMIDE	10 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	1.25 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	

Product Characteristics

Color	yellow	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	VW;LOGOcrescentmoonplus
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-209-20	2 in 1 CARTON	05/01/2025	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2025	

Labeler - RB Health (US) LLC (081049410)