

**ARAMARK PAIN FREE- acetaminophen, aspirin, caffeine, salicylamide tablet
Western First Aid Safety DBA Aramark**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Aramark Pain-Free Extra Strength Pain Reliever

Drug Facts

Active Ingredient (in each tablet)

Pain Free Formula L242

Acetaminophen 110 mg

Aspirin (NSAID) 162 mg

Caffeine 32.4 mg

Salicylamide (NSAID) 152 mg

Purpose

AcetaminophenPain Reliever/Fever Reducer
Aspirin (NSAID)Pain Reliever/Fever Reducer
CaffeineAdjuvent
Salicylamide (NSAID)Pain Reliever/Fever Reducer

Uses:

Temporarily Relieves minor aches and pains associated with:

- backach
- headaches
- colds
- toothache
- minor arthritis pain
- muscular aches
- menstrual cramps

Warnings:

- Reye's Syndrome: Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Reye's Syndrome, a rare but serious illness reported to be associated with aspirin.
- **Allergy Alert - Do not use:**
 - if you are allergic to aspirin or any other salicylate
 - if you have ever had an allergic reaction to any other pain reliever/fever reducer
- **Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
 - more than 8 tablets in 24 hours
 - with other drugs containing acetaminophen (prescription or nonprescription)
 - Ask a doctor or pharmacist before using with other drugs if you are not sure.
 - 3 or more alcoholic drinks every day while using this product.

- **Stomach Bleeding Warning:** This product contains nonsteroidal anti-inflammatory drugs (NSAIDs), which may cause stomach bleeding. The chance is higher if you:
 - are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
 - take more or for a longer time than directed
- **Do not use:**
 - with any other pain reliever/fever reducer
 - for pain for more than 10 days or for fever for more than 3 days unless directed by a doctor
 - with any other product containing acetaminophen

Ask a doctor before using if you have:

- upset stomach or stomach pain
- ulcers
- bleeding problems
- high blood pressure
- heart or kidney disease
- taken a diuretic
- reached age 60 or older

Ask a doctor or pharmacist before use if you are:

- taking any other drug containing an NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug

When using this product do not exceed the recommended dose.

Stop use and ask a doctor if:

- ringing in the ears or loss of hearing occurs
- pain or fever persists or gets worse
- new symptoms occur
- redness or swelling is present
- you feel faint, vomit blood, or have bloody or black stools.
- These are signs of stomach bleeding.

If pregnant or breast-feeding baby, ask a health professional before use. IT IS ESPECIALLY IMPORTANT NOT TO USE ASPIRIN DURING THE LAST 3 MONTHS OF PREGNANCY UNLESS SPECIFICALLY DIRECTED TO DO SO BY A DOCTOR BECAUSE IT MAY CAUSE PROBLEMS IN THE UNBORN CHILD OR COMPLICATIONS DURING DELIVERY.

KEEP OUT OF REACH OF CHILDREN.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical

attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions:	2 tablets as needed for pain every 4 hours. Do not exceed 8 tablets in 24 hours or as directed by a doctor.
• Adults and children 12 years of age and older:	
• Children under 12 years:	Consult a doctor.

Other Information:

Tamper evident. Do not use if packet is torn, cut or opened Store at controlled room temperature 15° to 30°C (59° to 86° F) Avoid excessive heat and humidity

Inactive Ingredients:

FD&C Yellow #6, Magnesium Stearate, Microcrystalline Cellulose, Povidone, Starch, Stearic Acid

Product Package

aramark

Pain-Free®

Extra Strength Pain Reliever

Temporary relief of minor aches and pains associated with a cold, headache, toothache and muscular aches

Part # 537R23

100 TABLETS

PER BOX

Compare active ingredient to:

EXCEDRIN®

Registered Trademark of GlaxoSmithKline

Retain carton for complete product information


MANUFACTURED FOR:

Aramark


Lenexa, KS 66219

(913) 269-9611

100 Tablet Box



250 Tablet Box



2-Tablet packet



PAIN FREE

Pain Reliever
2 Tablets

Active Ingredients (In Each Tablet)	Purpose
Acetaminophen 110 mg.....	Pain Reliever
Aspirin 162 mg.....	Pain Reliever
Caffeine 32.4 mg.....	Stimulant
Salicylamide 152 mg.....	Pain Reliever

Uses: For the temporary relief of pain from: •headaches •colds •muscular aches •menstrual cramps •toothaches •backaches •minor arthritic pain

Warnings: Reye's Syndrome Warning: Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Reye's Syndrome, a rare but serious illness reported to be associated with aspirin.

Liver Warning: This product contains Acetaminophen. Severe liver damage may occur if you take: more than 8 tablets in 24 hours. • with other drugs containing acetaminophen (prescription or non-prescription) • 3 or more alcoholic drinks every day while using this product.

FOR COMPLETE WARNINGS/CAUTIONS: SEE BOX

Warnings (continued)
• 3 or more alcoholic drinks every day while using this product.

Stomach Bleeding Warning: This product contains nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. **The chance is higher if you:** have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • are age 60 or older • have 3 or more alcoholic drinks every day while using this product • take with other drugs containing an NSAID • take more or for a longer time than directed.

Keep out of reach of children.

Directions: Adults and children: (12 years and older) Take 1 to 2 tablets with a full glass of water every 4 hours as needed. Do not exceed 6 tablets in 24 hours, or as directed by doctor. Children: (under 12 years) Do not give to children under 12 years of age

Other Information: Store at a controlled room temperature 15°-30° (59°-86°F). Tamper evident: Do not use packet if torn or cut.

Contact: Aramark Lenexa, KS 66219

REV 3/2021

Made in USA

res

ARAMARK PAIN FREE

acetaminophen, aspirin, caffeine, salicylamide tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:81238-0242
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	110 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	162 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	32.4 mg
SALICYLAMIDE (UNII: EM8BM710ZC) (SALICYLAMIDE - UNII:EM8BM710ZC)	SALICYLAMIDE	152 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	

STEARIC ACID (UNII: 4ELV7Z65AP)

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	FR2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:81238-0242-1	50 in 1 BOX	05/07/2021	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:81238-0242-2	125 in 1 BOX	05/07/2021	
2		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	05/07/2021	

Labeler - Western First Aid Safety DBA Aramark (043861524)

Registrant - Western First Aid Safety DBA Aramark (043861524)

Revised: 6/2021

Western First Aid Safety DBA Aramark