

ACETAMINOPHEN EXTRA STRENGTH- acetaminophen tablet

Bi-Mart

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reaction. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning:

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**
- **adults and children 12 years and over:** take 2 caplets every 6 hours while symptoms last; do not take more than 6 caplets in 24 hours, unless directed by a doctor; do not take for more than 10 days unless directed by a doctor
- **children under 12 years:** ask a doctor

Other information

- store between 20-25°C (68-77°F)

Inactive ingredients

Hypromellose, Magnesium Stearate, Microcrystalline Cellulose , Pregelatinized Starch, Polyethylene Glycol, Povidone, Stearic Acid, Talc, Titanium Dioxide

Questions?

call toll-free **1-844-912-4012**


Distributed by:

BI-MART

Eugene, OR 97402

PRINCIPAL DISPLAY PANEL-100 Count

PRINCIPAL DISPLAY PANEL-250 Count



BI-MART
Compare to the active ingredient in Tylenol® Extra Strength*

ACETAMINOPHEN
EXTRA STRENGTH

• PAIN RELIEVER
• FEVER REDUCER
• NON-ASPIRIN

250 CAPLETS • 500 mg EACH

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Drug Facts

Active ingredient (in each caplet) *Purpose*
Acetaminophen 500 mg.....Pain reliever/fever reducer

Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ backache ■ minor pain of arthritis ■ the common cold ■ toothache ■ premenstrual and menstrual cramps ■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

Do not use ■ with any other products containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ **do not take more than directed (see overdose warning)**
■ **adults and children 12 years and over:** take 2 caplets every 6 hours while symptoms last; do not take more than 6 caplets in 24 hours, unless directed by a doctor; do not take for more than 10 days unless directed by a doctor
■ **children under 12 years:** ask a doctor

Other information
■ store between 20-25°C (68-77°F)

Inactive ingredients
Hydroxymethylcellulose, Magnesium Stearate, Microcrystalline Cellulose, Pregelatinized Starch, Polyethylene Glycol, Povidone, Stearic Acid, Talc, Titanium Dioxide.

Questions? call toll-free 1-844-912-4012

*This product is not manufactured or distributed by the owner of the registered trademark "Tylenol".


Distributed by:
BI-MART
Eugene, OR 97402

Questions?
call toll-free 1-844-912-4012

Inactive ingredients Hydroxymethylcellulose, Magnesium Stearate, Microcrystalline Cellulose, Pregelatinized Starch, Polyethylene Glycol, Povidone, Stearic Acid, Talc, Titanium Dioxide.

THIS PRODUCT IS NOT MANUFACTURED OR DISTRIBUTED BY THE OWNER OF THE REGISTERED TRADEMARK "TYPENOL".
TYPENOL, TYPENOL LOGO, AND TYPENOL TABLETS ARE NOT TO BE USED IN CONNECTION WITH THIS PRODUCT.
BIO-ART, BIO-ART LOGO, AND BIO-ART TABLETS ARE NOT TO BE USED IN CONNECTION WITH THIS PRODUCT.
RETURN THE UNUSED PORTION AND YOUR ENTIRE PURCHASE PRICE WILL BE CHEERFULLY REFUNDED.

PRINCIPAL DISPLAY PANEL-500 Count



BI-MART
Compare to the active ingredient in Tylenol® Extra Strength*

ACETAMINOPHEN
EXTRA STRENGTH

• PAIN RELIEVER
• FEVER REDUCER
• NON-ASPIRIN

100 CAPLETS • 500 mg EACH

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Drug Facts

Active ingredient (in each caplet) *Purpose*
Acetaminophen 500 mg.....Pain reliever/fever reducer

Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ backache ■ minor pain of arthritis ■ the common cold ■ toothache ■ premenstrual and menstrual cramps ■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

Do not use ■ with any other products containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ **do not take more than directed (see overdose warning)**
■ **adults and children 12 years and over:** take 2 caplets every 6 hours while symptoms last; do not take more than 6 caplets in 24 hours, unless directed by a doctor; do not take for more than 10 days unless directed by a doctor
■ **children under 12 years:** ask a doctor

Other information
■ store between 20-25°C (68-77°F)

Inactive ingredients
Hydroxymethylcellulose, Magnesium Stearate, Microcrystalline Cellulose, Pregelatinized Starch, Polyethylene Glycol, Povidone, Stearic Acid, Talc, Titanium Dioxide.

Questions? call toll-free 1-844-912-4012

*This product is not manufactured or distributed by the owner of the registered trademark "Tylenol".


Distributed by:
BI-MART
Eugene, OR 97402

Questions?
call toll-free 1-844-912-4012

Inactive ingredients Hydroxymethylcellulose, Magnesium Stearate, Microcrystalline Cellulose, Pregelatinized Starch, Polyethylene Glycol, Povidone, Stearic Acid, Talc, Titanium Dioxide.

THIS PRODUCT IS NOT MANUFACTURED OR DISTRIBUTED BY THE OWNER OF THE REGISTERED TRADEMARK "TYPENOL".
TYPENOL, TYPENOL LOGO, AND TYPENOL TABLETS ARE NOT TO BE USED IN CONNECTION WITH THIS PRODUCT.
BIO-ART, BIO-ART LOGO, AND BIO-ART TABLETS ARE NOT TO BE USED IN CONNECTION WITH THIS PRODUCT.
RETURN THE UNUSED PORTION AND YOUR ENTIRE PURCHASE PRICE WILL BE CHEERFULLY REFUNDED.

PRINCIPAL DISPLAY PANEL-250 Count



BI-MART
Compare to the active ingredient in Tylenol® Extra Strength*

ACETAMINOPHEN
EXTRA STRENGTH

• PAIN RELIEVER
• FEVER REDUCER
• NON-ASPIRIN

250 CAPLETS • 500 mg EACH

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Drug Facts

Active ingredient (in each caplet) *Purpose*
Acetaminophen 500 mg.....Pain reliever/fever reducer

Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ backache ■ minor pain of arthritis ■ the common cold ■ toothache ■ premenstrual and menstrual cramps ■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

Do not use ■ with any other products containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ **do not take more than directed (see overdose warning)**
■ **adults and children 12 years and over:** take 2 caplets every 6 hours while symptoms last; do not take more than 6 caplets in 24 hours, unless directed by a doctor; do not take for more than 10 days unless directed by a doctor
■ **children under 12 years:** ask a doctor

Other information
■ store between 20-25°C (68-77°F)

Inactive ingredients Hydroxymethylcellulose, Magnesium Stearate, Microcrystalline Cellulose, Pregelatinized Starch, Polyethylene Glycol, Povidone, Stearic Acid, Talc, Titanium Dioxide.

Questions? call toll-free 1-844-912-4012

*This product is not manufactured or distributed by the owner of the registered trademark "Tylenol".


Distributed by:
BI-MART
Eugene, OR 97402

Questions?
call toll-free 1-844-912-4012

Inactive ingredients Hydroxymethylcellulose, Magnesium Stearate, Microcrystalline Cellulose, Pregelatinized Starch, Polyethylene Glycol, Povidone, Stearic Acid, Talc, Titanium Dioxide.

THIS PRODUCT IS NOT MANUFACTURED OR DISTRIBUTED BY THE OWNER OF THE REGISTERED TRADEMARK "TYPENOL".
TYPENOL, TYPENOL LOGO, AND TYPENOL TABLETS ARE NOT TO BE USED IN CONNECTION WITH THIS PRODUCT.
BIO-ART, BIO-ART LOGO, AND BIO-ART TABLETS ARE NOT TO BE USED IN CONNECTION WITH THIS PRODUCT.
RETURN THE UNUSED PORTION AND YOUR ENTIRE PURCHASE PRICE WILL BE CHEERFULLY REFUNDED.

PRINCIPAL DISPLAY PANEL-500 Count



BI-MART
Compare to the active ingredient in Tylenol® Extra Strength*

ACETAMINOPHEN
EXTRA STRENGTH

• PAIN RELIEVER
• FEVER REDUCER
• NON-ASPIRIN

100 CAPLETS • 500 mg EACH

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Drug Facts

Active ingredient (in each caplet) *Purpose*
Acetaminophen 500 mg.....Pain reliever/fever reducer

Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ backache ■ minor pain of arthritis ■ the common cold ■ toothache ■ premenstrual and menstrual cramps ■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

Do not use ■ with any other products containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ **do not take more than directed (see overdose warning)**
■ **adults and children 12 years and over:** take 2 caplets every 6 hours while symptoms last; do not take more than 6 caplets in 24 hours, unless directed by a doctor; do not take for more than 10 days unless directed by a doctor
■ **children under 12 years:** ask a doctor

Other information
■ store between 20-25°C (68-77°F)

Inactive ingredients
Hydroxymethylcellulose, Magnesium Stearate, Microcrystalline Cellulose, Pregelatinized Starch, Polyethylene Glycol, Povidone, Stearic Acid, Talc, Titanium Dioxide.

Questions? call toll-free 1-844-912-4012

*This product is not manufactured or distributed by the owner of the registered trademark "Tylenol".

Distributed by:
BI-MART
Eugene, OR 97402

Questions?
call toll-free 1-844-912-4012

Inactive ingredients Hydroxymethylcellulose, Magnesium Stearate, Microcrystalline Cellulose, Pregelatinized Starch, Polyethylene Glycol, Povidone, Stearic Acid, Talc, Titanium Dioxide.

THIS PRODUCT IS NOT MANUFACTURED OR DISTRIBUTED BY THE OWNER OF THE REGISTERED TRADEMARK "TYPENOL".
TYPENOL, TYPENOL LOGO, AND TYPENOL TABLETS ARE NOT TO BE USED IN CONNECTION WITH THIS PRODUCT.
BIO-ART, BIO-ART LOGO, AND BIO-ART TABLETS ARE NOT TO BE USED IN CONNECTION WITH THIS PRODUCT.
RETURN THE UNUSED PORTION AND YOUR ENTIRE PURCHASE PRICE WILL BE CHEERFULLY REFUNDED.

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37835-507-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2025	
2	NDC:37835-507-25	250 in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2025	
3	NDC:37835-507-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/01/2025	

Labeler - Bi-Mart (027630078)**Registrant** - Bi-Mart (027630078)**Establishment**

Name	Address	ID/FEI	Business Operations
Elysium Pharmaceuticals Ltd.		915664486	manufacture(37835-507)

Revised: 5/2025

Bi-Mart