

IBUPROFEN- ibuprofen tablet, film coated
Lidl US LLC

Lidl US LLC Ibuprofen Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen,

or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**

Adults and children 12 years and older:

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor

Children under 12 years: ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, hypromellose, iron oxide red, iron oxide yellow, microcrystalline cellulose, polyethylene glycol, polysorbate 80, stearic acid, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the active ingredient in Advil[®] Ibuprofen Tablets

SEE NEW WARNINGS INFORMATION

Ibuprofen TABLETS 200mg

Actual Size

PAIN RELIEVER/FEVER REDUCER (NSAID)

250 tablets



Drug Facts (continued)

Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, polyethylene glycol, polyethylene glycol, polyethylene glycol, starch, titanium dioxide.

Questions or comments? 1-800-774-0980

This product is not transdermal and is distributed by Wirth Consumer Healthcare, distributor of Advion Ibuprofen Tablets.

DISTRIBUTED BY:
 LDI, LDC, LLC
 3500 S. CLARK STREET
 ARLINGTON, VA 22202

GLUTEN FREE

NDC 71141-007-85

Ibuprofen

TABLETS 200mg

**PAIN RELIEVER/
FEVER REDUCER
(NSAID)**



Compare to the active ingredient in Advion Ibuprofen Tablets.¹

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.

Drug Facts (for each tablet)

Active ingredient (in each tablet): Ibuprofen (200mg)

Purpose: Pain reliever/fever reducer

Uses:

- temporarily relieves minor aches and pains due to:
- headache
- minor pain of arthritis
- the common cold
- temporarily reduces fever
- muscle aches
- toothache
- backache
- menstrual cramps

Warnings:

Alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- skin rash
- severe dizziness
- trouble breathing

If an allergic reaction to aspirin occurs, it may be more likely to occur with ibuprofen. Symptoms may include:

- stomach bleeding/warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you are age 65 or older, have had stomach ulcer or bleeding problems
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take other drugs containing prescription or nonprescription NSAIDs (ibuprofen, naproxen, or others)
- have 5 or more ulcers or bleeding problems every day while using this product
- have more or for a longer time than directed
- Do not use:
 - if you have ever had an allergic reaction to any other pain reliever/fever reducer
 - right before or after heart surgery


Ask a doctor before use if:

- you have problems or serious side effects from taking pain relievers of their kind
- you have been taking other drugs such as aspirin, ibuprofen, naproxen, or others
- you have a history of stomach problems, such as heartburn

Do not use:

- you have high blood pressure, heart disease, liver disease, or kidney disease
- you are taking a diuretic

: 6045 F F 1



Compare to the active ingredient in Advion Ibuprofen Tablets.¹

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are:

- taking other drugs such as aspirin, ibuprofen, naproxen, or others
- taking other drugs such as aspirin, ibuprofen, naproxen, or others
- taking other drugs such as aspirin, ibuprofen, naproxen, or others

When using this product:

- take with food or milk
- do not take more than directed
- stop use and ask a doctor if you experience any of the following signs or symptoms:
- stomach bleeding/warning: This product contains an NSAID, which may cause severe stomach bleeding.
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 5 or more ulcers or bleeding problems every day while using this product
- have more or for a longer time than directed

Stop use and ask a doctor if:

- you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Do not use:

- you have high blood pressure, heart disease, liver disease, or kidney disease
- you are taking a diuretic

Directions:

Do not take more than directed.

The smallest amount should be used.

Adults and children 12 years and older:

- take 1 tablet every 4 to 6 hours while symptoms persist.
- If pain or fever does not respond to 1 tablet, 2 tablets may be used.
- do not exceed 6 tablets in 24 hours, unless directed by a doctor.
- Children under 12 years: ask a doctor.

Other information:

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 30°C (86°F)

IBUPROFEN		
ibuprofen tablet, film coated		
Product Information		
Product Type	HUMAN OTC DRUG	Item Code (Source)
Route of Administration	ORAL	NDC:71141-007
Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg
Inactive Ingredients		
Ingredient Name	Strength	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		
STARCH, CORN (UNII: O8232NY3SJ)		
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)		
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3W0)		
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		
POLYSORBATE 80 (UNII: 6OZP39ZG8H)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
FERRIC OXIDE RED (UNII: 1K09F3G675)		
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)		

Product Characteristics

Color	BROWN	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	I2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71141-007-85	250 in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA072096	03/15/2017	

Labeler - Lidl US LLC (079389709)

Revised: 11/2019

Lidl US LLC