

COLD AND COUGH CHILDRENS- brompheniramine maleate, dextromethorphan hbr, phenylephrine hcl liquid
Rite Aid Corporation

Drug Facts

Active ingredients (in each 10 mL)

Brompheniramine Maleate 2 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Purposes

Antihistamine

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (Allergic rhinitis)
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily restores freer breathing through the nose

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease
- high blood pressure
- diabetes

- glaucoma
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

When using this product,

- **do not use more than directed**
- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be a sign of a serious condition

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- mL = milliliter

Age	Dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to 11 years	10 mL every 4 hours
children under 6 years	do not use

Other information

- **each 10 mL contains:** sodium 5 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose

Principal Display Panel

Compare to the active ingredients in Children's Dimetapp® Cold & Cough*

6 YEARS & OLDER

CHILDREN'S**COLD & COUGH**

BROMPHENIRAMINE MALEATE 2 mg

DEXTROMETHORPHAN HBr 10 mg

PHENYLEPHRINE HCl 5 mg

ANTIHISTAMINE

COUGH SUPPRESSANT

NASAL DECONGESTANT

relieves:

cough • itchy, watery eyes

runny nose • sneezing

stuffy nose

itching of the nose or throat

GRAPE FLAVOR

DOSAGE CUP INCLUDED

FL OZ (mL)

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Children's Dimetapp® Cold & Cough.

DISTRIBUTED BY:

RITE AID

30 HUNTER LANE

CAMP HILL, PA 17011

Product Label

Drug Facts (continued)

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product.
- mL = milliliter

Age	Dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to 11 years	10 mL every 4 hours
children under 6 years	do not use

Other information

- each 10 mL contains: sodium 5 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose

Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Children's Dimetapp® Cold & Cough.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org



DOSAGE CUP INCLUDED

DISTRIBUTED BY:
RITE AID
30 HUNTER LANE
CAMP HILL, PA 17011
www.riteaid.com

SATISFACTION GUARANTEE
If you're not satisfied, we'll happily refund your money.

NDC 11822-5259-3

Compare to the active ingredients in Children's Dimetapp® Cold & Cough*

FREE ALCOHOL FREE

6 YEARS & OLDER

CHILDREN'S COLD & COUGH

BROMPHENIRAMINE MALEATE 2 mg
DEXTROMETHORPHAN HBr 10 mg
PHENYLEPHRINE HCl 5 mg
ANTI-HISTAMINE COUGH SUPPRESSANT NASAL DECONGESTANT

relieves:
cough • itchy, watery eyes
runny nose • sneezing
stuffy nose
itching of the nose or throat




GRAPE FLAVOR
DOSAGE CUP INCLUDED
4 FL OZ (118 mL)

FREE ALCOHOL FREE

NDC 11822-5259-3

Compare to the active ingredients in Children's Dimetapp® Cold & Cough*

FREE ALCOHOL FREE

6 YEARS & OLDER

CHILDREN'S COLD & COUGH

BROMPHENIRAMINE MALEATE 2 mg
DEXTROMETHORPHAN HBr 10 mg
PHENYLEPHRINE HCl 5 mg
ANTI-HISTAMINE COUGH SUPPRESSANT NASAL DECONGESTANT

relieves:
cough • itchy, watery eyes
runny nose • sneezing
stuffy nose
itching of the nose or throat




GRAPE FLAVOR
DOSAGE CUP INCLUDED
4 FL OZ (118 mL)

Drug Facts

Active ingredients (in each 10 mL)

Brompheniramine maleate 2 mg	Antihistamine
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis)
- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat
- temporarily restores freer breathing through the nose

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.



0 11822 00093 2

PLD-6261D F0005817

Lot No.:
Exp. Date:

RITE AID Children's Cold & Cough Grape Flavor

COLD AND COUGH CHILDRENS

brompheniramine maleate, dextromethorphan hbr, phenylephrine hcl liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-5259
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BROMPHENIRAMINE MALEATE (UNII: IXA7C9ZN03) (BROMPHENIRAMINE - UNII:H57G17P2FN)	BROMPHENIRAMINE MALEATE	2 mg in 10 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-5259-3	1 in 1 BOX	11/05/2021	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/05/2021	

Labeler - Rite Aid Corporation (014578892)