

ANTI-DIARRHEAL- loperamide hydrochloride capsule, liquid filled
CHAIN DRUG MARKETING ASSOCIATION INC.

Loperamide HCl Capsules - Quality Choice

Active ingredient (in each capsule)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

- if you have bloody or black stool
- in children under 12 years of age

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea.**
- **not for use in children under 12 years of age**
- adults and children 12 years and over: 2 capsules after the first loose stool; 1 capsule after each subsequent loose stool; but no more than 4 capsules in 24 hours

Other information

- store at 20°-25°C (68°-77°F). Protect from Light.
- avoid excessive heat above 40°C (104°F)
- do not use if carton or blister unit is open or torn
- see side panel for lot number and expiration date

Inactive ingredients

butylated hydroxyanisole, edible ink, FD&C Blue #1, gelatin, glycerin, glyceryl caprylate, polyoxyl 40 hydrogenated castor oil, purified water

Questions or comments?

call toll free **1-888-235-2466**

THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER EVIDENT PACKAGE.
USE ONLY IF BLISTERS ARE INTACT.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Imodium ® A-D

L0000928

Lot #

Expiration Date

R1024

Distributed by CDMA, Inc.

Novi, MI 48375

www.qualitychoice.com

Questions: 800-935-2362

Principal Display Panel

QC ®

QUALITY CHOICE

NDC 83324-208-12

***Compare to the**

Active Ingredient in

Imodium® A-D

Anti-Diarrheal

Loperamide HCl Soft Gelatin Capsules, 2 mg

Controls the Symptoms of Diarrhea

Suitable for Adults and Children

12 Years and Over

†Each Liquid-Filled Capsule Contains 2 mg Loperamide HCl

12 Softgels †

Drug Facts

Active ingredient (in each capsule)
Loperamide HCl 2 mg

Purpose
Anti-diarrheal

Use
Controls symptoms of diarrhea, including Traveler's Diarrhea

Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.
Heart alert: Taking more than directed can cause serious heart problems or death.
Do not use if you have bloody or black stool or if you have been told by a doctor that you have a history of abnormal heart rhythm.
Do not use if you have fever, mucus in the stool, or a history of liver disease.

Ask a doctor before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product (drowsiness, dizziness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if:
■ symptoms get worse ■ diarrhea lasts for more than 2 days
■ you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ Drink plenty of clear fluids to help prevent dehydration caused by diarrhea.
■ not for use in children under 12 years of age
■ adults and children 12 years and over: 2 capsules after the first loose stool; 1 capsule after each subsequent loose stool, but no more than 4 capsules in 24 hours

Other information
■ store at 20°-25°C (68°-77°F). Protect from light.
■ avoid excessive heat above 40°C (104°F).
■ do not use if carton or blister unit is open or torn
■ see side panel for lot number and expiration date

Inactive ingredients
butylated hydroxyanisole, edible ink, FD&C Blue #1, gelatin, glycerin, glyceryl caprylate, polyoxy 40 hydrogenated castor oil, purified water

Questions or comments? call toll free 1-888-235-2466



Anti-Diarrheal

Loperamide HCl Soft Gelatin Capsules, 2 mg

Controls the Symptoms of Diarrhea

NDC 83324-208-12



*Compare to the
Active Ingredient in
Imodium® A-D

Anti-Diarrheal

Loperamide HCl Soft Gelatin Capsules, 2 mg

Controls the Symptoms of Diarrhea

Suitable for Adults and Children
12 Years and Over

*Each Liquid-Filled Capsule Contains 2 mg Loperamide HCl

12 Softgels*



Actual Size



Distributed by CDMA, Inc.
Novi, MI 48375
www.qualitychoice.com
Questions: 800-935-2362

*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Imodium® A-D

L0011523
Lot #
Expiration Date
R1024



See package for complete directions. Do not use if you have been told by a doctor that you have a history of abnormal heart rhythm.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

ANTI-DIARRHEAL				
loperamide hydrochloride capsule, liquid filled				
Product Information				
Product Type		HUMAN OTC DRUG	Item Code (Source)	
Route of Administration		ORAL	NDC:83324-208	
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)			LOPERAMIDE HYDROCHLORIDE	2 mg
Inactive Ingredients				
Ingredient Name				Strength
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
GELATIN (UNII: 2G86QN327L)				
GLYCERIN (UNII: PDC6A3C0OX)				
GLYCERYL CAPRYLATE (UNII: TM2TZD4G4A)				
POLYOXYL 40 HYDROGENATED CASTOR OIL (UNII: 7YC686GQ8F)				
WATER (UNII: 059QF0KO0R)				
Product Characteristics				
Color	blue	Score	no score	
Shape	CAPSULE	Size	10mm	
Flavor		Imprint Code	LP2	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83324-208-12	1 in 1 CARTON	05/21/2025	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA		NDA021855	05/21/2025	

Labeler - CHAIN DRUG MARKETING ASSOCIATION INC. (011920774)

Registrant - Bionpharma Inc. (079637826)

Establishment

Name	Address	ID/FEI	Business Operations
Patheon Softgels Inc.		002193829	manufacture(83324-208)

Revised: 5/2025

CHAIN DRUG MARKETING ASSOCIATION INC.