

GLYCERIN LAXATIVE- glycerin laxative suppositories suppository
Neilmed Pharmaceuticals Inc.

Glycerin Laxative foil wrapped Suppositories

Drug Facts: Active Ingredient

Active ingredient (in each suppository)
Glycerin, USP 2 grams

Drug Facts

Active ingredient (in each suppository)Purpose
Glycerin, USP 2 grams Laxative

Uses

- Relieves occasional constipation (irregularity)
- Generally produces bowel movement in 15 minutes to 1 hour

Warnings

For rectal use only

Do not use

laxative products for longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- Abdominal pain • Nausea • Vomiting • Noticed a sudden change in bowel habits that lasts over 2 weeks

When using the product

you may have rectal discomfort or a burning sensation.

Stop use and ask a doctor if

- You have rectal bleeding
- You fail to have a bowel movement after use of a laxative. These could be a signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

Adults and children 6 years of age and over	1 suppository per day or as directed by a doctor.
Children 2 to under 6 years	Use Neilmed Child Suppositories
Children under 2 years	Ask a doctor

Insert 1 suppository well into the rectum. To produce laxative action, suppository is designed to dissolve only partially and may not be noticeable.

Other Information

- Store at room temperature. Do not exceed 30° C (86° F)
- Lot No. & Exp. Date: see wrapper or box

Inactive ingredients

Purified water, sodium hydroxide, stearic acid

Questions?

Questions or comments? 1 (877) 477-8633

Important:

Please do not discard this printed box and any enclosed printed material. The inside final product may not have all the details you require for the ongoing use of the product.



Package Label

GLYCERIN LAXATIVE

glycerin laxative suppositories suppository

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:13709-329
Route of Administration	RECTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GLYCERIN (UNII: PDC6A3C0OX) (GLYCERIN - UNII:PDC6A3C0OX)	GLYCERIN	2 g in 2.5 g

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

STEARIC ACID (UNII: 4ELV7Z65AP)				
SODIUM HYDROXIDE (UNII: 55X04QC32I)				
WATER (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13709-329-01	30 in 1 CARTON	05/23/2025	
1		2.5 g in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M015	05/23/2025	

Labeler - Neilmed Pharmaceuticals Inc. (799295915)

Establishment

Name	Address	ID/FEI	Business Operations
NeilMed Pharmaceuticals Inc.		799295915	manufacture(13709-329)

Revised: 5/2025

Neilmed Pharmaceuticals Inc.