

NUMB CREAM- lidocaine paste

FlexiGo Inc



**TOPICAL PAIN TREATMENT
FOR TATTOOS**

- Recommended Steps for Treating before Tattooing:**
1. Clean your hands and the treatment area with soft soap and warm water, rinse thoroughly, then gently pat dry.
 2. Apply a thick layer of the numbing cream onto the treatment area.
 3. Wrap lightly with plastic wrap and wait 45-60 minutes for the numbness to peak. (The peak lasts up to 6 hours.)
 4. Wipe away the residue with a soft cloth or tissue.
 5. Begin the procedure.

Notice:

- ☑ Please be aware that the duration of the numbing effect may vary depending on factors such as temperature and individual skin types.
- ☑ Additionally, it is generally observed that longer wrapping times tend to produce better results in terms of numbness.
- ☑ Testing in a small area first is recommended to ensure the best numbness effect.



6
HOURS

ENT. WT.
1.7 OZ / 50ML

NUMBING WITH EMU OIL AND ARNICA

DRUG FACTS

ACTIVE INGREDIENT:	PURPOSE
LILOCAINE 4%	External Anesthetic
USES: For temporary relief of local and momentary discomfort associated with unexcited treatment or information	
WARNINGS:	
• For External Use Only	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
DO NOT USE:	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
WHEN USING:	
Do not exceed the recommended daily dosage unless directed by a doctor.	
Certain persons can develop allergic reactions to ingredients in this product.	
Do not put this product into the rectum by using fingers or any medical device or applicator.	

DRUG FACTS (continued)

STOP USE:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
ASK DOCTOR:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
KEEP OUT OF REACH OF CHILDREN:
In case of accidental ingestion, seek medical attention immediately.
DIRECTIONS:
Perform a patch test on the back of your hand before using it to check for skin sensitivity. When practice, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product.
Apply as to 2 times a day.
OTHER INFORMATION:
Keep away from direct sunlight or heat.
Store at room temperature (20-25°C / 68-77°F).
INACTIVE INGREDIENTS:
WATER, PROPYLENE GLYCOL, CARBOMER, GLYCERIN, DIODIUM EDTA, HYDROXY-ACETOPHENONE.

**TOPICAL PAIN TREATMENT
FOR TATTOOS**

- Recommended Steps for Treating before Tattooing:**
1. Clean your hands and the treatment area with soft soap and warm water, rinse thoroughly, then gently pat dry.
 2. Apply a thick layer of the numbing cream onto the treatment area.
 3. Wrap lightly with plastic wrap and wait 45-60 minutes for the numbness to peak. (The peak lasts up to 6 hours.)
 4. Wipe away the residue with a soft cloth or tissue.
 5. Begin the procedure.

Notice:

- ☑ Please be aware that the duration of the numbing effect may vary depending on factors such as temperature and individual skin types.
- ☑ Additionally, it is generally observed that longer wrapping times tend to produce better results in terms of numbness.
- ☑ Testing in a small area first is recommended to ensure the best numbness effect.



DRUG FACTS

ACTIVE INGREDIENT:	PURPOSE
LILOCAINE 4%	External Anesthetic
USES: For temporary relief of local and momentary discomfort associated with anesthetic treatment or information	
WARNINGS:	
• For External Use Only	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
DO NOT USE:	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
WHEN USING:	
Do not exceed the recommended daily dosage unless directed by a doctor.	
Certain persons can develop allergic reactions to ingredients in this product.	
Do not put this product into the rectum by using fingers or any medical device or applicator.	

DRUG FACTS (continued)

STOP USE:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
ASK DOCTOR:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
KEEP OUT OF REACH OF CHILDREN:
In case of accidental ingestion, seek medical attention immediately.
DIRECTIONS:
Perform a patch test on the back of your hand before using it to check for skin sensitivity. When practice, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with clean tissue or a soft cloth before application of this product.
Apply as to 2 times a day.
OTHER INFORMATION:
Keep away from direct sunlight or heat.
Store at room temperature (20-25°C / 68-77°F).
INACTIVE INGREDIENTS:
WATER, PROPYLENE GLYCOL, CARBOMER, GLYCERIN, DIODIUM EDTA, HYDROXY-ACETOPHENONE.

**TOPICAL PAIN TREATMENT
FOR TATTOOS**

- Recommended Steps for Treating before Tattooing:**
1. Clean your hands and the treatment area with soft soap and warm water, rinse thoroughly, then gently pat dry.
 2. Apply a thick layer of the numbing cream onto the treatment area.
 3. Wrap lightly with plastic wrap and wait 45-60 minutes for the numbness to peak. (The peak lasts up to 6 hours.)
 4. Wipe away the residue with a soft cloth or tissue.
 5. Begin the procedure.

Notice:

- ☑ Please be aware that the duration of the numbing effect may vary depending on factors such as temperature and individual skin types.
- ☑ Additionally, it is generally observed that longer wrapping times tend to produce better results in terms of numbness.
- ☑ Testing in a small area first is recommended to ensure the best numbness effect.



6
HOURS

ENT. WT.
1.7 OZ / 50ML

NUMBING WITH EMU OIL AND ARNICA

DRUG FACTS

ACTIVE INGREDIENT:	PURPOSE
LILOCAINE 4%	External Anesthetic
USES: For temporary relief of local and momentary discomfort associated with unexcited treatment or inflammation	
WARNINGS:	
• For External Use Only	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
DO NOT USE:	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
WHEN USING:	
Do not exceed the recommended daily dosage unless directed by a doctor.	
Certain persons can develop allergic reactions to ingredients in this product.	
Do not put this product into the rectum by using fingers or any medical device or applicator.	

DRUG FACTS (continued)

STOP USE:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
ASK DOCTOR:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
KEEP OUT OF REACH OF CHILDREN:
In case of accidental ingestion, seek medical attention immediately.
DIRECTIONS:
Perform a patch test on the back of your hand before using it to check for skin sensitivity. When practice, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with clean tissue or a soft cloth before application of this product.
Apply as to 2 times a day.
OTHER INFORMATION:
Keep away from direct sunlight or heat.
Store at room temperature (20-25°C / 68-77°F).
INACTIVE INGREDIENTS:
WATER, PROPYLENE GLYCOL, CARBOMER, GLYCERIN, DIODIUM EDTA, HYDROXY-ACETOPHENONE.

**TOPICAL PAIN TREATMENT
FOR TATTOOS**

- Recommended Steps for Treating before Tattooing:**
1. Clean your hands and the treatment area with soft soap and warm water, rinse thoroughly, then gently pat dry.
 2. Apply a thick layer of the numbing cream onto the treatment area.
 3. Wrap lightly with plastic wrap and wait 45-60 minutes for the numbness to peak. (The peak lasts up to 6 hours.)
 4. Wipe away the residue with a soft cloth or tissue.
 5. Begin the procedure.

Notice:

- ☑ Please be aware that the duration of the numbing effect may vary depending on factors such as temperature and individual skin types.
- ☑ Additionally, it is generally observed that longer wrapping times tend to produce better results in terms of numbness.
- ☑ Testing in a small area first is recommended to ensure the best numbness effect.



6
HOURS

ENT. WT.
1.7 OZ / 50ML

NUMBING WITH EMU OIL AND ARNICA

DRUG FACTS

ACTIVE INGREDIENT:	PURPOSE
LILOCAINE 4%	External Anesthetic
USES: For temporary relief of local and momentary discomfort associated with unexcited treatment or inflammation	
WARNINGS:	
• For External Use Only	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
DO NOT USE:	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
WHEN USING:	
Do not exceed the recommended daily dosage unless directed by a doctor.	
Certain persons can develop allergic reactions to ingredients in this product.	
Do not put this product into the rectum by using fingers or any medical device or applicator.	

DRUG FACTS (continued)

STOP USE:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
ASK DOCTOR:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
KEEP OUT OF REACH OF CHILDREN:
In case of accidental ingestion, seek medical attention immediately.
DIRECTIONS:
Perform a patch test on the back of your hand before using it to check for skin sensitivity. When practice, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with clean tissue or a soft cloth before application of this product.
Apply as to 2 times a day.
OTHER INFORMATION:
Keep away from direct sunlight or heat.
Store at room temperature (20-25°C / 68-77°F).
INACTIVE INGREDIENTS:
WATER, PROPYLENE GLYCOL, CARBOMER, GLYCERIN, DIODIUM EDTA, HYDROXY-ACETOPHENONE.

**TOPICAL PAIN TREATMENT
FOR TATTOOS**

- Recommended Steps for Treating before Tattooing:**
1. Clean your hands and the treatment area with soft soap and warm water, rinse thoroughly, then gently pat dry.
 2. Apply a thick layer of the numbing cream onto the treatment area.
 3. Wrap lightly with plastic wrap and wait 45-60 minutes for the numbness to peak. (The peak lasts up to 6 hours.)
 4. Wipe away the residue with a soft cloth or tissue.
 5. Begin the procedure.

Notice:

- ☑ Please be aware that the duration of the numbing effect may vary depending on factors such as temperature and individual skin types.
- ☑ Additionally, it is generally observed that longer wrapping times tend to produce better results in terms of numbness.
- ☑ Testing in a small area first is recommended to ensure the best numbness effect.



6
HOURS

ENT. WT.
1.7 OZ / 50ML

NUMBING WITH EMU OIL AND ARNICA

DRUG FACTS

ACTIVE INGREDIENT:	PURPOSE
LILOCAINE 4%	External Anesthetic
USES: For temporary relief of local and momentary discomfort associated with unexcited treatment or inflammation	
WARNINGS:	
• For External Use Only	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
DO NOT USE:	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
WHEN USING:	
Do not exceed the recommended daily dosage unless directed by a doctor.	
Certain persons can develop allergic reactions to ingredients in this product.	
Do not put this product into the rectum by using fingers or any medical device or applicator.	

DRUG FACTS (continued)

STOP USE:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
ASK DOCTOR:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
KEEP OUT OF REACH OF CHILDREN:
In case of accidental ingestion, seek medical attention immediately.
DIRECTIONS:
Perform a patch test on the back of your hand before using it to check for skin sensitivity. When practice, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with clean tissue or a soft cloth before application of this product.
Apply as to 2 times a day.
OTHER INFORMATION:
Keep away from direct sunlight or heat.
Store at room temperature (20-25°C / 68-77°F).
INACTIVE INGREDIENTS:
WATER, PROPYLENE GLYCOL, CARBOMER, GLYCERIN, DIODIUM EDTA, HYDROXY-ACETOPHENONE.

**TOPICAL PAIN TREATMENT
FOR TATTOOS**

- Recommended Steps for Treating before Tattooing:**
1. Clean your hands and the treatment area with soft soap and warm water, rinse thoroughly, then gently pat dry.
 2. Apply a thick layer of the numbing cream onto the treatment area.
 3. Wrap lightly with plastic wrap and wait 45-60 minutes for the numbness to peak. (The peak lasts up to 6 hours.)
 4. Wipe away the residue with a soft cloth or tissue.
 5. Begin the procedure.

Notice:

- ☑ Please be aware that the duration of the numbing effect may vary depending on factors such as temperature and individual skin types.
- ☑ Additionally, it is generally observed that longer wrapping times tend to produce better results in terms of numbness.
- ☑ Testing in a small area first is recommended to ensure the best numbness effect.



NUMBING WITH EMU OIL AND ARNICA

DRUG FACTS

ACTIVE INGREDIENT:	PURPOSE
Locaine 4%	External Anesthetic
USES: For temporary relief of local and momentary discomfort associated with unexcited treatment or inflammation	
WARNINGS:	
• For External Use Only	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
DO NOT USE:	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
WHEN USING:	
Do not exceed the recommended daily dosage unless directed by a doctor.	
Certain persons can develop allergic reactions to ingredients in this product.	
Do not put this product into the rectum by using fingers or any medical device or applicator.	

DRUG FACTS (continued)

STOP USE:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
ASK DOCTOR:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
KEEP OUT OF REACH OF CHILDREN:
In case of accidental ingestion, seek medical attention immediately.
DIRECTIONS:
Perform a patch test on the back of your hand before using it to check for skin sensitivity. When practice, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product.
Apply as to 2 times a day.
OTHER INFORMATION:
Keep away from direct sunlight or heat.
Store at room temperature (20-25°C / 68-77°F).
INACTIVE INGREDIENTS:
WATER, PROPYLENE GLYCOL, CARBOMER, GLYCERIN, DIODIUM EDTA, HYDROXY-ACETOPHENONE.

**TOPICAL PAIN TREATMENT
FOR TATTOOS**

- Recommended Steps for Treating before Tattooing:**
1. Clean your hands and the treatment area with soft soap and warm water, rinse thoroughly, then gently pat dry.
 2. Apply a thick layer of the numbing cream onto the treatment area.
 3. Wrap lightly with plastic wrap and wait 45-60 minutes for the numbness to peak. (The peak lasts up to 6 hours.)
 4. Wipe away the residue with a soft cloth or tissue.
 5. Begin the procedure.

Notice:

- ☑ Please be aware that the duration of the numbing effect may vary depending on factors such as temperature and individual skin types.
- ☑ Additionally, it is generally observed that longer wrapping times tend to produce better results in terms of numbness.
- ☑ Testing in a small area first is recommended to ensure the best numbness effect.



NUMBING WITH EMU OIL AND ARNICA

DRUG FACTS

ACTIVE INGREDIENT:	PURPOSE
Locaine 4%	External Anesthetic
USES: For temporary relief of local and momentary discomfort associated with unexcited treatment or inflammation	
WARNINGS:	
• For External Use Only	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
DO NOT USE:	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
WHEN USING:	
Do not exceed the recommended daily dosage unless directed by a doctor.	
Certain persons can develop allergic reactions to ingredients in this product.	
Do not put this product into the rectum by using fingers or any medical device or applicator.	

DRUG FACTS (continued)

STOP USE:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
ASK DOCTOR:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
KEEP OUT OF REACH OF CHILDREN:
In case of accidental ingestion, seek medical attention immediately.
DIRECTIONS:
Perform a patch test on the back of your hand before using it to check for skin sensitivity. When practice, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product.
Apply as to 2 times a day.
OTHER INFORMATION:
Keep away from direct sunlight or heat.
Store at room temperature (20-25°C / 68-77°F).
INACTIVE INGREDIENTS:
WATER, PROPYLENE GLYCOL, CARBOMER, GLYCERIN, DIODIUM EDTA, HYDROXY-ACETOPHENONE.

**TOPICAL PAIN TREATMENT
FOR TATTOOS**

- Recommended Steps for Treating before Tattooing:**
1. Clean your hands and the treatment area with soft soap and warm water, rinse thoroughly, then gently pat dry.
 2. Apply a thick layer of the numbing cream onto the treatment area.
 3. Wrap lightly with plastic wrap and wait 45-60 minutes for the numbness to peak. (The peak lasts up to 6 hours.)
 4. Wipe away the residue with a soft cloth or tissue.
 5. Begin the procedure.

Notice:

- ☑ Please be aware that the duration of the numbing effect may vary depending on factors such as temperature and individual skin types.
- ☑ Additionally, it is generally observed that longer wrapping times tend to produce better results in terms of numbness.
- ☑ Testing in a small area first is recommended to ensure the best numbness effect.



NUMBING WITH EMU OIL AND ARNICA

DRUG FACTS

ACTIVE INGREDIENT:	PURPOSE
LILOCAINE 4%	External Anesthetic
USES: For temporary relief of local and momentary discomfort associated with unexcited treatment or information	
WARNINGS:	
• For External Use Only	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
DO NOT USE:	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
WHEN USING:	
Do not exceed the recommended daily dosage unless directed by a doctor.	
Certain persons can develop allergic reactions to ingredients in this product.	
Do not put this product into the rectum by using fingers or any medical device or applicator.	

DRUG FACTS (continued)

STOP USE:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
ASK DOCTOR:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
KEEP OUT OF REACH OF CHILDREN:
In case of accidental ingestion, seek medical attention immediately.
DIRECTIONS:
Perform a patch test on the back of your hand before using it to check for skin sensitivity. When practice, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with clean tissue or a soft cloth before application of this product.
Apply as to 2 times a day.
OTHER INFORMATION:
Keep away from direct sunlight or heat.
Store at room temperature (20-25°C / 68-77°F).
INACTIVE INGREDIENTS:
WATER, PROPYLENE GLYCOL, CARBOMER, GLYCERIN, DIODIUM EDTA, HYDROXY-ACETOPHENONE.

**TOPICAL PAIN TREATMENT
FOR TATTOOS**

- Recommended Steps for Treating before Tattooing:**
1. Clean your hands and the treatment area with soft soap and warm water, rinse thoroughly, then gently pat dry.
 2. Apply a thick layer of the numbing cream onto the treatment area.
 3. Wrap lightly with plastic wrap and wait 45-60 minutes for the numbness to peak. (The peak lasts up to 6 hours.)
 4. Wipe away the residue with a soft cloth or tissue.
 5. Begin the procedure.

Notice:

- ☑ Please be aware that the duration of the numbing effect may vary depending on factors such as temperature and individual skin types.
- ☑ Additionally, it is generally observed that longer wrapping times tend to produce better results in terms of numbness.
- ☑ Testing in a small area first is recommended to ensure the best numbness effect.



NUMBING WITH EMU OIL AND ARNICA

DRUG FACTS

ACTIVE INGREDIENT:	PURPOSE
LILOCAINE 4%	External Anesthetic
USES: For temporary relief of local and momentary discomfort associated with anesthetic treatment or information	
WARNINGS:	
• For External Use Only	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
DO NOT USE:	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
WHEN USING:	
Do not exceed the recommended daily dosage unless directed by a doctor.	
Certain persons can develop allergic reactions to ingredients in this product.	
Do not put this product into the rectum by using fingers or any medical device or applicator.	

DRUG FACTS (continued)

STOP USE:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
ASK DOCTOR:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
KEEP OUT OF REACH OF CHILDREN:
In case of accidental ingestion, seek medical attention immediately.
DIRECTIONS:
Perform a patch test on the back of your hand before using it to check for skin sensitivity. When practice, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with clean tissue or a soft cloth before application of this product.
Apply as to 2 times a day.
OTHER INFORMATION:
Keep away from direct sunlight or heat.
Store at room temperature (20-25°C / 68-77°F).
INACTIVE INGREDIENTS:
WATER, PROPYLENE GLYCOL, CARBOMER, GLYCERIN, DIODIUM EDTA, HYDROXY-ACETOPHENONE.

**TOPICAL PAIN TREATMENT
FOR TATTOOS**

- Recommended Steps for Treating before Tattooing:**
1. Clean your hands and the treatment area with soft soap and warm water, rinse thoroughly, then gently pat dry.
 2. Apply a thick layer of the numbing cream onto the treatment area.
 3. Wrap lightly with plastic wrap and wait 45-60 minutes for the numbness to peak. (The peak lasts up to 6 hours.)
 4. Wipe away the residue with a soft cloth or tissue.
 5. Begin the procedure.

Notice:

- ☑ Please be aware that the duration of the numbing effect may vary depending on factors such as temperature and individual skin types.
- ☑ Additionally, it is generally observed that longer wrapping times tend to produce better results in terms of numbness.
- ☑ Testing in a small area first is recommended to ensure the best numbness effect.



NUMBING WITH EMU OIL AND ARNICA

DRUG FACTS

ACTIVE INGREDIENT:	PURPOSE
LILOCAINE 4%	External Anesthetic
USES: For temporary relief of local and momentary discomfort associated with unexcited treatment or inflammation	
WARNINGS:	
• For External Use Only	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
DO NOT USE:	
• Pregnant or breastfeeding, ask a health professional before use.	
• In case of accidental overdose, contact a doctor or Poison Control Center immediately.	
• Seal is broken or missing.	
WHEN USING:	
Do not exceed the recommended daily dosage unless directed by a doctor.	
Certain persons can develop allergic reactions to ingredients in this product.	
Do not put this product into the rectum by using fingers or any medical device or applicator.	

DRUG FACTS (continued)

STOP USE:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
ASK DOCTOR:
The symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase.
KEEP OUT OF REACH OF CHILDREN:
In case of accidental ingestion, seek medical attention immediately.
DIRECTIONS:
Perform a patch test on the back of your hand before using it to check for skin sensitivity. When practice, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with clean tissue or a soft cloth before application of this product.
Apply as to 2 times a day.
OTHER INFORMATION:
Keep away from direct sunlight or heat.
Store at room temperature (20-25°C / 68-77°F).
INACTIVE INGREDIENTS:
WATER, PROPYLENE GLYCOL, CARBOMER, GLYCERIN, DIODIUM EDTA, HYDROXY-ACETOPHENONE.



NUMB CREAM

lidocaine paste

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:85524-2506
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety		
Ingredient Name		Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)		0.04 g in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
EDETATE DISODIUM (UNII: 7FLD91C86K)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
GLYCERIN (UNII: PDC6A3C0OX)	

HYDROXYACETOPHENONE (UNII: G1L3HT4CMH)				
WATER (UNII: 059QF0KO0R)				
Product Characteristics				
Color		white	Score	
Shape			Size	
Flavor			Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85524-2506-5	50 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/26/2025	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M015	05/25/2025	

Labeler - FlexiGo Inc (118951527)

Establishment			
Name	Address	ID/FEI	Business Operations
FlexiGo Inc		118951527	manufacture(85524-2506) , label(85524-2506)

Revised: 5/2025

FlexiGo Inc