MILK OF MAGNESIA CHERRY- magnesium hydroxide liquid Power Health, Products, Inc.

gh mom chry

Active ingredient (in each 15 mL tablespoonful)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- usually produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are

taking a prescription drug.

This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- follow each dose with a full glass (8 oz) of liquid

age	dose
adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information

- each 15 mL tablespoonful contains: magnesium 500 mg, sodium 6 mg
- store at room temperature and avoid freezing
- close cap tightly after use

Inactive ingredients

carboxymethylcellulose sodium, citric acid, D and C red no. 28, flavor, glycerin, microcrystalline cellulose, purified water, sodium citrate, sodium hypochlorite, sucrose, xanthan gum.

package Label



Drug Facts

TAMPER-EVIDENT: Do not use if imprinted neckband is missing or broken.

Active ingredient (in each 15 mL) Magnesium hydroxide 1200 mg

Purpose Saline laxative

relieves occasional constipation (irregularity)

. usually produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use If you have • kidney disease

- · a magnesium-restricted diet · stomach pain, nausea, or vomiting
- · a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- · you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

shake well before use

- . do not exceed the maximum recommended daily dose in a 24 hour period
- · dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- follow each dosage with a full glass (8 oz) of liquid

age	dose
adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor



Other information • each 15 mL contains:

- magnesium 500 mg, sodium 6 mg
 store at room temperature and avoid freezing
- · close cap tightly after use

Inactive ingredients carboxymethylcellulose sodium, citric acid. D & C red no. 28, flavor, glycerin, microcrystalline cellulose, purified water, sodium citrate, sodium hypochlorite, sucrose, xanthan gum

*This product is not manufactured or distributed by the owner of the registered trademark Phillips ® Milk of Magnesia. REV GC0523

938-19312-5

Distributed by Power Health International Corp. 7919 20th Street Hollywood, Florida 33024 U.S.A.

MILK OF MAGNESIA CHERRY

magnesium hydroxide liquid

Product Information

HUMAN OTC DRUG NDC:85740-648 **Product Type** Item Code (Source)

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

Inactive Ingredients			
Ingredient Name	Strength		
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K6790BS311)			
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)			
D&C RED NO. 28 (UNII: 767IP0Y5NH)			
GLYCERIN (UNII: PDC6A3C0OX)			
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
WATER (UNII: 059QF0KO0R)			
SODIUM CITRATE (UNII: 1Q73Q2JULR)			
SODIUM HYPOCHLORITE (UNII: DY38VHM5OD)			
SUCROSE (UNII: C151H8M554)			
XANTHAN GUM (UNII: TTV12P4NEE)			

Product Characteristics			
Color	pink (opaque light pink)	Score	
Shape		Size	
Flavor	CHERRY (cherry)	Imprint Code	
Contains			

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2021	

Marketing Information			
Marketing Application Number or Monograph Category Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	01/01/2021	

Labeler - Power Health, Products, Inc. (005646227)

Registrant - GCP Laboratories (965480861)

Establishment			
Name	Address	ID/FEI	Business Operations
GCP LABORATORIES		965480861	manufacture(85740-648)

Revised: 5/2025 Power Health, Products, Inc.