

**FIBROMYALGIA SUPPORT- aconitum napellus, arsenicum album, belladonna, conium, gelsemium sempervirens, hypericum perforatum, kali bichromicum, lacticum acidum, phosphoricum acidum, rhus tox, uricum acidum tablet**  
**The Magni Company**

*Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.*

-----

**DRUG FACTS:**

**ACTIVE INGREDIENTS:**

Aconitum Napellus 30C, Arsenicum Album 30C, Belladonna 30C, Conium 30C, Gelsemium Sempervirens 30C, Hypericum Perforatum 3X, 30C, Kali Bichromicum 30C, Lacticum Acidum 30C, Phosphoricum Acidum 30C, Rhus Tox 30C, Uricum Acidum 30C.

**PURPOSE:**

Aconitum Napellus – For pain in joints and tendons, Arsenicum Album – For pain of the extremities, Belladonna – For pain and sensitivity, Conium – For muscle aches and pains, Gelsemium Sempervirens – For muscular weakness, Hypericum Perforatum – Reduces pain sensitivity, Kali Bichromicum – For pain and stiffness in joints, Lacticum Acidum – For arthritic pains, Phosphoricum Acidum – For weakness and pain, Rhus Tox – For pain and stiffness in joints and tendons, Uricum Acidum – For pain of the small joints

**USES:**

Temporarily relieves widespread aches, pains, and stiffness with fatigue

**WARNINGS:**

- **If pregnant or breastfeeding**, ask a health professional before use
- **Keep out of reach of children** In case of overdose, get medical help or contact a Poison Control Center right away
- If symptoms persist for more than 7 days or worsen, consult your physician

**KEEP OUT OF REACH OF CHILDREN:**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away

**DIRECTIONS:**

- Adults and children 12 years and above, dissolve 2 tablets under tongue 2 to 3 times daily or as directed by a physician
- Children under 12 years, consult a physician
- Take at least 10 minutes before or after eating or drinking

**OTHER INFORMATION:**

- Store tightly closed in a cool, dry place
- Tamper Sealed: Sealed for your protection. Do not use if seal is broken or missing
- The letters "HPUS" indicate that the components in this product are officially monographed in the Homeopathic Pharmacopeia of the United States
- These statements are supported by the traditional homeopathic principles

**INACTIVE INGREDIENTS:**

Lactose, Magnesium Stearate, Microcrystalline Cellulose

**QUESTIONS:**

Dist. by: The Magni Group, Inc. McKinney, TX 75072 USA

[www.magilife.com](http://www.magilife.com)

1-800-645-9199

**PACKAGE LABEL DISPLAY:**

**MAGNILIFE** Homeopathic  
**FIBROMYALGIA** NDC 43689-0058-1  
**SUPPORT**

**PAIN & FATIGUE RELIEF**

For symptoms of


Fibromyalgia

- ✓ Muscle Aches
- ✓ Arthritis
- ✓ Joint Pain
- ✓ Stiffness
- ✓ Fatigue

125 Dissolving

Tablets

<b>Drug Facts</b> (continued)	<div style="text-align: right;">Homeopathic NDC 43689-0058-1</div> <h2 style="text-align: center;">MAGNILIFE® FIBROMYALGIA SUPPORT</h2> <h3 style="text-align: center;">PAIN &amp; FATIGUE RELIEF</h3> <p style="text-align: center;"><i>For Symptoms of Fibromyalgia</i></p> <ul style="list-style-type: none"> <li>✓ Muscle Aches</li> <li>✓ Arthritis</li> <li>✓ Joint Pain</li> <li>✓ Stiffness</li> <li>✓ Fatigue</li> </ul>  <div style="text-align: center; font-weight: bold; font-size: 1.2em;">125 Dissolving Tablets</div>	<b>Drug Facts</b> <table border="0" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left; border-bottom: 1px solid black;">Active ingredients (in each tablet)</th> <th style="text-align: left; border-bottom: 1px solid black;">Purpose</th> </tr> </thead> <tbody> <tr> <td>Aconitum napellus 30C HPUS</td> <td>For pain in joints and tendons</td> </tr> <tr> <td>Arsenicum album 30C HPUS</td> <td>For pain of the extremities</td> </tr> <tr> <td>Belladonna 30C HPUS</td> <td>For pain and sensitivity</td> </tr> <tr> <td>Conium 30C HPUS</td> <td>For muscle aches and pains</td> </tr> <tr> <td>Gelsemium sempervirens 80C HPUS</td> <td>For muscular weakness</td> </tr> <tr> <td>Hypericum perforatum 3X, 30C HPUS</td> <td>Reduces pain sensitivity</td> </tr> <tr> <td>Kali bichromicum 30C HPUS</td> <td>For pain and stiffness in joints</td> </tr> <tr> <td>Lacticum acidum 90C HPUS</td> <td>For arthritic pains</td> </tr> <tr> <td>Phosphoricum acidum 30C HPUS</td> <td>For weakness and pain</td> </tr> <tr> <td>Rhus tox 30C HPUS</td> <td>For pain and stiffness in joints and tendons</td> </tr> <tr> <td>Uricum acidum 30C HPUS</td> <td>For pain of the small joints</td> </tr> </tbody> </table> <p><b>Uses</b> Temporarily relieves widespread aches, pains, and stiffness with fatigue →</p>	Active ingredients (in each tablet)	Purpose	Aconitum napellus 30C HPUS	For pain in joints and tendons	Arsenicum album 30C HPUS	For pain of the extremities	Belladonna 30C HPUS	For pain and sensitivity	Conium 30C HPUS	For muscle aches and pains	Gelsemium sempervirens 80C HPUS	For muscular weakness	Hypericum perforatum 3X, 30C HPUS	Reduces pain sensitivity	Kali bichromicum 30C HPUS	For pain and stiffness in joints	Lacticum acidum 90C HPUS	For arthritic pains	Phosphoricum acidum 30C HPUS	For weakness and pain	Rhus tox 30C HPUS	For pain and stiffness in joints and tendons	Uricum acidum 30C HPUS	For pain of the small joints
Active ingredients (in each tablet)		Purpose																								
Aconitum napellus 30C HPUS		For pain in joints and tendons																								
Arsenicum album 30C HPUS		For pain of the extremities																								
Belladonna 30C HPUS	For pain and sensitivity																									
Conium 30C HPUS	For muscle aches and pains																									
Gelsemium sempervirens 80C HPUS	For muscular weakness																									
Hypericum perforatum 3X, 30C HPUS	Reduces pain sensitivity																									
Kali bichromicum 30C HPUS	For pain and stiffness in joints																									
Lacticum acidum 90C HPUS	For arthritic pains																									
Phosphoricum acidum 30C HPUS	For weakness and pain																									
Rhus tox 30C HPUS	For pain and stiffness in joints and tendons																									
Uricum acidum 30C HPUS	For pain of the small joints																									
<b>Warnings</b> • If pregnant or breastfeeding, ask a health professional before use • Keep out of reach of children • In case of overdose, get medical help or contact a Poison Control Center right away • If symptoms persist for more than 7 days or worsen, consult your physician																										
<b>Directions</b> • Adults and children 12 years and above, dissolve 2 tablets under tongue 2 to 3 times daily or as directed by a physician • Children under 12 years, consult a physician • Take at least 10 minutes before or after eating or drinking																										
<b>Other information</b> • Store tightly closed in a dry place • Tamper Sealed: Sealed for your protection. Do not use if seal is broken or missing • The letters "HPUS" indicate that the components in this product are officially monographed in the Homeopathic Pharmacopoeia of the United States • These statements are supported by the traditional homeopathic principles																										
<b>Inactive ingredients</b> Lactose, Magnesium Stearate, Microcrystalline Cellulose																										
Dist. by: The Magni Group, Inc. McKinney, TX 75072 USA <a href="http://www.magnilife.com">www.magnilife.com</a> 1-800-645-9199																										

<div style="text-align: center;"> <p>TRY</p> <h2 style="margin: 0;">MAGNILIFE® FIBROMYALGIA SUPPORT</h2> </div> <div style="border: 1px solid black; padding: 5px;"> <p><b>Drug Facts</b></p> <p><b>Active ingredients (in each tablet)</b> . . . . . Purpose</p> <p>Aconitum napellus 30C HPUS . . . . . For pain in joints and tendons</p> <p>Arsenicum album 30C HPUS . . . . . For pain of the extremities</p> <p>Belladonna 30C HPUS . . . . . For pain and sensitivity</p> <p>Conium 30C HPUS . . . . . For muscle aches and pains</p> <p>Gelsemium sempervirens 80C HPUS . . . . . For muscular weakness</p> <p>Hypericum perforatum 3X, 30C HPUS . . . . . Reduces pain sensitivity</p> <p>Kali bichromicum 30C HPUS . . . . . For pain and stiffness in joints</p> <p>Lacticum acidum 90C HPUS . . . . . For arthritic pains</p> <p>Phosphoricum acidum 30C HPUS . . . . . For weakness and pain</p> <p>Rhus tox 30C HPUS . . . . . For pain and stiffness in joints and tendons</p> <p>Uricum acidum 30C HPUS . . . . . For pain of the small joints</p> <p><b>Uses</b> Temporarily relieves widespread aches, pains, and stiffness with fatigue.</p> <p><b>Warnings</b></p> <p>• If pregnant or breastfeeding, ask a health professional before use • Keep out of reach of children • In case of overdose, get medical help or contact a Poison Control Center right away • If symptoms persist for more than 7 days or worsen, consult your physician</p> <p><b>Directions</b> • Adults and children 12 years and above, dissolve 2 tablets under tongue 2 to 3 times daily or as directed by a physician • Children under 12 years, consult a physician • Take at least 10 minutes before or after eating or drinking</p> <p><b>Other information</b> • Store tightly closed in a dry place • Tamper Sealed: Sealed for your protection. Do not use if seal is broken or missing • The letters "HPUS" indicate that the components in this product are officially monographed in the Homeopathic Pharmacopoeia of the United States • These statements are supported by the traditional homeopathic principles</p> <p><b>Inactive ingredients</b> Lactose, Magnesium Stearate, Microcrystalline Cellulose</p> <p>Dist. by: The Magni Group, Inc. McKinney, TX 75072 USA  <a href="http://www.magnilife.com">www.magnilife.com</a> 1-800-645-9199</p> </div>	<div style="text-align: right;">Homeopathic NDC 43689-0058-2</div> <h2 style="text-align: center;">MAGNILIFE® FIBROMYALGIA SUPPORT</h2> <h3 style="text-align: center;">PAIN &amp; FATIGUE RELIEF</h3> <p style="text-align: center;"><i>For Symptoms of Fibromyalgia</i></p> <ul style="list-style-type: none"> <li>✓ Muscle Aches</li> <li>✓ Arthritis</li> <li>✓ Joint Pain</li> <li>✓ Stiffness</li> <li>✓ Fatigue</li> </ul> <p>Temporarily relieves muscle aches, joint pain, arthritic pains, and stiffness usually accompanied by extreme fatigue</p> <p>Provides natural pain relief</p>  <div style="text-align: center; font-weight: bold; font-size: 1.2em;">125 Dissolving Tablets</div> <p style="font-size: 0.8em;">Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective. Not intended to treat or cure Fibromyalgia</p>	<h2 style="text-align: center;">MAGNILIFE® FIBROMYALGIA SUPPORT</h2> <h3 style="text-align: center;">PAIN &amp; FATIGUE RELIEF</h3> <p style="text-align: center;"><i>For Symptoms of Fibromyalgia</i></p> <ul style="list-style-type: none"> <li>✓ Muscle Aches</li> <li>✓ Arthritis</li> <li>✓ Joint Pain</li> <li>✓ Stiffness</li> <li>✓ Fatigue</li> </ul> <p>Temporarily relieves muscle aches, joint pain, arthritic pains, and stiffness usually accompanied by extreme fatigue</p> <p>Provides natural pain relief</p>  <div style="text-align: center; font-weight: bold; font-size: 1.2em;">125 Dissolving Tablets</div>	<h2 style="text-align: center;">MAGNILIFE® FIBROMYALGIA SUPPORT</h2> <p style="text-align: center;">No known contraindications</p> <p style="text-align: center;">No harmful side effects</p> <p style="text-align: center;">Tablets dissolve under the tongue</p> 																																																				
				<p><b>Product Information</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%;"><b>Product Type</b></td> <td style="width: 30%;">HUMAN OTC DRUG</td> <td style="width: 20%;"><b>Item Code (Source)</b></td> <td style="width: 20%;">NDC:43689-0058</td> </tr> <tr> <td><b>Route of Administration</b></td> <td>ORAL</td> <td></td> <td></td> </tr> <tr> <td colspan="4"><b>Active Ingredient/Active Moiety</b></td> </tr> </table>	<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058	<b>Route of Administration</b>	ORAL			<b>Active Ingredient/Active Moiety</b>				<p><b>Product Information</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%;"><b>Product Type</b></td> <td style="width: 30%;">HUMAN OTC DRUG</td> <td style="width: 20%;"><b>Item Code (Source)</b></td> <td style="width: 20%;">NDC:43689-0058</td> </tr> <tr> <td><b>Route of Administration</b></td> <td>ORAL</td> <td></td> <td></td> </tr> <tr> <td colspan="4"><b>Active Ingredient/Active Moiety</b></td> </tr> </table>	<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058	<b>Route of Administration</b>	ORAL			<b>Active Ingredient/Active Moiety</b>				<p><b>Product Information</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%;"><b>Product Type</b></td> <td style="width: 30%;">HUMAN OTC DRUG</td> <td style="width: 20%;"><b>Item Code (Source)</b></td> <td style="width: 20%;">NDC:43689-0058</td> </tr> <tr> <td><b>Route of Administration</b></td> <td>ORAL</td> <td></td> <td></td> </tr> <tr> <td colspan="4"><b>Active Ingredient/Active Moiety</b></td> </tr> </table>	<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058	<b>Route of Administration</b>	ORAL			<b>Active Ingredient/Active Moiety</b>				<p><b>Product Information</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%;"><b>Product Type</b></td> <td style="width: 30%;">HUMAN OTC DRUG</td> <td style="width: 20%;"><b>Item Code (Source)</b></td> <td style="width: 20%;">NDC:43689-0058</td> </tr> <tr> <td><b>Route of Administration</b></td> <td>ORAL</td> <td></td> <td></td> </tr> <tr> <td colspan="4"><b>Active Ingredient/Active Moiety</b></td> </tr> </table>	<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058	<b>Route of Administration</b>	ORAL			<b>Active Ingredient/Active Moiety</b>			
					<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058																																															
					<b>Route of Administration</b>	ORAL																																																	
<b>Active Ingredient/Active Moiety</b>																																																							
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058																																																				
<b>Route of Administration</b>	ORAL																																																						
<b>Active Ingredient/Active Moiety</b>																																																							
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058																																																				
<b>Route of Administration</b>	ORAL																																																						
<b>Active Ingredient/Active Moiety</b>																																																							
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058																																																				
<b>Route of Administration</b>	ORAL																																																						
<b>Active Ingredient/Active Moiety</b>																																																							
<p><b>Product Information</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%;"><b>Product Type</b></td> <td style="width: 30%;">HUMAN OTC DRUG</td> <td style="width: 20%;"><b>Item Code (Source)</b></td> <td style="width: 20%;">NDC:43689-0058</td> </tr> <tr> <td><b>Route of Administration</b></td> <td>ORAL</td> <td></td> <td></td> </tr> <tr> <td colspan="4"><b>Active Ingredient/Active Moiety</b></td> </tr> </table>	<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058	<b>Route of Administration</b>	ORAL			<b>Active Ingredient/Active Moiety</b>				<p><b>Product Information</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%;"><b>Product Type</b></td> <td style="width: 30%;">HUMAN OTC DRUG</td> <td style="width: 20%;"><b>Item Code (Source)</b></td> <td style="width: 20%;">NDC:43689-0058</td> </tr> <tr> <td><b>Route of Administration</b></td> <td>ORAL</td> <td></td> <td></td> </tr> <tr> <td colspan="4"><b>Active Ingredient/Active Moiety</b></td> </tr> </table>	<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058	<b>Route of Administration</b>	ORAL			<b>Active Ingredient/Active Moiety</b>				<p><b>Product Information</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%;"><b>Product Type</b></td> <td style="width: 30%;">HUMAN OTC DRUG</td> <td style="width: 20%;"><b>Item Code (Source)</b></td> <td style="width: 20%;">NDC:43689-0058</td> </tr> <tr> <td><b>Route of Administration</b></td> <td>ORAL</td> <td></td> <td></td> </tr> <tr> <td colspan="4"><b>Active Ingredient/Active Moiety</b></td> </tr> </table>	<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058	<b>Route of Administration</b>	ORAL			<b>Active Ingredient/Active Moiety</b>				<p><b>Product Information</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%;"><b>Product Type</b></td> <td style="width: 30%;">HUMAN OTC DRUG</td> <td style="width: 20%;"><b>Item Code (Source)</b></td> <td style="width: 20%;">NDC:43689-0058</td> </tr> <tr> <td><b>Route of Administration</b></td> <td>ORAL</td> <td></td> <td></td> </tr> <tr> <td colspan="4"><b>Active Ingredient/Active Moiety</b></td> </tr> </table>	<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058	<b>Route of Administration</b>	ORAL			<b>Active Ingredient/Active Moiety</b>							
	<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058																																																			
<b>Route of Administration</b>	ORAL																																																						
<b>Active Ingredient/Active Moiety</b>																																																							
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058																																																				
<b>Route of Administration</b>	ORAL																																																						
<b>Active Ingredient/Active Moiety</b>																																																							
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058																																																				
<b>Route of Administration</b>	ORAL																																																						
<b>Active Ingredient/Active Moiety</b>																																																							
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058																																																				
<b>Route of Administration</b>	ORAL																																																						
<b>Active Ingredient/Active Moiety</b>																																																							
<p><b>Product Information</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%;"><b>Product Type</b></td> <td style="width: 30%;">HUMAN OTC DRUG</td> <td style="width: 20%;"><b>Item Code (Source)</b></td> <td style="width: 20%;">NDC:43689-0058</td> </tr> <tr> <td><b>Route of Administration</b></td> <td>ORAL</td> <td></td> <td></td> </tr> <tr> <td colspan="4"><b>Active Ingredient/Active Moiety</b></td> </tr> </table>	<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058	<b>Route of Administration</b>	ORAL			<b>Active Ingredient/Active Moiety</b>				<p><b>Product Information</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%;"><b>Product Type</b></td> <td style="width: 30%;">HUMAN OTC DRUG</td> <td style="width: 20%;"><b>Item Code (Source)</b></td> <td style="width: 20%;">NDC:43689-0058</td> </tr> <tr> <td><b>Route of Administration</b></td> <td>ORAL</td> <td></td> <td></td> </tr> <tr> <td colspan="4"><b>Active Ingredient/Active Moiety</b></td> </tr> </table>	<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058	<b>Route of Administration</b>	ORAL			<b>Active Ingredient/Active Moiety</b>				<p><b>Product Information</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%;"><b>Product Type</b></td> <td style="width: 30%;">HUMAN OTC DRUG</td> <td style="width: 20%;"><b>Item Code (Source)</b></td> <td style="width: 20%;">NDC:43689-0058</td> </tr> <tr> <td><b>Route of Administration</b></td> <td>ORAL</td> <td></td> <td></td> </tr> <tr> <td colspan="4"><b>Active Ingredient/Active Moiety</b></td> </tr> </table>	<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058	<b>Route of Administration</b>	ORAL			<b>Active Ingredient/Active Moiety</b>				<p><b>Product Information</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%;"><b>Product Type</b></td> <td style="width: 30%;">HUMAN OTC DRUG</td> <td style="width: 20%;"><b>Item Code (Source)</b></td> <td style="width: 20%;">NDC:43689-0058</td> </tr> <tr> <td><b>Route of Administration</b></td> <td>ORAL</td> <td></td> <td></td> </tr> <tr> <td colspan="4"><b>Active Ingredient/Active Moiety</b></td> </tr> </table>	<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058	<b>Route of Administration</b>	ORAL			<b>Active Ingredient/Active Moiety</b>							
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058																																																				
<b>Route of Administration</b>	ORAL																																																						
<b>Active Ingredient/Active Moiety</b>																																																							
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058																																																				
<b>Route of Administration</b>	ORAL																																																						
<b>Active Ingredient/Active Moiety</b>																																																							
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058																																																				
<b>Route of Administration</b>	ORAL																																																						
<b>Active Ingredient/Active Moiety</b>																																																							
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058																																																				
<b>Route of Administration</b>	ORAL																																																						
<b>Active Ingredient/Active Moiety</b>																																																							

## FIBROMYALGIA SUPPORT

aconitum napellus, arsenicum album, belladonna, conium, gelsemium sempervirens, hypericum perforatum, kali bichromicum, lacticum acidum, phosphoricum acidum, rhus tox, uricum acidum tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:43689-0058
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACONITUM NAPELLUS WHOLE</b> (UNII: U0NQ8555JD) (ACONITUM NAPELLUS - UNII:U0NQ8555JD)	ACONITUM NAPELLUS WHOLE	30 [hp_C]
<b>ARSENIC TRIOXIDE</b> (UNII: S7V92P67HO) (ARSENIC CATION (3+) - UNII:C96613F5AV)	ARSENIC TRIOXIDE	30 [hp_C]
<b>ATROPA BELLADONNA</b> (UNII: WQZ3G9PF0H) (ATROPA BELLADONNA - UNII:WQZ3G9PF0H)	ATROPA BELLADONNA	30 [hp_C]
<b>CONIINE</b> (UNII: C479P32L2D) (CONIINE - UNII:C479P32L2D)	CONIINE	30 [hp_C]
<b>GELSEMIUM SEMPERVIRENS ROOT</b> (UNII: 639KR60Q1Q) (GELSEMIUM SEMPERVIRENS ROOT - UNII:639KR60Q1Q)	GELSEMIUM SEMPERVIRENS ROOT	30 [hp_C]
<b>HYPERICUM PERFORATUM WHOLE</b> (UNII: XK4IUX8MNB) (HYPERICUM PERFORATUM - UNII:XK4IUX8MNB)	HYPERICUM PERFORATUM WHOLE	3 [hp_X]
<b>POTASSIUM DICHROMATE</b> (UNII: T4423S18FM) (DICHROMATE ION - UNII:9LKY4BFN2V)	POTASSIUM DICHROMATE	30 [hp_C]
<b>LACTIC ACID, DL-</b> (UNII: 3B8D35Y7S4) (LACTIC ACID, DL- - UNII:3B8D35Y7S4)	LACTIC ACID, DL-	30 [hp_C]
<b>PHOSPHORIC ACID</b> (UNII: E4GA8884NN) (PHOSPHORIC ACID - UNII:E4GA8884NN)	PHOSPHORIC ACID	30 [hp_C]
<b>TOXICODENDRON PUBESCENS LEAF</b> (UNII: 6IO182RP7A) (TOXICODENDRON PUBESCENS LEAF - UNII:6IO182RP7A)	TOXICODENDRON PUBESCENS LEAF	30 [hp_C]
<b>URIC ACID</b> (UNII: 268B43MJ25) (URIC ACID - UNII:268B43MJ25)	URIC ACID	30 [hp_C]

### Inactive Ingredients

Ingredient Name	Strength
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND (Round Convex)	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	ML
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43689-0058-2	1 in 1 CARTON	01/25/2023	
1	NDC:43689-0058-1	125 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved homeopathic		01/25/2023	

**Labeler** - The Magni Company (113501902)

**Registrant** - Apotheca Company (844330915)

**Establishment**

Name	Address	ID/FEI	Business Operations
Apotheca Company		844330915	manufacture(43689-0058) , api manufacture(43689-0058) , label(43689-0058) , pack(43689-0058)

Revised: 10/2025

The Magni Company