

NIGHTTIME CHERRY COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate liquid
BETTER LIVING BRANDS LLC.

750L-Albertsons-Acetaminophen 650mg , Dextromethorphan Hydrobromide 30mg, and Doxylamine Succinate 12.5mg

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash,

nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL contains:** sodium 52 mg
- store at no greater than 25°C and do not refrigerate

Inactive ingredients

alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum

Questions or comments? Call **1-877-290-4008**

COMPARE TO

Vicks® NyQuil™ Cold & Flu
Cherry Liquid Active Ingredients*

NDC 21130-907-01

Signature
SELECT

**NIGHTTIME
COLD & FLU
RELIEF**

**FOR AGES 12
& OVER**

**CHERRY
FLAVOR**

ACETAMINOPHEN 650 mg

- Pain Reliever/Fever Reducer

DEXTROMETHORPHAN HBr 30 mg

- Cough Suppressant

DOXYLAMINE SUCCINATE 12.5 mg

- Antihistamine

• Relieves headaches, fever, sore
throat, minor aches & pains

• Sneezing, runny nose, cough

• Nighttime relief



ALCOHOL 10%

12 FL OZ (354 mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND SEAL AROUND THE NECK IS BROKEN OR MISSING

← PEEL BACK FOR DRUG FACTS

Drug Facts

Active Ingredients (in each 30 mL)

Acetaminophen 650 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 30 mg.....Cough suppressant
Doxylamine succinate 12.5 mg.....Antihistamine

Purpose

Uses temporarily relieves common cold/flu symptoms:
■ cough due to minor throat and bronchial irritation
■ sore throat
■ headache
■ minor aches and pains
■ fever
■ runny nose and sneezing

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
■ more than 4 doses in 24 hours, which is the maximum daily amount
■ for this product

DISTRIBUTED BY:
BETTER LIVING BRANDS LLC
P.O. BOX 99
PLEASANTON, CA 94566-0009
1-888-723-3029

750L 0525 Made in India
RD 25128 S6850

PLASTIC BOTTLE CUP

NOT FOR RESALE

LOT:
EXP.:
Print/Varnish Omit Area

smartlabel™ Scan here for more information

PARENTS: Learn about teen medicine abuse
www.StopMedicineAbuse.org

BACK

Drug Facts (continued)

■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions.
Symptoms may include:
■ skin reddening
■ hives
■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
■ liver disease
■ glaucoma
■ cough that occurs with too much phlegm (mucus)
■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are
■ taking sedatives or tranquilizers
■ taking the blood thinning drug warfarin

When using this product
■ excitability may occur, especially in children
■ marked drowsiness may occur
■ avoid alcoholic drinks
■ be careful when driving a motor vehicle or operating machinery

BACK PANEL 1

Drug Facts (continued)

■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if
■ pain or cough gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ take only as directed
■ only use the dose cup provided
■ do not exceed 4 doses per 24 hrs
adults & children 12 yrs & over 30 mL every 6 hrs
children 4 to under 12 yrs ask a doctor
children under 4 yrs do not use

Other information
■ each 30 mL contains: sodium 52 mg
■ store at no greater than 25°C and do not refrigerate

Inactive ingredients alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum

Questions or comments? Call 1-877-290-4008

*This product is not manufactured or distributed by Procter & Gamble Company, owner of the registered trademarks Vicks® NyQuil® Cold & Flu Cherry Liquid.

BACK PANEL 2

TWIN PACK

COMPARE TO Vicks® NyQuil® Cold & Flu Cherry Liquid Active Ingredients*
NDC 21130-907-02

Signature SELECT

NIGHTTIME COLD & FLU RELIEF

FOR AGES 12 & OVER **CHERRY FLAVOR**

ACETAMINOPHEN 650 mg
- Pain Reliever/Fever Reducer
DEXTROMETHORPHAN HBr 30 mg
- Cough Suppressant
DOXYLAMINE SUCCINATE 12.5 mg
- Antihistamine

Relieves headaches, fever, sore throat, minor aches & pains
Sneezing, runny nose, cough
Nighttime relief

ALCOHOL 10%

2 BOTTLES - 12 FL OZ (354 mL) EACH 24 FL OZ (708 mL) TOTAL

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 30 mg.....Cough suppressant
Doxylamine succinate 12.5 mg.....Antihistamine

Purpose

Uses temporarily relieves common cold/flu symptoms:
■ cough due to minor throat and bronchial irritation
■ sore throat
■ headache
■ minor aches and pains
■ fever
■ runny nose and sneezing

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
■ more than 4 doses in 24 hours, which is the maximum daily amount for this product
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions.
Symptoms may include: ■ skin reddening ■ hives ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
■ liver disease
■ glaucoma
■ cough that occurs with too much phlegm (mucus)
■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are
■ taking sedatives or tranquilizers
■ taking the blood thinning drug warfarin

When using this product
■ excitability may occur, especially in children
■ marked drowsiness may occur
■ avoid alcoholic drinks
■ be careful when driving a motor vehicle or operating machinery
■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if
■ pain or cough gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for

Drug Facts (continued)

adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ take only as directed
■ only use the dose cup provided
■ do not exceed 4 doses per 24 hrs
adults & children 12 yrs & over 30 mL every 6 hrs
children 4 to under 12 yrs ask a doctor
children under 4 yrs do not use

Other information
■ each 30 mL contains: sodium 52 mg
■ store at no greater than 25°C and do not refrigerate

Inactive ingredients alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum

Questions or comments? Call 1-877-290-4008

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND SEAL AROUND THE NECK OF BOTTLE IS BROKEN OR MISSING

*This product is not manufactured or distributed by Procter & Gamble Company, owner of the registered trademarks Vicks® NyQuil® Cold & Flu Cherry Liquid.

smartlabel™ Scan here for more information

DISTRIBUTED BY:
BETTER LIVING BRANDS LLC
P.O. BOX 99
PLEASANTON, CA 94566-0009
1-888-723-3029

Made in India 750L 0425

RD25101 S6850

3 211307879415

PARENTS: Learn about teen medicine abuse
www.StopMedicineAbuse.org

LOT:
EXP.:
Print/Varnish Omit Area

NIGHTTIME CHERRY COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-907
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL

DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM SACCHARIN (UNII: SB8ZUX40TY)	
GLYCERIN (UNII: PDC6A3C0OX)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
ALCOHOL (UNII: 3K9958V90M)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
XANTHAN GUM (UNII: TTV12P4NEE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
WATER (UNII: 059QF0KO0R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-907-02	2 in 1 PACKAGE	06/01/2025	
1	NDC:21130-907-01	354 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/01/2025	

Labeler - BETTER LIVING BRANDS LLC. (009137209)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment			
Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		677604129	manufacture(21130-907)

Revised: 6/2025

BETTER LIVING BRANDS LLC.