

**NIGHTTIME COLD AND FLU- acetaminophen, doxylamine succinate and dextromethorphan hbr liquid**  
**BI-MART**

-----  
**NIGHTTIME COLD & FLU**

***Drug Facts***

***Active ingredients (in each 30 mL dose cup)***

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine Succinate 12.5 mg

***Purpose***

Pain Reliever/Fever Reducer

Cough Suppressant

Antihistamine

***Uses***

temporarily relieves common cold/flu symptoms:

- sore throat
- minor aches and pains
- runny nose and sneezing
- headache
- fever
- cough due to minor throat and bronchial irritation.

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if:

- adults and children over 12 years of age take more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occur with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to an enlarged prostate gland.

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin.

### **When using this product**

- do not use more than directed (see overdose warning)
- avoid alcoholic drinks
- excitability may occur, especially in children
- marked drowsiness may occur
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness.

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of the reach of children.**

**Overdose warning:**

Taking more than the recommended dose (overdose) could cause serious health problems, including liver damage. In case of accidental overdose, seek professional assistance or contact a Poison Control Center (1-800-222-1222) immediately. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

- take only as directed (see overdose warning)
- measure only with dosing cup provided and keep dosage cup with product
- mL=milliliter
- do not exceed 4 doses per 24 hours

<b>Age</b>	<b>Dose</b>
adults and children 12 years and over	30 mL every 6 hours
children under 12 years	do not use

***Other Information***

- **Each 30 mL dose cup contains:**sodium 10 mg
- store at room temperature 15-30°C (59-86°F) and do not refrigerate

***Inactive ingredients***

citric acid, D&C Yellow #10, FD&C Green #3, FD&C Yellow #6, flavor, glycerin, menthol, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum.

***Questions or comments?***

1-844-912-4012

TAMPER-EVIDENT FEATURE: DO NOT USE IF PRINTED  
NECK WRAP IS BROKEN OR MISSING.

  
0 71357 91245 12

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

BI-MART UNCONDITIONAL MONEY BACK GUARANTEE  
If you are not completely satisfied with your BI-MART  
QUALITY PRODUCT, regardless of the reason, return the  
unused portion and your entire purchase price will be  
cheerfully refunded.

\*This product is not manufactured  
or distributed by The Procter &  
Gamble Company, distributor of  
Vicks® NyQuil® Cough & Flu.

Distributed by:  
BI-MART  
Eugene, OR 97402

525-12-BM 04/25  
Rev. 1

no print / no varnish area  
lot no. & exp. date

**BI-MART**  
NDC 37835-525-12

Compare to the active  
ingredients in Vicks® NyQuil®  
Cold & Flu\*

**NIGHTTIME**  
**COLD & FLU**  
**Acetaminophen** (Pain reliever/fever reducer)  
**Doxylamine Succinate** (Antihistamine)  
**Dextromethorphan HBr** (Cough suppressant)  
  
Headache, Fever, Sore Throat,  
Minor Aches & Pains,  
Sneezing, Runny Nose,  
Cough  
  
Ages 12 Years and Over  
  
12 FL OZ (355 mL)

**Drug Facts**  
**Active ingredients** (in each 30 mL dose cup)  
**Acetaminophen** 650 mg **Pain Reliever/Fever Reducer**  
**Dextromethorphan HBr** 30 mg **Cough Suppressant**  
**Doxylamine Succinate** 12.5 mg **Antihistamine**  
**Uses** temporarily relieves common cold/flu symptoms: • sore throat  
• minor aches and pains • runny nose and sneezing • headache • fever  
• cough due to minor throat and bronchial irritation  
**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage  
may occur if:  
• adults and children over 12 years of age take more than 4 doses (30 mL  
each) in 24 hours, which is the maximum daily amount for this product  
• taken with other drugs containing acetaminophen  
• adult has 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms  
may include: • skin redness • blisters • rash. If a skin reaction occurs,  
stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, is  
accompanied or followed by fever, headache, rash, nausea, or vomiting,  
consult a doctor promptly.  
**Do not use** • with any other drug containing acetaminophen (prescription or  
nonprescription). If you are not sure whether a drug contains  
acetaminophen, ask a doctor or pharmacist. • If you are now taking a  
prescription monoamine oxidase inhibitor (MAOI) (certain drugs for  
depression, psychiatric, or emotional conditions, or Parkinson's disease), or  
for 2 weeks after stopping the MAOI drug. If you do not know if your  
prescription drug contains an MAOI, ask a doctor or pharmacist before taking  
this product. • If you have ever had an allergic reaction to this product or any  
of its ingredients.  
**Ask a doctor before use if you have** • liver disease • glaucoma • cough  
that occurs with too much phlegm (mucus) • a breathing problem or chronic  
cough that lasts or as occur with smoking, asthma, chronic bronchitis, or  
emphysema • trouble urinating due to an enlarged prostate gland  
  
→ DRUG FACTS CONTINUED ON THE BACK →  
PEEL BACK →

**Drug Facts (continued)**  
**Ask a doctor or pharmacist before use if you are** • taking sedatives or  
tranquilizers • taking the blood thinning drug warfarin  
**When using this product** • do not use more than directed (see overdose  
warning) • avoid alcoholic drinks • excitability may occur, especially in children  
• marked drowsiness may occur • be careful when driving a motor vehicle or  
operating machinery • alcohol, sedatives, and tranquilizers may increase  
drowsiness.  
**Stop use and ask a doctor if** • pain or cough gets worse or lasts more than 7  
days • fever gets worse or lasts more than 3 days • redness or swelling is  
present • new symptoms occur • cough comes back or occurs with rash or  
headache that lasts. These could be signs of a serious condition.  
**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of the reach of children.**  
**Overdose warning:** Taking more than the recommended dose (overdose) could  
cause serious health problems, including liver damage. In case of accidental  
overdose, seek professional assistance or contact a Poison Control Center  
(1-800-222-1222) immediately. Quick medical attention is critical for adults as  
well as for children even if you do not notice any signs or symptoms.  
**Directions** • take only as directed (see overdose warning)  
• measure only with dose cup provided and keep dose cup with product  
• **ml = mL** • do not exceed 4 doses per 24 hours  

Age	Dose
adults and children 12 years and over	30 mL every 6 hours
children under 12 years	do not use

**Other information** • Each 30 mL dose cup contains: sodium 10 mg  
• store at room temperature 15–30°C (59–86°F) and do not refrigerate  
**Inactive ingredients** citric acid, D&C Yellow #10, FD&C Green #3, FD&C  
Yellow #6, flavor, glycerin, menthol, propylene glycol, purified water, sodium  
benzoate, sorbitol solution, sucralose, xanthan gum.  
**Questions or comments?** 1-844-912-4012

NIGHTTIME COLD AND FLU

acetaminophen, doxylamine succinate and dextromethorphan hbr liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37835-525
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>FD&amp;C GREEN NO. 3</b> (UNII: 3P3ONR6O1S)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>MENTHOL</b> (UNII: L7T10EIP3A)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>SORBITOL SOLUTION 70%</b> (UNII: 8KW3E207O2)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	

### Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37835-525-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2025	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2025	

**Labeler -** BI-MART (027630078)

**Registrant -** BI-MART (027630078)

### Establishment

Name	Address	ID/FEI	Business Operations
Rnv LLC		118917568	manufacture(37835-525)

Revised: 5/2025

BI-MART