

PAIN RELIEF- acetaminophen tablet
AAA Pharmaceutical, Inc.

1001-RES-2025-0602

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if the user is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has liver disease.

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 tablets every 4 to 6 hours while symptoms last▪ do not take more than 10 tablets in 24 hours▪ do not use for more than 10 days unless directed by a doctor
children 6-11 years	<ul style="list-style-type: none">▪ take 1 tablet every 4 to 6 hours while symptoms last▪ do not take more than 5 tablets in 24 hours▪ do not use for more than 5 days unless directed by a doctor
children under 6 years	<ul style="list-style-type: none">▪ ask a doctor

- store between 20-25°C (68-77°F)
- retain carton for complete product information

povidone, pregelatinized starch, sodium starch glycolate, stearic acid

1-844-705-4384

Restore U

NDC 57344-001-06

†COMPARE TO THE ACTIVE INGREDIENT IN TYLENOL® REGULAR STRENGTH TABLETS
REGULAR STRENGTH


Pain Relief

Acetaminophen

Pain Reliever / Fever Reducer

actual size

500 TABLETS - 325 MG EACH



REGULAR STRENGTH

Pain Relief

Acetaminophen

Pain Reliever / Fever Reducer

500 TABLETS - 325 MG EACH

COMPARE TO THE ACTIVE INGREDIENT IN
TYLENOL® REGULAR STRENGTH TABLETS

NDC 57344-001-06

DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)
Acetaminophen 325 mg

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- hives
- blistering
- itching
- if a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if the user is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has liver disease.

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1232). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- adults and children
 - take 2 tablets every 4 to 6 hours while symptoms last
 - do not take more than 10 tablets in 24 hours
 - do not use for more than 10 days unless directed by a doctor
- children 6-11 years
 - take 1 tablet every 4 to 6 hours while symptoms last
 - do not take more than 5 tablets in 24 hours
 - do not use for more than 5 days unless directed by a doctor
- children under 6 years
 - ask a doctor

Other information

- store between 20-25°C (68-77°F)

Inactive ingredients


povidone, pregelatinized starch, sodium starch glycolate, stearic acid

Questions or comments? 1-844-705-4384

This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol Regular Strength Tablets.

L100106RES_R0

Distributed by:
A&A Pharmaceutical, Inc.
681 Main Street
Lumberton, NJ 08048



0 99581 91239 0

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57344-301
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	325 mg

Inactive Ingredients	
Ingredient Name	Strength
POVIDONE (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics			
Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	M2A3;57344
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57344-301-06	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/02/2025	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/13/2012	

Labeler -
AAA Pharmaceutical, Inc. (181192162)