

**PAIN RELIEF- acetaminophen tablet**  
**AAA Pharmaceutical, Inc.**

-----  
**1001-RES-2025-0602**

***Drug Facts***

**Active ingredient (in each tablet)**

Acetaminophen 325 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold
  - headache
  - backache
  - minor pain of arthritis
  - toothache
  - muscular aches
  - premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert**

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if the user is allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if the user has** liver disease.

**Ask a doctor or pharmacist before use if the user is** taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"><li>▪ take 2 tablets every 4 to 6 hours while symptoms last</li><li>▪ do not take more than 10 tablets in 24 hours</li><li>▪ do not use for more than 10 days unless directed by a doctor</li></ul>
children 6-11 years	<ul style="list-style-type: none"><li>▪ take 1 tablet every 4 to 6 hours while symptoms last</li><li>▪ do not take more than 5 tablets in 24 hours</li><li>▪ do not use for more than 5 days unless directed by a doctor</li></ul>
children under 6 years	<ul style="list-style-type: none"><li>▪ ask a doctor</li></ul>

## Other information

- store between 20-25°C (68-77°F)
- retain carton for complete product information

## Inactive ingredients

povidone, pregelatinized starch, sodium starch glycolate, stearic acid

## Questions or comments?

1-844-705-4384

## PRINCIPAL DISPLAY PANEL

Restore U

NDC 57344-001-06

†COMPARE TO THE ACTIVE INGREDIENT IN TYLENOL® REGULAR STRENGTH TABLETS  
REGULAR STRENGTH

Pain Relief

Acetaminophen

Pain Reliever / Fever Reducer

actual size

500 TABLETS - 325 MG EACH

**RESTORE U**  
NDC 57344-301-06  
†COMPARE TO THE ACTIVE INGREDIENT IN TYLENOL® REGULAR STRENGTH TABLETS  
**REGULAR STRENGTH**  
**Pain Relief**  
**Acetaminophen**  
**Pain Reliever / Fever Reducer**  
actual size  
**500 TABLETS - 325 MG EACH**

**Drug Facts** **DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING**  
Acetaminophen 325 mg..... Pain reliever/fever reducer

**Purpose**  
Pain reliever/fever reducer

**Active ingredient (in each tablet)**  
Acetaminophen 325 mg.....

**Uses**  
Temporarily relieves minor aches and pains due to:  
▪ the common cold  
▪ headache  
▪ toothache  
▪ muscular aches  
▪ temporarily reduces fever

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if  
▪ adult takes more than 4,000 mg of acetaminophen in 24 hours  
▪ child takes more than 5 doses in 24 hours, which is the maximum daily amount  
▪ taken with other drugs containing acetaminophen  
▪ adult has 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
▪ skin redness  
▪ blisters  
▪ rash  
If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**  
▪ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
▪ if the user is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if the user has liver disease.**

**Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin**

**Stop use and ask a doctor if**  
▪ pain gets worse or lasts more than 10 days in adults  
▪ pain gets worse or lasts more than 5 days in children under 12 years  
▪ fever gets worse or lasts more than 3 days  
▪ new symptoms occur  
▪ redness or swelling is present  
These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
▪ **do not take more than directed (see overdose warning)**  
adults and children  
▪ take 2 tablets every 4 to 6 hours while symptoms last  
▪ do not take more than 10 tablets in 24 hours  
▪ do not use for more than 10 days unless directed by a doctor  
children 6-11 years  
▪ take 1 tablet every 4 to 6 hours while symptoms last  
▪ do not take more than 5 tablets in 24 hours  
▪ do not use for more than 5 days unless directed by a doctor  
children under 6 years  
▪ ask a doctor

**Other information**  
▪ store between 20-25°C (68-77°F)

**Inactive ingredients**  
povidone, pregelatinized starch, sodium starch glycolate, stearic acid

**Questions or comments?** 1-844-705-4384  
This product is not manufactured or distributed by the manufacturer of Tylenol® Regular Strength Tablets.  
L11001049ES\_R0  
Distributed by:  
AAA Pharmaceutical, Inc.  
681 Main Street  
Lumberton, NJ 08048

0 99581 91239 0

## PAIN RELIEF

acetaminophen tablet

### Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:57344-301

<b>Route of Administration</b>	ORAL			
<b>Active Ingredient/Active Moiety</b>				
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
	ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg	
<b>Inactive Ingredients</b>				
	<b>Ingredient Name</b>	<b>Strength</b>		
	POVIDONE (UNII: FZ989GH94E)			
	STARCH, PREGELATINIZED CORN (UNII: O8232NY3S)			
	SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)			
	STEARIC ACID (UNII: 4ELV7Z65AP)			
<b>Product Characteristics</b>				
<b>Color</b>	white	<b>Score</b>	2 pieces	
<b>Shape</b>	ROUND	<b>Size</b>	10mm	
<b>Flavor</b>		<b>Imprint Code</b>	M2A3;57344	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:57344-301-06	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/02/2025	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
OTC Monograph Drug	M013	12/13/2012		

**Labeler** - AAA Pharmaceutical, Inc. (181192162)

Revised: 7/2025

AAA Pharmaceutical, Inc.