

PAIN RELIEF- acetaminophen tablet
AAA Pharmaceutical, Inc.

1003-RES-2025-0818

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 tablets every 6 hours while symptoms last▪ do not take more than 6 tablets in 24 hours, unless directed by a doctor▪ do not use for more than 10 days unless directed by a doctor
children under 12 years	<ul style="list-style-type: none">▪ ask a doctor

Other information

- store between 20-25°C (68-77°F)
- retain carton for complete product information

Inactive ingredients

povidone, pregelatinized starch, sodium starch glycolate, stearic acid

Questions or comments?

1-844-705-4384

PRINCIPAL DISPLAY PANEL

RESTORE U

NDC 57344-003-07

†COMPARE TO THE ACTIVE INGREDIENT IN TYLENOL® EXTRA STRENGTH

For Adults

EXTRA STRENGTH

Pain Relief

Acetaminophen

Pain Reliever / Fever Reducer

actual size

500 TABLETS - 500 MG EACH

RESTORE U
NDC 57344-103-07
†COMPARE TO THE ACTIVE INGREDIENT IN TYLENOL® EXTRA STRENGTH For Adults
EXTRA STRENGTH
Pain Relief
Acetaminophen
Pain Reliever / Fever Reducer
actual size
500 TABLETS - 500 MG EACH

DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts
Acetaminophen 500 mg

Active ingredient (in each tablet)
Acetaminophen 500 mg

Purpose
Pain reliever/fever reducer

Uses
Temporarily relieves minor aches and pains due to:

- the common cold
- headache
- toothache
- temporarily reduces fever
- minor pain of arthritis
- backache
- muscular aches
- premenstrual and menstrual cramps

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.
Do not use

- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease.
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin
Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur

These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets every 6 hours while symptoms last
- do not take more than 6 tablets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20-25°C (68-77°F)

Inactive ingredients
povidone, pregelatinized starch, sodium starch glycolate, stearic acid

Questions or comments? 1-844-705-4384

Distributed by:
AAA Pharmaceutical, Inc.
681 Main Street
Lumberton, NJ 08048

This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Extra Strength

L100307RES_R0
0 195319 1240 6

PAIN RELIEF

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57344-103
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	500 mg	
Inactive Ingredients				
Ingredient Name			Strength	
POVIDONE (UNII: FZ989GH94E)				
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)				
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
Product Characteristics				
Color	white	Score	no score	
Shape	ROUND	Size	12mm	
Flavor		Imprint Code	M2A4;57344	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57344-103-07	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/02/2025	
2	NDC:57344-103-15	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/15/2025	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	06/02/2025		

Labeler - AAA Pharmaceutical, Inc. (181192162)

Revised: 12/2025

AAA Pharmaceutical, Inc.