

**SHOPRITE NI CALM- acetaminophen, dextromethorphan hbr, doxylamine succinate solution**  
**Wakefern Food Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**ShopRite Ni-Calm Drug Facts**

**Active ingredients (in each 30 mL)**

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

**Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- fever
- sore throat
- headache
- minor aches and pains
- runny nose and sneezing

**Warnings**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks

**Stop use and ask a doctor if**

- redness or swelling is present
- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

## Other information

- each 30 mL contains: sodium 39 mg
- store at 20-25°C (68-77°F)

## Inactive ingredients

alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

## Questions or comments?

1-800-SHOPRITE

## Package/Label Principal Display Panel

Compare to: Active Ingredients in Vicks® NyQuil®

NI-CALM

PAIN RELIEVER/FEVER REDUCER

Acetaminophen – Aches/Fever

COUGH SUPPRESSANT

Dextromethorphan HBr – Cough

ANTIHISTAMINE

Doxylamine Succinate – Sneezing/Runny Nose

Relieves major flu symptoms so you can rest

Multi-symptom – 10% ALCOHOL

Cherry Flavor

12 FL OZ (355 mL)

**Drug Facts** (continued)

- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks

**Stop use and ask a doctor if** ■ redness or swelling is present  
 ■ pain or cough gets worse or lasts more than 7 days  
 ■ fever gets worse or lasts more than 3 days ■ new symptoms occur  
 ■ cough comes back or occurs with rash or headache that lasts.  
 These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ take only as directed – see Overdose warning  
 ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs  
 adults & children 12 yrs & over 30 mL every 6 hrs  
 children 4 to under 12 yrs ask a doctor  
 children under 4 yrs do not use

**Other information** ■ each 30 mL contains: sodium 39 mg  
 ■ store at 20-25°C (68-77°F)

**Inactive ingredients** alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

**Questions or comments?** 1-800-SHOPRITE

\*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil®.

Distributed By: Wakefern Food Corp.  
 5000 Riverside Drive  
 Keasbey, NJ 08832  
 ©2016

**PARENTS:**  
 Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)



Compare to: Active Ingredients in Vicks® NyQuil®



ShopRite®

# NI-CALM

## PAIN RELIEVER/FEVER REDUCER

Acetaminophen....Aches/Fever

## COUGH SUPPRESSANT

Dextromethorphan HBr....Cough

## ANTIHISTAMINE

Doxylamine Succinate....Sneezing/Runny Nose

Relieves major flu symptoms so you can rest  
 Multi-symptom • 10% ALCOHOL

Cherry Flavor

12 FL OZ (355 mL)

**Drug Facts** **DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine

**Uses** temporarily relieves common cold/flu symptoms: ■ fever  
 ■ cough due to minor throat and bronchial irritation ■ sore throat  
 ■ minor aches and pains ■ runny nose and sneezing ■ headache

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours  
 ■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma  
 ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem such as emphysema or chronic bronchitis ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough as occurs with smoking, asthma, or emphysema ■ a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers  
 ■ taking the blood thinning drug warfarin

**When using this product**  
 ■ excitability may occur, especially in children  
 ■ marked drowsiness may occur

45940 6B F4

## SHOPRITE NI CALM

acetaminophen, dextromethorphan hbr, doxylamine succinate solution

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41190-459
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	

<b>SACCHARIN SODIUM</b> (UNI: SB8ZUX40TY)	
<b>SODIUM CITRATE</b> (UNI: 1Q73Q2JULR)	

### Product Characteristics

<b>Color</b>	RED (Clear/Dark Red)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHEERRY (Menthol Aroma)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41190-459-34	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/17/2013	
2	NDC:41190-459-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/07/2013	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	06/17/2013	

**Labeler** - Wakefern Food Corporation (069722418)

Revised: 12/2018

Wakefern Food Corporation