HEMORRHOID MASTER- lidocaine, phenylephrine hcl ointment Clinical Resolution Laboratory, Inc.

Active Ingredients

Lidocaine 4.00%

Phenylephrine HCI 0.25%

Directions

- Adults: When practical, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product. Children under 12 years of age: consult a doctor.
- For External Use: Apply externally to the affected area up to 4 times daily.
- For Intrarectal use: Attach applicator to tube. Lubricate applicator well, then gently insert applicator into the rectum. Apply to the affected area up to 4 times daily.

Warnings:For external and/ or intrarectal use only.

- If condition worsens or does not improve within 7 days, consult a doctor.
- Do not exceed the recommended daily dosage unless directed by a doctor.
- In case of bleeding, consult a doctor promptly.
- Certain persons can develop allergic reactions to ingredients in this product. If the symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase, discontinue use and consult a doctor.
- Do not use this product if you have heart disease, high blood pressure, thyroid disease, diabetes, or difficulty in urination due to enlargement of the prostate gland unless directed by a doctor.
- Ask a doctor or pharmacist before use if you are presently taking a prescription drug for high blood pressure or depression.
- Do not use this product with an applicator if the introduction of the applicator into the rectum causes additional pain. Consult a doctor promptly.

Uses

- For the temporary relief of local and anorectal itching and discomfort associated with anorectal disorders and anorectal inflammation.
- For the temporary relief of pain, burning and soreness.
- Temporarily shrinks hemorrhoidal tissue.
- Temporarily reduces the swelling associated with irritation in hemorrhoids and other anorectal disorders.
- Adults: When practical, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product. Children under 12 years of age: consult a doctor.
- For External Use: Apply externally to the affected area up to 4 times daily.

■ For Intrarectal use: Attach applicator to tube. Lubricate applicator well, then gently

insert applicator into the rectum. Apply to the affected area up to 4 times daily.

Inactive Ingredients

Aloe Barbadensis Leaf Juice, Ascorbic Acid, Ascorbyl Palmitate, Caprylyl Glycol, Carthamus Tinctorius (Safflower) Seed Oil, Cetearyl Alcohol, Chamomilla Recutita (Matricaria) Flower Extract, Cholecalciferol, Cholesterol, Diisopropyl Sebacate, Dimethyl Isosorbide, Dodecane, Ethoxydiglycol, Ethylhexylglycerin, Glycerin, Hexylene Glycol, Hydrocortisone, Isododecane, Isopropyl Myristate, Lecithin, Mentha Piperita (Peppermint) Oil, Microcrystalline Wax, Mineral Oil, Octyldodecanol, Panax Ginseng Root Extract, PEG-8 Dimethicone, Petrolatum, Phenoxyethanol, Phospholipids, Polyethylene, Propylene Glycol, Punica Granatum Extract, Pyridoxine HCl, Retinyl Palmitate, Silica, Sodium Propoxyhydroxypropyl Thiosulfate Silica, Stearic Acid, Tocopheryl Acetate, Triethoxycaprylylsilane, Water, Zea Mays (Corn) Oil, Zinc Oxide

Purpose

Local Anesthetic

Vasoconstrictor

Other Information

- Keep away from direct sunlight or heat.
- Store at room temperature 15°-30°C (59°-86°F).
- Keep out of reach of children.
- In case of accidental overdose or ingestion, call a doctor or poison control center immediately.
- Do not use this product if seal is broken or missing.

Ebanel

4% LIDOCAINE & 0.25% PHENYLEPHRINE HCI

LIPOSOMAL HEMORRHOID MASTER



- For the Temporary Relief of Pain, Burning, and Soreness
- Temporarily Shrinks Hemorrhoidal Tissue

1.6 oz e 45 a **HEMORRHOIDAL OINTMENT**

- Temporarily Reduces the Swelling Associated with Irritation in Hemorrhoids

DRUG FACTS

Active Ingredients
Lidocaine 4.00%.....

Phenylephrine HCI 0.25%

Local Anesthetic

Uses: To rule temporary relief of local and anorectal itching and discomfort associated with anorectal disorders and anorectal inflammation. For the temporary relief of pain, burning and sorene Temporarily shrinks hemorrhoidal tissue. Temporarily reduces the swelling associated with irritation in hemorrhoids and other anorectal disorders.

Warnings: For external and /or intrarectal use only.

If condition worsens or does not improve within 7 days, consult a doctor. Do not exceed the recommended daily dosage unless directed by a doctor. In case of bleeding, consult a doctor promptly. Certain persons can develop allergic reactions to ingredients in this product. If the symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase, discontinue use and consult a doctor. Do not use this product if you have heard disease, high blood pressure, thyroid disease, diabetes, or difficulty in urination due to enlargement of the prostate gland unless directed by a doctor. Ask a doctor or pharmactis before use if you are presently taking a prescription drug for high blood pressure or depression. Do not use this product with an applicator if the introduction of the applicator into the rectum causes additional pain. Consult a doctor promptly.

Directions: Adults: When practical, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product. Children under 12 years of age: consult a doctor.

For Intrarectal Use: Attach applicator to tube. Lubricate applicator well, then gently insert applicator into the rectum. Apply to the affected area up to 4 times daily.

Other Intrarectal:

Keep away from direct sunlight or heat.

Store at room temperature 15°-30°C (59°-86°F).

Keep out of reach of children.

On other this product if seal is broken or missing.

In case of accidental overdose or ingestion, call a doctor poison control to the rectum.

Apply to the affected area up to 4 times daily.

Inactive Ingredients: Aloe Barbadensis Leaf Juice, Ascorbic Acid, Ascorbyl Palmitate, Caprylyl Glycol, Carthamus Tinctorius (Safflower) Seed Oil, Cetearyl Alcohol, Chamomilla Recutita (Matricaria) Flower Extract, Cholecalciferol, Cholesterol, Diisopropyl Sebacate, Dimethyl Isosorbide, Dodecane, Ethoxydiglycol, Ethylhexylglycerin, Glycerin, Hexylene Glycol, Hydrocortisone, Isododecane, Isopropyl Myristate, Lecithin, Mentha Piperita (Pepperpernit) Oil, Microcrystalline Wax, Mineral Oil, Octyldodecanol, Panax Ginseng Root Unrethicone, Petrolatum, Phenoxyethanol, Phospholipids, Polyethylene, Propylene Glycol, Punica Granatum Extract, Pyridoxine HCI, Retinyl Palmitate, Silica, Sodium Propoxyhoyloroxypropyl Thiosulfate Silica, Stearic Acid, Tocopheryl Acetate, Triethoxycaprylysilane, Water, Zea Mays (Corn) Oil, Zinc Oxide

Ebanel

4% LIDOCAINE & 0.25% PHENYLEPHRINE HCI

LIPOSOMAL HEMORRHOID MASTER



- For the Temporary Relief of Pain, Burning, and Soreness
- Temporarily Shrinks Hemorrhoidal Tissue

Temporarily Reduces the Swelling Associated with Irritation in Hemorrhoids

1.6 oz e 45 g

HEMORRHOIDAL OINTMENT

DRUG FACTS

Active Ingredients

Purpose Local Anesthetic

Uses: To rithe temporary relief of local and anorectal itching and discomfort associated with anorectal disorders and anorectal inflammation. To rithe temporary relief of pain, burning and soreness ■ Temporarily shrinks hemorrhoidal tissue. ■ Temporarily reduces the swelling associated with irritation in hemorrhoids and other anorectal disorders.

Warnings: For external and /or intrarectal use only.

If condition worsens or does not improve within 7 days, consult a doctor. Do not exceed the recommended daily dosage unless directed by a doctor. In case of bleeding, consult a doctor promptly. Certain persons can develop allergic reactions to ingredients in this product. If the symptom being treated does not subside or if redness, firstiation, swelling, pain, or other resymptoms develop or increase, high blood pressure, thyroid disease, diabetes, or difficulty in urination due to enlargement of the prostate gland unless directed by a doctor.

Ask a doctor or pharmacist before use if you are presently taking a prescription drug for high blood pressure or depression. Do not use this product with an applicator if the introduction of the applicator into the rectum causes additional pain. Consult a doctor promptly.

Directions: Adults: When practical, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product. Children under 12 years of age: consult a doctor.

For External Use: Apply externally to the affected area up to 4 times daily.

For Intrarectal Use: Attach applicator to tube. Lubricate applicator well, then gently insert applicator into the rectum. Apply to the affected area up to 4 times daily.

Inactive Ingredients: Aloe Barbadensis Leaf Juice, Ascorbic Acid, Ascorbyl Palmitate, Caprylyl Glycol, Carthamus Tinctorius (Safflower) Seed Oil, Cetearyl Alcohol, Chamomilla Recutita (Matricaria) Flower Extract, Cholecalciferol, Cholesterol, Diisopropyl Sebacate, Dimethyl Isosorbide, Dodecane, Ethoxydiglycol, Ethylhexylglycerin, Glycerin, Hexylene Glycol, Hydrocortisone, Isododecane, Isopropyl Myristate, Lecithin, Mentha Piperita (Peppermin) Oil, Microcrystaline Wax, Mineral Oil, Octyldodecanol, Panax Ginseng Root Inmethicone, Petrolatum, Phenoxyethanol, Phospholipids, Polyethylene, Propylene Glycol, Punica Granatum Extract, Pyridoxine HCl, Retinyl Palmitate, Silica, Sodium Propoxyhrdyroxypropyl Thiosulfate Silica, Stearic Acid, Tocopheryl Acetate, Triethoxycaprylylsilane, Water, Zea Mays (Com) Oil, Zinc Oxide

HEMORRHOID MASTER

lidocaine, phenylephrine hcl ointment

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63742-045	
Route of Administration	TOPICAL			

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	4 mg in 1 g		
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 1 g		

Ingredient Name	Strength
ETHYLHEXYLGLYCERIN (UNII: 147D247K3P)	
GLYCERIN (UNII: PDC6A3C0OX)	
PETROLATUM (UNII: 4T6H12BN9U)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
MINERAL OIL (UNII: T5L8T28FGP)	
ALOE BARBADENSIS LEAF JUICE (UNII: RUE8E6T4NB)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
HYDROCORTISONE (UNII: W4X0X7BPJ)	
WATER (UNII: 059QF0KO0R)	
ZEA MAYS (CORN) OIL (UNII: 8470G57WFM)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
MENTHA PIPERITA (PEPPERMINT) OIL (UNII: AV092KU4JH)	
PANAX GINSENG ROOT (UNII: CUQ3A77YXI)	
SILICA (UNII: ETJ7Z6XBU4)	
CHOLECALCIFEROL, (5E)- (UNII: W28SLM7V6D)	
DIMETHYL ISOSORBIDE (UNII: SA6A6V432S)	
DODECANE (UNII: 11A386X1QH)	
ETHOXYDIGLYCOL (UNII: A1A1I8X02B)	
HEXYLENE GLYCOL (UNII: KEH0A3F75J)	
PEG-8 DIMETHICONE (UNII: GIA7T7640D)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
EGG PHOSPHOLIPIDS (UNII: 1Z74184RGV)	
POLYETHYLENE (UNII: UG00KM4WR7)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
RETINYL PALMITATE (UNII: 1D1K0N0VVC)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
ZINC OXIDE (UNII: SOI2LOH54Z)	
ISODODECANE (UNII: A8289P68Y2)	
ASCORBYL PALMITATE (UNII: QN83US2B0N)	

OCTYLDODECANOL (UNII: 461N1O614Y)	
POMEGRANATE (UNII: 56687D1Z4D)	
PYRIDOXINE HCL (UNII: 68Y4CF58BV)	
SODIUM PROPOXYHYDROXYPROPYL THIOSULFATE SILICA (UNII: 208G222332)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
.ALPHATOCOPHEROL ACETATE, D- (UNII: A7E6112E4N)	
CAPRYLYL GLYCOL (UNII: 00YIU5438U)	
CHAMOMILE (UNII: FGL3685T2X)	
CARTHAMUS TINCTORIUS (SAFFLOWER) SEED OIL (UNII: 65UEH262IS)	
CETEARYL ALCOHOL (UNII: 2DMT128M1S)	
CHOLESTEROL (UNII: 97C5T2UQ7J)	
DIISOPROPYL SEBACATE (UNII: J8T3X564IH)	

Product Characteristics			
Color	white (Yellowish White)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
		45 g in 1 TUBE, WTH APPLICATOR; Type 0: Not a Combination Product	11/01/2019	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M015	11/01/2019	

Labeler - Clinical Resolution Laboratory, Inc. (825047942)

Registrant - Clinical Resolution Laboratory, Inc. (825047942)

Establishment				
Name	Address	ID/FEI	Business Operations	
Clinical Resolution Laboratory, Inc.		825047942	manufacture(63742-045)	

Revised: 6/2025 Clinical Resolution Laboratory, Inc.