


CHLORTHALIDONE- chlorthalidone tablet

Nivagen Pharmaceuticals, INC.

DESCRIPTION

Chlorthalidone is an oral antihypertensive/diuretic. It is a monosulfamyl diuretic that differs chemically from thiazide diuretics in that a double-ring system is incorporated in its structure. It is 2-chloro-5-(1-hydroxy-3-oxo-1-isoindolinyl) benzenesulfonamide with the following structural formula:



Molecular Formula: C₁₄H₁₁ClN₂O₄S

Molecular weight: 338.776

Chlorthalidone, USP is practically insoluble in water, in ether, and in chloroform; soluble in methanol; slightly soluble in alcohol.

Chlorthalidone tablets are available containing either 25 mg or 50 mg of chlorthalidone USP and the following inactive ingredients: microcrystalline cellulose, partially pregelatinized maize starch, sodium starch glycolate, colloidal silicon dioxide, magnesium stearate, ingredients of aquadry blend yellow like iron oxide yellow, lactose monohydrate, Ferrosoferric Oxide for 25 mg and ingredients of aquadry blend green like D & C Yellow #10 aluminum lake, FD & C blue #1/ Brilliant Blue FCF Aluminum Lake for 50 mg.

CLINICAL PHARMACOLOGY

Chlorthalidone is an oral diuretic with prolonged action (48–72 hours) and low toxicity. The major portion of the drug is excreted unchanged by the kidneys. The diuretic effect of the drug occurs in approximately 2.6 hours and continues for up to 72 hours. The mean half-life following a 50 to 200 mg dose is 40 hours. In the first order of absorption, the elimination half-life is 53 hours following a 50 mg dose, and 60 hours following a 100 mg dose. Approximately 75 percent of the drug is bound to plasma proteins, 58 percent of the drug being bound to albumin. This is caused by an increased affinity of the drug to erythrocyte carbonic anhydrase. Nonrenal routes of elimination have yet to be clarified. Data are not available regarding percentage of dose as unchanged drug and metabolites, concentration of the drug in body fluids, degree of uptake by a particular organ or in the fetus, or passage across the blood-brain barrier.

The drug produces copious diuresis with greatly increased excretion of sodium and chloride. At maximal therapeutic dosage, chlorthalidone is approximately equal in its diuretic effect to comparable maximal therapeutic doses of benzothiadiazine diuretics. The site of action appears to be the cortical diluting segment of the ascending limb of Henle's loop of the nephron.

INDICATIONS AND USAGE

Diuretics such as chlorthalidone are indicated in the management of hypertension either alone or in combination with other antihypertensive drugs.

Chlorthalidone is indicated as adjunctive therapy in edema associated with congestive heart failure, hepatic cirrhosis, and corticosteroid and estrogen therapy.

Chlorthalidone has also been found useful in edema due to various forms of renal dysfunction, such as nephrotic syndrome, acute glomerulonephritis, and chronic renal failure.

Usage in Pregnancy

The routine use of diuretics in an otherwise healthy woman is inappropriate and exposes mother and fetus to unnecessary hazard. Diuretics do not prevent development of toxemia of pregnancy, and there is no satisfactory evidence that they are useful in the treatment of developed toxemia. Edema during pregnancy may arise from pathologic causes or from the physiologic and mechanical consequences of pregnancy.

Chlorthalidone is indicated in pregnancy when edema is due to pathologic causes, just as it is in the absence of pregnancy (however, see PRECAUTIONS, below). Dependent edema in pregnancy, resulting from restriction of venous return by the expanded uterus, is properly treated through elevation of the lower extremities and use of support hose; use of diuretics to lower intravascular volume in this case is illogical and unnecessary. There is hypervolemia during normal pregnancy that is harmful to neither the fetus nor the mother (in the absence of cardiovascular disease), but that is associated with edema, including generalized edema, in the majority of pregnant women. If this edema produces discomfort, increased recumbency will often provide relief. In rare instances, this edema may cause extreme discomfort that is not relieved by rest. In these cases, a short course of diuretics may provide relief and be appropriate.

CONTRAINDICATIONS

Anuria.

Known hypersensitivity to chlorthalidone or other sulfonamide-derived drugs.

WARNINGS

Chlorthalidone should be used with caution in severe renal disease. In patients with renal disease, chlorthalidone or related drugs may precipitate azotemia. Cumulative effects of the drug may develop in patients with impaired renal function.

Chlorthalidone should be used with caution in patients with impaired hepatic function or progressive liver disease, since minor alterations of fluid and electrolyte balance may

precipitate hepatic coma.

Sensitivity reactions may occur in patients with a history of allergy or bronchial asthma.

The possibility of exacerbation or activation of systemic lupus erythematosus has been reported with thiazide diuretics, which are structurally related to chlorthalidone.

However, systemic lupus erythematosus has not been reported following chlorthalidone administration.

PRECAUTIONS

General

Hypokalemia may develop with chlorthalidone as with any other diuretic, especially with brisk diuresis when severe cirrhosis is present or during concomitant use of corticosteroids or ACTH.

Interference with adequate oral electrolyte intake will also contribute to hypokalemia. Digitalis therapy may exaggerate metabolic effects of hypokalemia especially with reference to myocardial activity.

Any chloride deficit is generally mild and usually does not require specific treatment except under extraordinary circumstances (as in liver disease or renal disease). Dilutional hyponatremia may occur in edematous patients in hot weather, appropriate therapy is water restriction, rather than administration of salt except in rare instances when the hyponatremia is life threatening. In actual salt depletion, appropriate replacement is the therapy of choice.

Hyperuricemia may occur or frank gout may be precipitated in certain patients receiving chlorthalidone.

Thiazide-like diuretics have been shown to increase the urinary excretion of magnesium; this may result in hypomagnesemia.

The antihypertensive effects of the drug may be enhanced in the post-sympathectomy patient.

If progressive renal impairment becomes evident, as indicated by a rising nonprotein nitrogen or blood urea nitrogen, a careful reappraisal of therapy is necessary with consideration given to withholding or discontinuing diuretic therapy.

Calcium excretion is decreased by thiazide-like drugs. Pathological changes in the parathyroid gland with hypercalcemia and hypophosphatemia have been observed in few patients on thiazide therapy. The common complications of hyperparathyroidism such as renal lithiasis, bone resorption and peptic ulceration have not been seen.

Information for Patients

Patients should inform their physician if they have: (1) had an allergic reaction to chlorthalidone or other diuretics or have asthma, (2) kidney disease, (3) liver disease, (4) gout, (5) systemic lupus erythematosus, or (6) been taking other drugs such as cortisone, digitalis, lithium carbonate, or drugs for diabetes.

Patients should be cautioned to contact their physician if they experience any of the following symptoms of potassium loss: excess thirst, tiredness, drowsiness, restlessness, muscle pains or cramps, nausea, vomiting, or increased heart rate or

pulse.

Patients should also be cautioned that taking alcohol can increase the chance of dizziness occurring.

Laboratory Tests

Periodic determination of serum electrolytes to detect possible electrolyte imbalance should be performed at appropriate intervals.

All patients receiving chlorthalidone should be observed for clinical signs of fluid or electrolyte imbalance: namely, hyponatremia, hypochloremic alkalosis, and hypokalemia. Serum and urine electrolyte determinations are particularly important when the patient is vomiting excessively or receiving parenteral fluids.

Drug Interactions

Chlorthalidone may add to or potentiate the action of other antihypertensive drugs. Potentiation occurs with ganglionic peripheral adrenergic blocking drugs.

Medication such as digitalis may also influence serum electrolytes. Warning signs, irrespective of cause, are: dryness of mouth, thirst, weakness, lethargy, drowsiness, restlessness, muscle pains or cramps, muscular fatigue, hypotension, oliguria, tachycardia, and gastrointestinal disturbances such as nausea and vomiting.

Insulin requirements in diabetic patients may be increased, decreased, or unchanged. Higher dosage of oral hypoglycemic agents may be required. Latent diabetes mellitus may become manifest during chlorthalidone administration.

Chlorthalidone and related drugs may increase the responsiveness to tubocurarine.

Chlorthalidone and related drugs may decrease arterial responsiveness to norepinephrine. This diminution is not sufficient to preclude effectiveness of the pressor agent for therapeutic use.

Drug /Laboratory Test Interactions

Chlorthalidone and related drugs may decrease serum PBI levels without signs of thyroid disturbance.

Carcinogenesis, Mutagenesis, Impairment of Fertility

No information is available.

Pregnancy

Teratogenic Effects. Pregnancy Category B

Reproduction studies have been performed in the rat and the rabbit at doses up to 420 times the human dose and have revealed no evidence of harm to the fetus due to chlorthalidone. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nonteratogenic Effects

Thiazides cross the placental barrier and appear in cord blood. The use of chlorthalidone and related drugs in pregnant women requires that the anticipated benefits of the drug be weighed against possible hazards to the fetus. These hazards include fetal or neonatal jaundice, thrombocytopenia, and possibly other adverse reactions that have occurred in the adult.

Nursing Mothers

Thiazides are excreted in human milk. Because of the potential for serious adverse reactions in nursing infants from chlorthalidone, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

Safety and effectiveness in children have not been established.

Geriatric Use

Clinical studies of the 15 mg chlorthalidone tablets did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience with 15 mg chlorthalidone tablets has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

ADVERSE REACTIONS

The following adverse reactions have been observed, but there is not enough systematic collection of data to support an estimate of their frequency.

Gastrointestinal System Reactions:anorexia, gastric irritation, nausea, vomiting, cramping, diarrhea, constipation, jaundice (intrahepatic cholestatic jaundice), pancreatitis.

Central Nervous System Reactions:dizziness, vertigo, paresthesias, headache, xanthopsia.

Hematologic Reactions:leukopenia, agranulocytosis, thrombocytopenia, aplastic anemia.

Dermatologic-Hypersensitivity Reactions:purpura, photosensitivity, rash, urticaria, necrotizing angitis (vasculitis, cutaneous vasculitis), Lyell's syndrome (toxic epidermal necrolysis).

Cardio vascular Reactions:orthostatic hypotension may occur and may be aggravated by alcohol, barbiturates, or narcotics.

Other Adverse Reactions:hyperglycemia, glycosuria, hyperuricemia, muscle spasm, weakness, restlessness, impotence.

Whenever adverse reactions are moderate or severe, chlorthalidone dosage should be reduced or therapy withdrawn.

OVERDOSAGE

Symptoms of acute overdosage include nausea, weakness, dizziness, and disturbances of electrolyte balance. The oral LD₅₀ of the drug in the mouse and the rat is more than 25,000 mg/kg body weight. The minimum lethal dose (MLD) in humans has not been established. There is no specific antidote, but gastric lavage is recommended, followed by supportive treatment. Where necessary, this may include intravenous dextrose-saline with potassium, administered with caution.

DOSAGE AND ADMINISTRATION

Therapy should be initiated with the lowest possible dose, then titrated according to individual patient response. A single dose given in the morning with food is recommended; divided daily doses are unnecessary.

Hypertension

Initiation

Therapy, in most patients, should be initiated with a single daily dose of 15 mg. If the response is insufficient after a suitable trial, the dosage may be increased to a single daily dose of 25 mg. If additional control is required, the dosage of chlorthalidone may be increased to 30-50 mg once daily or 100 mg once daily. A second antihypertensive drug (step 2 therapy) may be added, as necessary. Dosage above 100 mg daily usually does not increase effectiveness. Increases in serum uric acid and decreases in serum potassium are dose-related over the 15 to 100 mg/day range.

Maintenance

Maintenance doses may be lower than initial doses and should be adjusted according to individual patient response. Effectiveness is well sustained during continued use.

Edema

Initiation

Adults, initially 30 to 100 mg daily, or 100 mg on alternate days. Some patients may require 90 to 200 mg at these intervals or up to 200 mg daily. Dosages above this level, however, do not usually produce a greater response.

Maintenance

Maintenance doses may often be lower than initial doses and should be adjusted according to individual patient response. Effectiveness is well sustained during continued use.

HOW SUPPLIED

Chlorthalidone Tablets, USP are available containing 25 mg or 50 mg of Chlorthalidone, USP.

25 mg Tablets are Light yellow colour, round, unscored tablet, debossed with "N" on one side and plain on other side. 25 mg Tablets are supplied as follows:

Bottles of 100 tablets NDC # 75834-109-01

Bottles of 1000 tablets NDC # 75834-109-10

50 mg Tablets are Light green colour, round tablet, debossed with "N" and score line on one side and plain on other side. 50 mg Tablets are supplied as follows:

Bottles of 100 tablets NDC # 75834-110-01

Bottles of 1000 tablets NDC # 75834-110-10

Store at 20° to 25°C (68° to 77°F). [See USP for Controlled Room Temperature.]

Protect from light.

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.

ANIMAL PHARMACOLOGY

Biochemical studies in animals have suggested reasons for the prolonged effect of chlorthalidone. Absorption from the gastrointestinal tract is slow due to its low solubility. After passage to the liver, some of the drug enters the general circulation, while some is excreted in the bile, to be reabsorbed later. In the general circulation, it is distributed widely to the tissues, but is taken up in highest concentrations by the kidneys, where amounts have been found 72 hours after ingestion, long after it has disappeared from other tissues. The drug is excreted unchanged in the urine.

Manufactured for:

Nivagen Pharmaceuticals, Inc.
Sacramento, CA 95827 USA
Toll free number: 1-877-977-0687

Manufactured by:

Umedica Laboratories Pvt. Ltd.
Plot No.221 and 221/1,
GIDC, IInd Phase, Vapi,
Gujarat 396195, India (IND)

Manufactured by:

Square Pharmaceuticals Limited
Dhaka Unit, Kaliakoir,
Gazipur – 1750,
Bangladesh.

Revised July 2023, V-03

PACKAGE LABEL PRINCIPAL DISPLAY PANEL


Chlorthalidone Tablets, USP - 25 mg NDC - 75834-109-01 - 100 Tablets - Bottle Label (Umedica)

Rx only NDC 75834-109-01 Chlorthalidone Tablets, USP 25 mg 100 Tablets 	Each uncoated tablet contains Chlorthalidone USP.....25 mg Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure. Keep container tightly closed. Keep this and all medication out of the reach of children. Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.] Protect from light. Usual Dosage: See accompanying prescribing information. July 2023, V-03	 XXXXXX	Mfg. Lic. No. : G/25/491 Manufactured for: Nivagen Pharmaceuticals, Inc. Sacramento, CA 95827 USA Toll free number: 1-877-977-0687 Manufactured by: Umedica Laboratories Pvt. Ltd. Plot No. 221 and 221/1, GIDC, IInd Phase, Vapi, Gujarat 396195, INDIA (IND) Batch No. : Exp. Date :
---	--	---	---

Chlorthalidone Tablets, USP - 25 mg NDC - 75834-109-10 - 1000 Tablets - Bottle Label (Umedica)

Rx only NDC 75834-109-10 Chlorthalidone Tablets, USP 25 mg 1000 Tablets NIVAGEN	Each uncoated tablet contains Chlorthalidone USP.....25 mg Dispense in a tight, light-resistant container as defined in the USP using a continuous threads closure. Keep container tightly closed. Keep this and all medication out of the reach of children. Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.] Protect from light. Usual Dosage: See accompanying prescribing information. July 2023, V-03	 XXXXXX	Mfg. Lic. No. : G/25/491 Manufactured for: Nivagen Pharmaceuticals, Inc. Sacramento, CA 95827 USA Toll free number: 1-877-977-0687 Manufactured by: Umedica Laboratories Pvt. Ltd. Plot No. 221 and 221/1, GIDC, IInd Phase, Vapi, Gujarat 396195, INDIA (IND)
---	--	---	---

Chlorthalidone Tablets, USP - 50 mg NDC - 75834-110-01 - 100 Tablets - Bottle Label (Umedica)

Rx only NDC 75834-110-01 Chlorthalidone Tablets, USP 50 mg 100 Tablets 	Each uncoated tablet contains Chlorthalidone USP.....50 mg Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure. Keep container tightly closed. Keep this and all medication out of the reach of children. Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.] Protect from light. Usual Dosage: See accompanying prescribing information. July 2023, V-03	 XXXXXX	Mfg. Lic. No. : G/25/491 Manufactured for: Nivagen Pharmaceuticals, Inc. Sacramento, CA 95827 USA Toll free number: 1-877-977-0687 Manufactured by: Umedica Laboratories Pvt. Ltd. Plot No. 221 and 221/1, GIDC, IInd Phase, Vapi, Gujarat 396195, INDIA (IND) Batch No. : UNVARNISHED ZONE Exp. Date : 50 mm × 18 mm
--	---	---	--

Chlorthalidone Tablets, USP - 50 mg NDC - 75834-110-10 - 1000 Tablets - Bottle Label (Umedica)

	Each uncoated tablet contains Chlorthalidone USP.....50 mg Dispense in a tight, light-resistant container as defined in the USP using a continuous threads closure. Keep container tightly closed. Keep this and all medication out of the reach of children. Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.] Protect from light. Usual Dosage: See accompanying prescribing information. July 2023, V-03	 XXXXXX	Mfg. Lic. No. : G/25/491
			Manufactured for: Nivagen Pharmaceuticals, Inc. Sacramento, CA 95827 USA Toll free number: 1-877-977-0687 Manufactured by: Umedica Laboratories Pvt. Ltd. Plot No. 221 and 221/1, GIDC, IInd Phase, Vapi, Gujarat 396195, INDIA (IND)
			Batch No. : UNVARNISHED ZONE Exp. Date : 62 mm × 20 mm

Chlorthalidone Tablets, USP - 25 mg NDC - 75834-109-01 - 100 Tablets - Bottle Label (Square Pharma)

	Each uncoated tablet contains Chlorthalidone USP.....25 mg Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure. Keep container tightly closed. Keep this and all medication out of the reach of children. Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.] Protect from light. Usual Dosage: See accompanying prescribing information. July 2023, V-03	 10044	Mfg. Lic. No. : 460
			Manufactured for: Nivagen Pharmaceuticals, Inc. Sacramento, CA 95827 USA Toll free number: 1-877-977-0687 Manufactured by: Square Pharmaceuticals LTD. Dhaka Unit, Kaliakoir, Gazipur-1750 Bangladesh
			UNVARNISHED ZONE 50 mm × 18 mm

Chlorthalidone Tablets, USP - 25 mg NDC - 75834-109-10 - 1000 Tablets - Bottle Label (Square Pharma)

	Each uncoated tablet contains Chlorthalidone USP.....25 mg Dispense in a tight, light-resistant container as defined in the USP using a continuous threads closure. Keep container tightly closed. Keep this and all medication out of the reach of children. Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.] Protect from light. Usual Dosage: See accompanying prescribing information. July 2023, V-03	 10045	Mfg. Lic. No. : 460
			Manufactured for: Nivagen Pharmaceuticals, Inc. Sacramento, CA 95827 USA Toll free number: 1-877-977-0687 Manufactured by: Square Pharmaceuticals LTD. Dhaka Unit, Kaliakoir, Gazipur-1750 Bangladesh
			UNVARNISHED ZONE 62 mm × 20 mm

Chlorthalidone Tablets, USP - 50 mg NDC - 75834-110-01 - 100 Tablets -

Rx only

NDC 75834-110-01

Chlorthalidone
Tablets, USP

50 mg

100 Tablets

NIVAGEN

Each uncoated tablet contains Chlorthalidone USP.....50 mg

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.

Keep container tightly closed.

Keep this and all medication out of the reach of children.

Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]

Protect from light.

Usual Dosage:
See accompanying prescribing information.

July 2023, V-03

10046

Mfg. Lic. No. : 460

Manufactured for: Nivagen Pharmaceuticals, Inc.
Sacramento, CA 95827 USA
Toll free number: 1-877-977-0687

Manufactured by: Square Pharmaceuticals LTD.
Dhaka Unit, Kaliakoir, Gazipur-1750
Bangladesh

UNVARNISHED ZONE

50 mm × 18 mm

2000026857

Rx only

NDC 75834-110-10

Chlorthalidone Tablets, USP

50 mg

1000 Tablets

NIVAGEN

Each uncoated tablet contains Chlorthalidone USP.....50 mg

Dispense in a tight, light-resistant container as defined in the USP using a continuous threads closure.

Keep container tightly closed.

Keep this and all medication out of the reach of children.

Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]

Protect from light.

Usual Dosage:
See accompanying prescribing information.

July 2023, V-03

3 75834 11010 3

10047

Mfg. Lic. No. : 460

Manufactured for: **Nivagen Pharmaceuticals, Inc.**
Sacramento, CA 95827 USA
Toll free number: 1-877-977-0687

Manufactured by: **Square Pharmaceuticals LTD.**
Dhaka Unit, Kaliakoir, Gazipur-1750
Bangladesh

UNVARNISHED ZONE
62 mm × 20 mm

Ingredient Name	Strength
-----------------	----------

CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
STARCH, CORN (UNII: O8232NY3SJ)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
Product Characteristics				
Color	green (Light Green)		Score	2 pieces
Shape	ROUND		Size	7mm
Flavor			Imprint Code	N
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75834-110-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/28/2018	
2	NDC:75834-110-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	05/28/2018	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA207222		05/28/2018	

CHLORTHALIDONE			
chlorthalidone tablet			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:75834-109
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
CHLORTHALIDONE (UNII: Q0MQD1073Q) (CHLORTHALIDONE - UNII:Q0MQD1073Q)		CHLORTHALIDONE	25 mg
Inactive Ingredients			
Ingredient Name			Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
STARCH, CORN (UNII: O8232NY3SI)			

SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

Product Characteristics

Color	yellow (Light Yellow)	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	N
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75834-109-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/28/2018	
2	NDC:75834-109-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	05/28/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA207222	05/28/2018	

Labeler - Nivagen Pharmaceuticals, INC. (052032418)

Registrant - Umedica Laboratories Pvt. Ltd. (650219934)

Establishment

Name	Address	ID/FEI	Business Operations
SQUARE PHARMACEUTICALS LIMITED, Dhaka unit		850366520	manufacture(75834-109, 75834-110) , analysis(75834-109, 75834-110) , pack(75834-109, 75834-110) , label(75834-110, 75834-110)

Establishment

Name	Address	ID/FEI	Business Operations
Umedica Laboratories Pvt. Ltd.		920635096	manufacture(75834-109, 75834-110) , analysis(75834-109, 75834-110) , pack(75834-109, 75834-110)