

PRILOSEC OTC- omeprazole magnesium tablet, delayed release
The Procter & Gamble Manufacturing Company

Prilosec OTC ®

Drug Facts

Active ingredient (in each tablet)

Omeprazole delayed-release tablet 20 mg (equivalent to 20.6 mg omeprazole magnesium)

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs ***2 or more*** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- Do not use if you are allergic to omeprazole.
- Omeprazole may cause severe skin reactions. Symptoms may include:
 - skin reddening • blisters • rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck, or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

taking a prescription drug.

Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- **swallow whole. Do not chew, crush, or suck tablets.**

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20 to 25°C (68 to 77°F) and protect from moisture

Inactive ingredients

acetyl tributyl citrate, colloidal silicon dioxide, corn starch, croscarmellose sodium,

hydroxypropyl cellulose, hypromellose 2910, magnesium stearate, methacrylic acid copolymer type c, microcrystalline cellulose, polyethylene glycol 400, polyethylene glycol 3350, polyethylene glycol 8000, polyvinyl alcohol, red iron oxide, sucrose, talc, titanium dioxide, triethyl citrate

Questions?

1-800-289-9181

Safety Feature - Do not use if tablet blister unit is open or torn.

**Dist. by Procter & Gamble,
Cincinnati, OH 45202**

PRINCIPAL DISPLAY PANEL - 42 Tablet Carton

NEW FORMULA

SAME ACTIVE INGREDIENT.

SEE CURRENT DRUG FACTS.

NDC 84126-300-42

Treats FREQUENT Heartburn! 24 HR

Prilosec OTC [®]

omeprazole delayed-release tablets

20 mg / acid reducer

Three 14-day courses of treatment

May take 1 to 4 days for full effect.

42 TABLETS

Treats FREQUENT Heartburn!

24
HR

Prilosec
OTC®

NEW FORMULA
SAME ACTIVE INGREDIENT.
SEE CURRENT DRUG FACTS.

NDC 84126-300-42

Treats FREQUENT Heartburn!

24
HR

Prilosec

omeprazole delayed-release tablets
20 mg / acid reducer **OTC®**

42 TABLETS

Three 14-day courses of treatment
May take 1 to 4 days for full effect

24
ACTUAL SIZE



Prilosec
OTC®

If you are not satisfied with Prilosec OTC®, simply return the UPC Code from this package and original sales receipt within 60 days of purchase for a full refund. For offer details, visit PrilosecOTCGuarantee.com.

P&G

www.pg.com
Patents: www.pg.com/patents

Safety Feature -
Do not use if
tablet blister
unit is open
or torn.

Dist. by Procter & Gamble,
Cincinnati, OH 45202

21259300



Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole delayed-release tablet 20 mg Acid reducer
(equivalent to 20.6 mg omeprazole magnesium)

Use • treats frequent heartburn (occurs 2 or more days a week)
• not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:
• Do not use if you are allergic to omeprazole.
• Omeprazole may cause severe skin reactions. Symptoms may include:
• skin reddening • blisters • rash
If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:
• trouble or pain swallowing food, vomiting with blood, or bloody or black stools
• heartburn with lightheadedness, sweating or dizziness
• chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
• frequent chest pain
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
• had heartburn over 3 months. This may be a sign of a more serious condition.
• frequent wheezing, particularly with heartburn
• unexplained weight loss • nausea or vomiting • stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
• your heartburn continues or worsens
• you need to take this product for more than 14 days
• you need to take more than 1 course of treatment every 4 months
• you get diarrhea • you develop a rash or joint pain

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

• for adults 18 years of age and older
• this product is to be used once a day (every 24 hours), every day for 14 days
• it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
• swallow 1 tablet with a glass of water before eating in the morning
• take every day for 14 days
• do not take more than 1 tablet a day
• do not use for more than 14 days unless directed by your doctor
• swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)
• you may repeat a 14-day course every 4 months
• do not take for more than 14 days or more often than every 4 months unless directed by a doctor
• children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

• read the directions and warnings before use
• keep the carton. It contains important information.

• store at 20° to 25°C (68° to 77°F) and protect from moisture

Inactive ingredients acetyl tributyl citrate, colloidal silicon dioxide, corn starch, croscarmellose sodium, hydroxypropyl cellulose, hypromellose 2910, magnesium stearate, methacrylic acid copolymer type c, microcrystalline cellulose, polyethylene glycol 400, polyethylene glycol 3350, polyethylene glycol 8000, polyvinyl alcohol, red iron oxide, sucrose, talc, titanium dioxide, triethyl citrate

Questions? 1-800-289-9181

PRILOSEC OTC

omeprazole magnesium tablet, delayed release							
Product Information							
Product Type		HUMAN OTC DRUG	Item Code (Source)	NDC:84126-300			
Route of Administration		ORAL					
Active Ingredient/Active Moiety							
Ingredient Name			Basis of Strength	Strength			
OMEPRAZOLE MAGNESIUM (UNII: 426QFE7XLK) (OMEPRAZOLE - UNII:KG60484QX9)			OMEPRAZOLE	20.6 mg			
Inactive Ingredients							
Ingredient Name				Strength			
ACETYL TRIBUTYL CITRATE (UNII: 0ZBX0N59RZ)							
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)							
STARCH, CORN (UNII: O8232NY3SJ)							
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)							
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)							
HYPROMELLOSE 2910 (15 MPA.S) (UNII: 36SFW2JZ0W)							
MAGNESIUM STEARATE (UNII: 70097M6I30)							
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)							
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)							
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)							
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)							
POLYVINYL ALCOHOL (UNII: 532B59J990)							
SUCROSE (UNII: C151H8M554)							
TALC (UNII: 7SEV7J4R1U)							
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)							
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)							
FERRIC OXIDE RED (UNII: 1K09F3G675)							
POLYETHYLENE GLYCOL 8000000 (UNII: 8EIY1IHX76)							
Product Characteristics							
Color	red	Score	score with uneven pieces				
Shape	OVAL	Size	13mm				
Flavor		Imprint Code	24				
Contains							
Packaging							
#	Item Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:84126-300-14	1 in 1 CARTON	09/01/2025				
1		14 in 1 BLISTER PACK; Type 0: Not a Combination					

1		Product		
2	NDC:84126-300-42	3 in 1 CARTON	09/01/2025	
2		14 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:84126-300-24	2 in 1 CARTON	09/01/2025	
3		14 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206582	09/01/2025	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Registrant - The Procter & Gamble Manufacturing Company (004238200)